

## ***Cranial Electrostimulation Therapy (CES)***

**Effective:** February 1, 2023

**Next Review:** November 2023

**Last Review:** December 2022

### **IMPORTANT REMINDER**

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

### **DESCRIPTION**

Cranial electrostimulation therapy (CES), also called cranial electrotherapy stimulation, involves passing small electrical impulses across the head, usually from electrodes placed on or near both ears.

### **MEDICAL POLICY CRITERIA**

Cranial electrostimulation therapy is considered **investigational** for all indications, including but not limited to treatment of:

- A. Alzheimer's disease
- B. Anxiety
- C. Apathy related to traumatic brain injury
- D. Chemical dependence / substance abuse
- E. Chronic pain related to spinal cord injury
- F. Cognitive dysfunction
- G. Depressive symptoms
- H. Fibromyalgia

- I. Headache
- J. Irritable Bowel Syndrome (IBS)
- K. Smoking cessation
- L. Sleep disturbances
- M. Stress related conditions
- N. Tinnitus
- O. Traumatic brain injury

*NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.*

## POLICY GUIDELINES

Some cranial electrostimulation therapy (CES) devices may also be FDA approved to apply electrical stimulation to peripheral nerves [e.g., transcutaneous electrical nerve stimulation (TENS)]. This policy addresses cranial electrical stimulation that targets the brain only; electrical stimulation of peripheral nerves for the treatment of pain or other indications is addressed in separate policies (see Medical Policy, see Cross References, DME-83 for an index of other electrical stimulation policies).

## CROSS REFERENCES

1. [External Trigeminal Nerve Stimulation for the Treatment of Attention Deficit Hyperactivity Disorder](#), Durable Medical Equipment, Policy No. 83.14

## BACKGROUND

Although the mechanism of action is not clearly understood, it is hypothesized that electrical currents emitted from CES may positively impact the limbic system, the reticular activating system and/or the hypothalamus, resetting the brain to improved homeostasis levels.<sup>[1]</sup>

CES is proposed for use in treating a variety of chronic conditions including, but not limited to stress, alcoholism and drug addiction, headache, cognitive dysfunction in head injured patients, psychiatric conditions, irritable bowel syndrome, reflex sympathetic dystrophy and multiple sclerosis. Because many of these indications require long-term therapy with medications which may be costly, CES has been proposed as a cost-effective, non-invasive alternative to standard treatment.

## REGULATORY STATUS

The U.S. Food and Drug Administration (FDA) has granted 510(k)\* approval for a number of cranial electrotherapy stimulators including, but not limited to the following:

- Alpha-Stim® Cs (Electromedical Products, Inc)
- BR-2 Biorest (Biorest, Inc)
- Biotron18 (Biotronics Corp)
- CES Ultra™ (Neuro-Fitness, LLC)
- Elexoma Medic (Redplane AG)
- FM 10/C (Johari Digital Healthcare, Ltd)

- HP-1 Healthpax or Nurtipax (Health Directions, Inc)
- LB-2000 (Life Balance Intl., Inc)
- LISS SBI202-B and SBI201-M (Medical Consultants Intl., Ltd)
- NET-2000 Microcurrent Stimulator (Auri-Stim Medical, Inc)
- NF-1 Mindpeace (NeuroFitness)
- NH 2002 (Life Balance Intl., Inc.)
- NTI-1000 (Neurotek, Inc)
- TESA-1 (Kalaco Scientific, Inc.)

\*Marketing clearance via the 510(k) process does not require data regarding clinical efficacy.

In June of 2019, the FDA granted De Novo approval of the IB-Stim (Innovative Health Solutions) to aid in the reduction of functional abdominal pain in patients 11-18 years of age with irritable bowel syndrome (IBS). The device subsequently was reclassified to a Class II device with the broader regulation name of “non-implanted nerve stimulator for functional abdominal pain relief”.<sup>[2]</sup>

## EVIDENCE SUMMARY

The principal outcomes associated with treatment of pain due to any cause may include: relief of pain, improved functional level, and return to work. Relief of pain is a subjective outcome that is typically associated with a placebo effect. Therefore, data from adequately powered, blinded, randomized controlled trials (RCT) are required to control for the placebo effect, determine its magnitude, and determine whether any treatment effect from an electrical stimulation device provides a significant advantage over the placebo. Treatment of mood disorders (anxiety, depression) and chemical dependency issues require the same level of evidence to ensure valid conclusions regarding superiority over placebo.

Treatment with an electrical stimulation device must also be evaluated in general groups of patients against the existing standard of care for the condition being treated. For example, in patients with pain symptoms, treatment with an electrical stimulation device should be compared to other forms of conservative therapy such as pain medications. In patients with mood disorders or chemical dependency issues, treatment must be compared with the standard of care: psychotherapy or behavioral therapy, respectively, with or without medication.

## SYSTEMATIC REVIEWS AND META ANALYSES

Ching (2022) published a Systematic Review (SR) with meta-analysis of Cranial Electrotherapy Stimulation (CES) in the treatment of anxiety with a secondary outcome of CES on depressive symptoms.<sup>[3]</sup> To be included, studies met the following eligibility criteria: randomized clinical trials (RCTs) investigating the effects of CES as monotherapy or combination with other treatment (e.g., biofeedback therapy and antidepressants) for management of anxiety symptoms; (ii) diagnosis of anxiety disorders which meet the criteria in DSM-IV, DSM-IV TR, DSM-V or ICD10; (iii) the definition of anxiety symptoms based on screening tool; (iv) a comparison between an intervention group and a control group (e.g., biofeedback therapy, antidepressants); (v) sufficient data for both the intervention group and the control group; (vi) articles written in English. Cranial Electrotherapy Stimulation significantly improved anxiety symptoms with moderate effect size (number of trials = 11, n =

692, Hedge's  $g = -0.625$ , 95% CIs =  $-0.952$  to  $-0.298$ ,  $p < 0.001$ ) compared to control/sham group. Additionally, CES significantly reduced depressive symptoms in patients with anxiety disorders (number of trials = 8,  $n = 552$ , Hedges'  $g = -0.648$ , 95% CIs =  $-1.062$  to  $-0.234$ ,  $p = 0.002$ ;) compared to control/sham group. Significant heterogeneity was noted for both outcomes. Majority of the studies used Alpha-stim products. Limitations included five out of 11 studies (45.4%) had an overall RoB2 ranking of some risk of bias concerns, studies used different stimulation protocols and the protocols were only used for acute treatment, limiting the ability to evaluate long term outcomes. The authors note that the placebo affect is a critical issue in these types of studies.

Price (2021) published a SR with meta-analyses of CES(Alpha-Stim) for the treatment of depression.<sup>[4]</sup> Two separate meta-analyses were conducted, one on data from RCTs (five studies,  $n = 242$ ) and another on data from non-randomized studies (12 studies,  $n = 1173$ ). To be considered as a RCT, trials were required to have subjects blinded, a sham versus active condition, use of valid and reliable measurement instruments, at a minimum, a pretest-posttest design (additional repeated measures were acceptable), and rated as "good" or "fair" in a scoring system in which scoring categories were 0–1 limitations (rating = good); 2–4 limitations (rating = fair); 5–9 limitations (rating = limited). Due to variation in reporting of results across the five RCTs, only the difference at posttest between groups was used in the calculation of the effect of Alpha-Stim CES on depression. The average (population) effect for the five RCTs was  $d = -0.69$  (i.e., the mean depression level at posttest for the active group was  $-0.69$  standard deviations lower than the mean depression level for the sham group), which is considered a medium effect in favor of the treatment group. Analysis of the nonrandomized study data revealed an average (population) effect of  $d = -0.43$ , which is a small effect size. Significant heterogeneity for the effect sizes was found for this analysis. Limitations in the available RCT data include the small samples of patients who met DSM-V criteria for major depressive disorder. The systematic review and meta-analyses were sponsored by Electromedical Products International, Inc. and several study authors are affiliated with the company.

In a 2020 SR with meta-analysis published by Yu, the effectiveness of noninvasive brain stimulation in the treatment of neuropathic pain in individuals with spinal cord injury was evaluated.<sup>[5]</sup> Although eleven studies of various noninvasive brain stimulation treatments were included, only one used CES as the intervention.<sup>[6]</sup> The pooled analysis found no significant effect of any of the treatment modalities on neuropathic pain reduction after spinal cord injury. While no beneficial effect over sham stimulation was found for depression scales, reduction in anxiety immediately following CES treatment was identified. The authors concluded that their findings do not support the routine use of noninvasive brain stimulation for neuropathic pain in individuals with spinal cord injury.

A 2018 SR prepared by Shekelle for the Department of Veterans Affairs Evidence-based Synthesis Program (ESP) synthesized evidence on CES for chronic pain, depression, anxiety, insomnia, and posttraumatic stress disorder (PTSD).<sup>[7]</sup> The authors identified 28 RCTs that met inclusion criteria. A meta-analysis could not be completed because there were too few studies of the same patient population and treatment protocol. The quality of all included RCTs was found to be low, and all had a high risk of bias. Therefore, although the results of the included RCTs indicated that CES may have a modest beneficial effect on symptoms of anxiety and depression in selected patients, the authors urged caution in interpreting the results.

A Cochrane SR and meta-analysis evaluated the use of CES as a non-invasive treatment for chronic pain, originally published in 2011 and updated in 2014 and again in 2018.<sup>[8-10]</sup> No differences were found in health outcomes when CES was compared with sham in the 11 studies that met the inclusion criteria. The review concluded that all available studies were at risk of bias, and that available data failed to suggest that CES provided a clear benefit over sham treatment.

Boldt (2014) evaluated non-pharmacological interventions for chronic pain in people with spinal cord injury in a Cochrane SR, including two trials that assessed the effects of transcranial direct current stimulation (tDCS), three trials that used repetitive transcranial magnetic stimulation (rTMS), and three studies that used CES.<sup>[11]</sup> In all of these trials sham controls were used. For the use of tDCS, the overall evidence for the effectiveness of tDCS in reducing chronic pain in spinal cord injury was scarce and inconclusive. For the use of rTMS, the data from the three studies was inconsistent regarding the treatments effectiveness in reducing chronic pain in this population. The two studies on CES had methodological limitations including selective reporting and imbalances in baseline characteristics between groups, and a third study was inconclusive.

A 2014 Cochrane SR by Kavirajan assessed the efficacy and safety of CES as a treatment of acute depression compared to sham or simulated CES treatment.<sup>[12]</sup> Authors searched for properly blinded randomized trials of CES in adults aged 18-75 with depressive disorder, however, no studies met inclusion criteria. The authors concluded, “(t)here are insufficient methodologically rigorous studies of CES in treatment of acute depression. There is a need for double-blind RCTs of CES in the treatment of acute depression.”

A 2009 Cochrane SR for treatment of apathy in traumatic brain injury found only one RCT which met inclusion criteria for the review.<sup>[13]</sup> However, the reviewers cautioned against making conclusions from this RCT due to the small study size (n=21).

## **RANDOMIZED CONTROLLED TRIALS**

A randomized controlled trial (RCT) by Kim (2021) evaluated CES as a treatment for anxiety in which nonclinical volunteers experiencing daily anxiety were randomly assigned to either active or sham groups<sup>[14]</sup>. Outcomes assessed after three weeks of 20 self-administered treatment sessions included the State-Trait Anxiety Inventory (STAI), Beck Depression Inventory (BDI), Wisconsin Card Sort Test (WCST), and resting-state electroencephalography (EEG). A significant improvement in state-anxiety was found in the active treatment group compared to sham, however, no significant difference in depression or WCST was found between groups. Increased EEG signal in specific brain regions in the active treatment group correlating to changes in state-anxiety were observed, the clinical utility of which is not clearly established. This study is limited by small sample size and no long-term outcomes.

Kovacic (2017) published the results of a RCT of the Neuro-Stim device for the treatment of abdominal pain-related functional gastrointestinal disorders in adolescents (aged 11-18 years).<sup>[15]</sup> Patients were randomly assigned (1:1) to an active stimulation or sham (no electrical charge) for four weeks and patient-reported worst abdominal pain and composite pain scores (Pain Frequency-Severity-Duration [PFSD] scale) were assessed as the primary outcomes. A total of 57 patients in the active treatment group and 47 patients in the sham group were included in the primary analysis. A greater reduction in worst pain after three weeks of treatment was found in the treatment compared to the sham group (treatment: median score 5.0 [IQR 4.0–7.0]; sham: 7.0 [5.0–9.0]; least square means estimate of change in worse pain

2.15 [95% CI 1.37–2.93],  $p < 0.0001$ ). PFSD composite scores also decreased significantly in the treatment group (from 24.5 [IQR 16.8–33.3] to 8.4 [3.2–16.2]) compared with sham (from 22.8 [IQR 8.4–38.2] to 15.2 [4.4–36.8]) with a mean decrease of 11.48 (95% CI 6.63–16.32;  $p < 0.0001$ ) after three weeks. These differences persisted into follow-up (median follow-up 9.2 weeks [IQR 6.4–13.4]). There were no serious adverse events reported.

A subset analysis of the Kovacic RCT was published by Krasaelap (2020) which evaluated outcomes for patients with Irritable Bowel Syndrome (IBS), specifically.<sup>[16]</sup> This cohort consisted of patients who received active stimulation ( $n=27$ ) or sham stimulation ( $n=23$ ) for four weeks. The number of patients with a reduction of 30% or more in worst abdominal pain severity after three weeks was considered the primary outcome. Reduction in composite abdominal pain severity score, reduction in usual abdominal pain severity, and improvement in global symptom based on a symptom response scale after three weeks were evaluated as secondary outcomes. The primary outcome was found in 59% of patients who received the active stimulation and 26% of patients who received sham stimulation ( $p=0.024$ ). Composite pain and usual pain median scores were reduced in the active stimulation compared to the sham group ( $p=0.026$ ,  $0.029$ , respectively). A symptom response scale score of two or more was observed in 82% of patients who received active treatment compared to 26% of patients in the sham group ( $p \leq 0.001$ ). The authors reported no significant side effects. This study was limited by small sample size, lack of long-term follow-up, and specific endpoints not being defined *a priori*. The stimulation devices were provided by Innovative Health Solutions.

Roh and Wi-Young (2017) published a RCT that evaluated how CES effects symptoms of depression and anxiety, by evaluating behaviors and certain hormones.<sup>[17]</sup> Fifty postmenopausal women received active CES ( $n=25$ ) or a sham treatment ( $n=25$ ). The active group received 20 minutes of CES three times a week for eight weeks. Cortisol, adrenocorticotrophic hormone (ACTH), brain derived neurotrophic factor (BDNF), and nerve growth factor (NGF) levels were evaluated prior to the treatments and after the eight-week sessions. No differences in the levels were found. The CES group had less depression and tension-anxiety, but no changes were seen for anger-hostility, vigor-anxiety, fatigue-inertia, and confusion-bewilderment. This study had methodological limitations including small sample size and lack of long-term follow-up.

A number of RCTs explored the efficacy of CES for a variety of conditions not addressed in the Cochrane SRs noted above, including Alzheimer's disease, smoking cessation, anxiety in patients receiving dental care, preoperative anxiety, chemical dependence, sleep disturbances, fibromyalgia, constipation, dysfunctional gait, preoperative blood pressure, pediatric tic disorders, and tinnitus.<sup>[18-30]</sup> In addition, several RCTs not included in the reviews above were also identified.<sup>[31, 32]</sup> Overall, data from these studies were unreliable due to a variety of limitations, including small study populations,<sup>[18-24, 27, 33, 34]</sup> short follow-up of study subjects,<sup>[18-24, 26, 33-35]</sup> confounding use of co-therapies such as fibromyalgia medications<sup>[23]</sup> and antidepressants<sup>[31]</sup>, weak or unclear randomization methods,<sup>[18, 21, 23, 24]</sup> and the use of flawed data analysis methodologies such as deleting a subset of patients based on their diagnosis after they had been randomized and treated<sup>[21]</sup>, rendered the study findings unreliable.

Overall, the RCTs did not adequately explain the clinical significance of the changes observed in their outcomes of interest.<sup>[36]</sup> The treatment parameters used in the studies varied in their frequency, intensity, duration of individual CES sessions, as well as the overall treatment duration. Only two studies evaluated how changes in treatment parameters influenced the

same outcome of interest. They did not find a significant difference between the two, but these studies were subject to other major design flaws.<sup>[18, 19]</sup>

## PRACTICE GUIDELINE SUMMARY

There are no evidence-based clinical practice guidelines that recommend the use of cranial electrical stimulation devices for the treatment of pain or any other indication.

## SUMMARY

There is not enough research to show that cranial electrostimulation therapy (CES) improves health outcomes for people with pain or any other condition. In addition, no clinical guidelines based on research recommend CES as a treatment for any condition. Therefore, cranial electrostimulation therapy (CES) is considered investigational for all indications.

## REFERENCES

1. Gunther M, Phillips KD. Cranial electrotherapy stimulation for the treatment of depression. *Journal of psychosocial nursing and mental health services*. 2010;48(11):37-42. PMID: 20669869
2. The US Food and Drug Administration (FDA) DEN180057 Non-implanted nerve stimulator for functional abdominal pain relief. [cited 12/01/2022]. 'Available from:' [https://www.accessdata.fda.gov/cdrh\\_docs/pdf18/DEN180057.pdf](https://www.accessdata.fda.gov/cdrh_docs/pdf18/DEN180057.pdf).
3. Ching PY, Hsu TW, Chen GW, et al. Efficacy and Tolerability of Cranial Electrotherapy Stimulation in the Treatment of Anxiety: A Systemic Review and Meta-Analysis. *Front Psychiatry*. 2022;13:899040. PMID: 35757229
4. Price L, Briley J, Haltiwanger S, et al. A meta-analysis of cranial electrotherapy stimulation in the treatment of depression. *J Psychiatr Res*. 2021;135:119-34. PMID: 33477056
5. Yu B, Qiu H, Li J, et al. Noninvasive Brain Stimulation Does Not Improve Neuropathic Pain in Individuals With Spinal Cord Injury: Evidence From a Meta-Analysis of 11 Randomized Controlled Trials. *Am J Phys Med Rehabil*. 2020;99(9):811-20. PMID: 32175926
6. Tan G, Rintala DH, Jensen MP, et al. Efficacy of cranial electrotherapy stimulation for neuropathic pain following spinal cord injury: a multi-site randomized controlled trial with a secondary 6-month open-label phase. *J Spinal Cord Med*. 2011;34(3):285-96. PMID: 21756567
7. Shekelle P, Cook I, Miake-Lye IM, et al. *The Effectiveness and Risks of Cranial Electrical Stimulation for the Treatment of Pain, Depression, Anxiety, PTSD, and Insomnia: A Systematic Review*. Washington DC, 2018, pp.
8. O'Connell NE, Wand BM, Marston L, et al. Non-invasive brain stimulation techniques for chronic pain. A report of a Cochrane systematic review and meta-analysis. *Eur J Phys Rehabil Med*. 2011;47(2):309-26. PMID: 21494222
9. O'Connell NE, Wand BM, Marston L, et al. Non-invasive brain stimulation techniques for chronic pain. *Cochrane Database Syst Rev*. 2014;4:CD008208. PMID: 24729198
10. O'Connell NE, Marston L, Spencer S, et al. Non-invasive brain stimulation techniques for chronic pain. *Cochrane Database Syst Rev*. 2018;4:CD008208. PMID: 29652088

11. Boldt I, Eriks-Hoogland I, Brinkhof MW, et al. Non-pharmacological interventions for chronic pain in people with spinal cord injury. *Cochrane Database Syst Rev.* 2014;11:CD009177. PMID: 25432061
12. Kavirajan HC, Lueck K, Chuang K. Alternating current cranial electrotherapy stimulation (CES) for depression. *Cochrane Database Syst Rev.* 2014;7:CD010521. PMID: 25000907
13. Lane-Brown A, Tate R. Interventions for apathy after traumatic brain injury. *Cochrane Database Syst Rev.* 2009(2):CD006341. PMID: 19370632
14. Kim J, Kim H, Kim DH, et al. Effects of cranial electrotherapy stimulation with novel in-ear electrodes on anxiety and resting-state brain activity: A randomized double-blind placebo-controlled trial. *Journal of affective disorders.* 2021;295:856-64. PMID: 34706456
15. Kovacic K, Hainsworth K, Sood M, et al. Neurostimulation for abdominal pain-related functional gastrointestinal disorders in adolescents: a randomised, double-blind, sham-controlled trial. *Lancet Gastroenterol Hepatol.* 2017;2(10):727-37. PMID: 28826627
16. Krasaelap A, Sood MR, Li BUK, et al. Efficacy of Auricular Neurostimulation in Adolescents With Irritable Bowel Syndrome in a Randomized, Double-Blind Trial. *Clin Gastroenterol Hepatol.* 2020;18(9):1987-94 e2. PMID: 31622740
17. Roh HT, So WY. Cranial electrotherapy stimulation affects mood state but not levels of peripheral neurotrophic factors or hypothalamic- pituitary-adrenal axis regulation. *Technology and health care : official journal of the European Society for Engineering and Medicine.* 2017;25(3):403-12. PMID: 27886020
18. Scherder E, Knol D, van Someren E, et al. Effects of low-frequency cranial electrostimulation on the rest-activity rhythm and salivary cortisol in Alzheimer's disease. *Neurorehabil Neural Repair.* 2003;17(2):101-8. PMID: 12814055
19. Scherder EJ, van Tol MJ, Swaab DF. High-frequency cranial electrostimulation (CES) in patients with probable Alzheimer's disease. *Am J Phys Med Rehabil.* 2006;85(7):614-8. PMID: 16788393
20. Winick RL. Cranial electrotherapy stimulation (CES): a safe and effective low cost means of anxiety control in a dental practice. *Gen Dent.* 1999;47(1):50-5. PMID: 10321152
21. Schmitt R, Capo T, Boyd E. Cranial electrotherapy stimulation as a treatment for anxiety in chemically dependent persons. *Alcohol Clin Exp Res.* 1986;10(2):158-60. PMID: 3521373
22. Rose KM, Taylor AG, Bourguignon C. Effects of cranial electrical stimulation on sleep disturbances, depressive symptoms, and caregiving appraisal in spousal caregivers of persons with Alzheimer's disease. *Appl Nurs Res.* 2009;22(2):119-25. PMID: 19427574
23. Lichtbroun AS, Mei-Ming C. The treatment of fibromyalgia with cranial electrotherapy stimulation. *J Clin Psychiatry.* 1984;45(2):60-3. PMID: No PMID Entry
24. Kapkin O, Satar B, Yetiser S. Transcutaneous electrical stimulation of subjective tinnitus. A placebo-controlled, randomized and comparative analysis. *ORL J Otorhinolaryngol Relat Spec.* 2008;70(3):156-61. PMID: 18401195
25. Lichtbroun AS, Raicer MM, Smith RB. The treatment of fibromyalgia with cranial electrotherapy stimulation. *J Clin Rheumatol.* 2001;7(2):72-8; discussion 78. PMID: 17039098
26. Pickworth WB, Fant RV, Butschky MF, et al. Evaluation of cranial electrostimulation therapy on short-term smoking cessation. *Biol Psychiatry.* 1997;42(2):116-21. PMID: 9209728



27. Lee SH, Kim WY, Lee CH, et al. Effects of cranial electrotherapy stimulation on preoperative anxiety, pain and endocrine response. *J Int Med Res.* 2013;41:1788-95. PMID: 24265330
28. Gong BY, Ma HM, Zang XY, et al. Efficacy of Cranial Electrotherapy Stimulation Combined with Biofeedback Therapy in Patients with Functional Constipation. *Journal of neurogastroenterology and motility.* 2016;22(3):497-508. PMID: 26932836
29. Kang HW, Kim HJ, Kim WY, et al. Effects of cranial electrotherapy stimulation on preoperative anxiety and blood pressure during anesthetic induction in patients with essential hypertension. *J Int Med Res.* 2020;48(8):300060520939370. PMID: 32865096
30. Wu WJ, Wang Y, Cai M, et al. A double-blind, randomized, sham-controlled study of cranial electrotherapy stimulation as an add-on treatment for tic disorders in children and adolescents. *Asian J Psychiatr.* 2020;51:101992. PMID: 32145674
31. Barclay TH, Barclay RD. A clinical trial of cranial electrotherapy stimulation for anxiety and comorbid depression. *Journal of affective disorders.* 2014;164:171-7. PMID: 24856571
32. McClure D, Greenman SC, Koppolu SS, et al. A Pilot Study of Safety and Efficacy of Cranial Electrotherapy Stimulation in Treatment of Bipolar II Depression. *The Journal of nervous and mental disease.* 2015;203(11):827-35. PMID: 26414234
33. Ochi M, Saeki S, Oda T, et al. Effects of anodal and cathodal transcranial direct current stimulation combined with robotic therapy on severely affected arms in chronic stroke patients. *Journal of rehabilitation medicine : official journal of the UEMS European Board of Physical and Rehabilitation Medicine.* 2013;45(2):137-40. PMID: 23306448
34. Tyler ME, Kaczmarek KA, Rust KL, et al. Non-invasive neuromodulation to improve gait in chronic multiple sclerosis: a randomized double blind controlled pilot trial. *J Neuroeng Rehabil.* 2014;11:79. PMID: 24885412
35. Lande RG, Gragnani C. Efficacy of cranial electric stimulation for the treatment of insomnia: a randomized pilot study. *Complementary therapies in medicine.* 2013;21(1):8-13. PMID: 23374200
36. Brignani D, Ruzzoli M, Mauri P, et al. Is transcranial alternating current stimulation effective in modulating brain oscillations? *PLoS One.* 2013;8:e56589. PMID: 23457586

## CODES

Codes	Number	Description
CPT	0720T	Percutaneous electrical nerve field stimulation, cranial nerves, without implantation
	64999	Unlisted procedure, nervous system
HCPCS	A4596	Cranial electrotherapy stimulation (ces) system supplies and accessories, per month
	E1399	Durable medical equipment, miscellaneous
	K1002	Cranial electrotherapy stimulation (ces) system, any type

**Date of Origin:** April 2007