

**Regence BlueCross BlueShield of Oregon • Regence BlueShield  
Regence BlueCross BlueShield of Utah • Regence BlueShield of Idaho  
Independent licensees of the Blue Cross and Blue Shield Association**

**Medication Policy Manual**

**Policy No:** dru157

**Topic:** Boniva<sup>®</sup>, ibandronate oral

**Date of Origin:** July 18, 2008

**Revised/Effective:** July 17, 2009

**Next Review Date:** July 2010

**IMPORTANT REMINDER**

This Medical Policy has been developed through consideration of medical necessity, generally accepted standards of medical practice, and review of medical literature and government approval status.

Benefit determinations should be based in all cases on the applicable contract language. To the extent there are any conflicts between these guidelines and the contract language, the contract language will control.

The purpose of medical policy is to provide a guide to coverage. Medical Policy is not intended to dictate to providers how to practice medicine. Providers are expected to exercise their medical judgment in providing the most appropriate care.

**Description**

Ibandronate (Boniva<sup>®</sup>) is a bisphosphonate, a medication used to treat or prevent osteoporosis (bone loss).

## **Policy/Criteria**

**I.** Most contracts require prior authorization approval of oral ibandronate prior to coverage. Oral ibandronate may be considered medically necessary when:

**A.** a generic bisphosphonate (such as alendronate) has not been tolerated or is contraindicated.

### **AND**

**B.** risedronate has not been tolerated or is contraindicated.

**II.** Administration and Authorization Period

**A.** Regence considers oral ibandronate to be a self-administered medication.

**B.** Authorization may be reviewed at least annually to confirm that current medical necessity criteria are met.

## **Position Statement**

### *Summary*

- Osteoporosis is characterized by low bone mass and structural deterioration, leading to bone fragility and an increased risk of fractures of the hip, spine and wrist.<sup>[7]</sup> The objective of treatment of osteoporosis is to reduce the incidence of fractures.
- Generic alendronate provides the best value for the prevention and treatment of osteoporosis.
- There is no reliable evidence that one bisphosphonate is more effective for fracture prevention than another.
- There is no reliable evidence that one bisphosphonate is safer or better tolerated than another.
- There is no reliable comparative evidence to differentiate harms.
- All oral bisphosphonates carry some degree of risk of gastric irritation (oral formulations more so than injectables).

### *Clinical Efficacy*

- There is evidence that alendronate (Fosamax<sup>®</sup>), ibandronate (Boniva), risedronate (Actonel<sup>®</sup>), and zoledronic acid (Reclast<sup>®</sup>) decrease the risk of fracture.
- There are clinical trials of at least three years duration showing that alendronate (Fosamax), risedronate (Actonel), ibandronate (Boniva), and zoledronic acid (Reclast) each reduce the risk of fracture in patients with osteoporosis. Fourteen to 23 patients need to be treated for 3 years to prevent one vertebral fracture. This is similar between the bisphosphonate medications.
- All fracture data for the oral bisphosphonates are based on the daily dosage form only. There are no fracture data for the weekly or monthly products or other modified dosing regimens.
- Bisphosphonates increase bone mineral density (BMD), which correlates to a decrease in the risk of fractures.
- There is no evidence demonstrating superiority of any one bisphosphonate over another for the prevention of fractures. None of the head-to-head comparisons between agents had large enough sample sizes to detect differences. <sup>[4]</sup>

### *Safety*

- Side effects are similar for all oral bisphosphonate medications and include gastrointestinal problems such as difficulty swallowing, gastric ulcer and inflammation of the esophagus. <sup>[4,17]</sup>
- Based on the FDA safety review regarding the possibility of an association between atrial fibrillation and bisphosphonate therapy the FDA states that healthcare professionals should not alter their prescribing patterns for bisphosphonates and patients should not stop taking their bisphosphonate medication. <sup>[18]</sup>
- All of the bisphosphonates have extremely long presence in bone - years for oral, decades for IV zoledronic acid. The long term toxicity is unknown.
- Injectable products are options for patients with contraindications to the oral formulations, and are associated with fewer GI adverse events.
- There have been reports of osteonecrosis of the jaw (particularly following intravenous bisphosphonate treatment for patients with cancer). The level of risk for osteonecrosis in patients being treated for osteoporosis with bisphosphonates is not known, but appears extremely small for at least up to 5 years. <sup>[17]</sup>

## References

1. Wells GA, Cranney A, Peterson J, Boucher M, Shea B, Robinson V, Coyle D, Tugwell P. Alendronate for the primary and secondary prevention of osteoporotic fractures in postmenopausal women. *Cochrane Database of Systematic Reviews* 2008, Issue 1. Art. No.:CD001155. DOI: 10.1002/14651858.CD001155.pub2.
2. The care of patients with fragility fracture. National Institute for Health Research (NIHR) Health Technology Assessment Programme. Published by the British Orthopaedic Association September 2007.
3. Mosekilde L, Vestergaard P, Langdahl B. Fracture prevention in postmenopausal women. *BMJ Clin Evid* 2008;01:1109.
4. MacLean C, Alexander A, Carter J, Chen S, Desai SB, Grossman J, Maglione M, McMahon M, McNamara M, Mojica W, Newberry S, Ranganath V, Suttorp M, Timmer M, Tringale C, Valentine D, Zhou A. Comparative Effectiveness of Treatments To Prevent Fractures in Men and Women With Low Bone Density or Osteoporosis. Comparative Effectiveness Review No. 12. (Prepared by Southern California/RAND Evidence-based Practice Center under Contract No. 290-02-0003). Rockville, MD: Agency for Healthcare Research and Quality. December 2007. Available at: [www.effectivehealthcare.ahrq.gov/reports/final.cfm](http://www.effectivehealthcare.ahrq.gov/reports/final.cfm).
5. Wells G, Cranney A, Peterson J, Boucher M, Shea B, Robinson V, Coyle D, Tugwell P. Risedronate for the primary and secondary prevention of osteoporotic fractures in postmenopausal women. *Cochrane Database of Systematic Reviews* 2008, Issue 1. Art. No.: CD004523. DOI: 10.1002/14651858.CD004523.pub3.
6. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the prevention and treatment of postmenopausal osteoporosis: 2001 edition, with selected updates for 2003. *Endocr Pract* 2003;9(6):544-64.
7. Osteoporosis Overview. National Institutes of Health Osteoporosis and Related Bone Diseases ~ National Resource Center. Updated December 2007. Available at [http://www.niams.nih.gov/Health\\_Info/Bone/](http://www.niams.nih.gov/Health_Info/Bone/) accessed March 12, 2008.
8. Thomson Micromedex © 1974-2008. Micromedex Healthcare Series, Vol 120.
9. Actonel<sup>®</sup> (risedronate sodium) Tablet Prescribing Information. April 2008 Procter & Gamble Pharmaceuticals, Inc. Cincinnati, OH.

10. Fosamax<sup>®</sup> (alendronate sodium) Prescribing Information. February 2008. Merck & Co., Inc. Whitehouse Station, NJ.
11. Boniva<sup>®</sup> (ibandronate sodium) Tablets Prescribing Information. November 2008. Roche Laboratories Inc., Nutley, NJ.
12. Forteo<sup>®</sup> teriparatide (rDNA origin) injection Prescribing Information. June 2008. Eli Lilly and Company - Indianapolis, IN
13. Evista<sup>®</sup> (raloxifene hydrochloride) Tablets Prescribing Information. October 2008. Eli Lilly and Company - Indianapolis, IN.
14. Reclast<sup>®</sup> (zoledronic acid) Injection Prescribing Information. May 2009. Novartis Pharmaceuticals Corporation, East Hanover, NJ.
15. Black DM, Delmas PD, Eastell R, Reid IR, Boonen S, et al. Once-yearly zoledronic acid for the treatment of postmenopausal osteoporosis. *N Engl J Med*. 2007;356:1809-1822.
16. U.S. Food and Drug Administration [page on the internet] 2007. Early communication of an ongoing safety review. Available at:  
[http://www.fda.gov/cder/drug/early\\_comm/bisphosphonates.htm](http://www.fda.gov/cder/drug/early_comm/bisphosphonates.htm). Accessed on 10/17/07
17. Clinician's Guide To Prevention And Treatment Of Osteoporosis. National Osteoporosis Foundation, 2008. Available at: [http://www.nof.org/professionals/Clinicians\\_Guide.htm](http://www.nof.org/professionals/Clinicians_Guide.htm). Accessed March 12, 2008.
18. U.S. Food and Drug Administration [page on the internet] 2008. November 12, 2008 [Update to the October 1, 2007 Early Communication about the Ongoing Safety Review of Bisphosphonates](#). Available at:  
<http://www.fda.gov/Safety/MedWatch/SafetyInformation/SafetyAlertsforHumanMedicalProducts/ucm079352.htm> accessed June 22, 2009

<b>Cross References</b>
Actonel <sup>®</sup> , risedronate dru155
Boniva <sup>®</sup> , ibandronate injection dru156
Forteo <sup>®</sup> , teriparatide dru085
Reclast <sup>®</sup> , zoledronic acid dru158

<b>Codes</b>	<b>Number</b>	<b>Description</b>
HCPCS	J1740	Injection, ibandronate sodium, 1 mg (Boniva 3 MG/3ML KIT)
HCPCS	J3488	Injection, zoledronic acid (Reclast), 1 mg (Reclast 5 MG/100ML SOLN)

**Appendix 1:** Indications for specified osteoporosis therapy

	<b>Alendronate</b>	<b>Risedronate</b>	<b>Ibandronate</b>	<b>Zoledronic acid</b>
Postmenopausal Osteoporosis: Prevention	✓	✓	✓	✓
Postmenopausal Osteoporosis: Treatment	✓	✓	✓	✓
Glucocorticoid-Induced Osteoporosis: Prevention	-	✓	-	✓
Glucocorticoid-Induced Osteoporosis: Treatment	✓	✓	-	-✓
Osteoporosis in Men	✓	✓	-	-✓
Paget's disease	✓	✓	-	✓

## Appendix 2: Dosing regimens of bisphosphonate therapy

Drug Products	Usual Dose/Route	Administration Considerations
alendronate (Fosamax <sup>®</sup> ) <sup>10</sup>	<ul style="list-style-type: none"> <li>• 5 mg p.o. daily</li> <li>• 10 mg p.o. daily</li> <li>• 35 mg p.o. weekly</li> <li>• 70 mg p.o. weekly</li> <li>• 40mg p.o. daily x 6mo</li> </ul>	<ul style="list-style-type: none"> <li>• Taken 30 minutes before first food, beverage, or medication</li> <li>• Swallowed with a full glass of water (180 to 240 mL).</li> <li>• Followed by at least 60 mL (one-fourth cup) of water</li> <li>• Do not lie down for at least 30 minutes and until after their first food of the day</li> </ul>
ibandronate (Boniva <sup>®</sup> ) <sup>11</sup>	<ul style="list-style-type: none"> <li>• 2.5 mg p.o. daily</li> <li>• 150 mg p.o. monthly</li> <li>• 3 mg IV every 3 months</li> </ul>	<ul style="list-style-type: none"> <li>• Taken 60 minutes before the first food or drink (other than water)</li> <li>• Swallowed whole with a full glass of plain water (180 to 240 mL; 6 to 8 oz)</li> <li>• Do not lie down for 60 minutes</li> </ul>
risedronate (Actonel <sup>®</sup> ) <sup>9</sup>	<ul style="list-style-type: none"> <li>• 5 mg p.o. daily</li> <li>• 35 mg p.o. weekly</li> <li>• 30 mg p.o. daily x 2mo (Paget's)</li> <li>• 75 mg p.o. two consecutive days/month</li> <li>• 150mg p.o. monthly</li> </ul>	<ul style="list-style-type: none"> <li>• Taken 30 minutes before the first food or drink of the day other than water</li> <li>• Full glass of plain water (6 to 8 oz).</li> <li>• Do not lie down for 30 minutes</li> </ul>
zoledronic acid injectable (Reclast <sup>®</sup> ) <sup>14</sup>	<ul style="list-style-type: none"> <li>• 5 mg IV once annually</li> </ul>	<ul style="list-style-type: none"> <li>• Given over no less than 15 minutes</li> </ul>