

Gene-Based Tests for Screening, Detection, and Management of Prostate or Bladder Cancer

Effective: October 1, 2020

Next Review: September 2021

Last Review: September 2020

IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

DESCRIPTION

There are a variety of gene-based biomarkers that have been associated with prostate cancer or bladder cancer. These tests have the potential to improve the accuracy of risk prediction, diagnosis, staging, or prognosis of prostate cancer.

MEDICAL POLICY CRITERIA

Genetic tests for the screening, detection, and management of prostate cancer or bladder cancer are considered **investigational**, including but not limited to the following:

- A. Single-nucleotide variants (SNVs) for risk assessment;
- B. PCA3 for disease diagnosis;
- C. TMPRSS fusion genes for diagnosis and prognosis;
- D. Gene hypermethylation for diagnosis and prognosis;
- E. Mitochondrial DNA variant testing for diagnosis;
- F. Gene expression analysis, including but not limited to Prolaris®, Oncotype DX® Prostate, Decipher® Prostate (Biopsy and RP), and Decipher® Bladder.

NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.

CROSS REFERENCES

1. [Gene Expression-Based Assays for Cancers of Unknown Primary](#), Genetic Testing, Policy No. 15
2. [Genetic and Molecular Diagnostic Testing](#), Genetic Testing, Policy No. 20
3. [Circulating Tumor DNA and Circulating Tumor Cells for Management \(Liquid Biopsy\) of Solid Tumor Cancers](#), Laboratory, Policy No. 46
4. [Protein Biomarkers for Screening, Detection, and/or Management of Prostate Cancer](#), Laboratory, Policy No. 69

BACKGROUND

Prostate cancer and bladder cancer are complex, heterogeneous diseases. Some forms may be indolent and never progress to cause harm, while other metastasize quickly. Current challenges in cancer care are assessing risk; providing early and accurate detection; monitoring low-risk patients undergoing surveillance only; predicting recurrence after initial treatment; detecting recurrence after treatment; and assessing efficacy of treatment for advanced disease.

In response to the need for better biomarkers for risk assessment, diagnosis, and prognosis, a variety of exploratory research is ongoing. Some products of this work have already been translated or are in the process of being translated into commercially available tests, including:

- Single-nucleotide variants (SNVs) for risk assessment
- The Gen-Probe PROGENSA® *PCA3* Assay (*PCA3*) for diagnosis
- *TMPRSS* fusion genes for diagnosis and prognosis
- Gene hypermethylation for diagnosis and prognosis
- Mitochondrial DNA variant testing for diagnosis
- Gene expression analysis for risk assessment and diagnosis, including Prolaris®, Oncotype DX® Prostate, Decipher® Prostate, and Decipher® Bladder
- Exosomal RNA-based urine assay ExoDx™ Prostate (Intelliscore) for diagnosis

While studies using these tests generate information that may help elucidate the biologic mechanisms of prostate or bladder cancer and eventually help design treatments, the above-mentioned tests are currently in a developmental phase, without evidence of clinical utility for diagnosis, prognosis, or risk assessment. Many of the tests listed above have not been submitted to the U.S. Food and Drug Administration (FDA) for marketing clearance but, if available, are offered as laboratory-developed tests by Clinical Laboratory Improvement Amendments (CLIA) licensed laboratories.

SNV testing as part of genome-scanning tests with risk assessment for prostate cancer is offered by a variety of laboratories including Navigenics, LabCorp (23andme), and ARUP (deCode) as laboratory-developed tests. The *PCA3* test is offered in the U.S. by reference laboratories including ARUP, Mayo Medical Laboratories, and LabCorp. The reagents used in testing are developed by Gen-Probe. A test for hypermethylation of *GSTP1* was available from LabCorp (“Glutathione S-transferase Gene [*GSTP1*, pi-class] Methylation Assay”); but as of January 2015, this test is no longer offered.

ConfirmMDx is offered from MDxHealth. The tissue-based DNA methylation multigene assay aims to improve stratification of men being considered for repeat prostate biopsy. Hypermethylation of *GSTP1*, *APC*, and *RASSF1* are assessed in core biopsy samples.

SelectMDx for Prostate Cancer is also offered from MDxHealth. The reverse transcription PCR (RT-PCR) assay is performed on post-DRE (digital rectal examination), first-void urine specimens from patients with clinical risk factors for prostate cancer, who are being considered for biopsy. The test measures the mRNA levels of the DLX1 and HOXC6 biomarkers, using KLK3 expression as internal reference, to aid in patient selection for prostate biopsy.

Regulatory Status

One mitochondrial DNA test, Mitomics (Broomfield, CO), is currently available. Mitomics offers the Prostate Core Mitomics Test which measures mitochondrial DNA variants in a negative prostate biopsy to determine whether a patient should undergo repeat biopsy. The test is performed on the initial negative prostate biopsy tissue.

The PROGENSA® PCA3 Assay was approved by the FDA on February 15, 2012 through the premarket approval process. According to the approval granted by the FDA:^[1]

“The PROGENSA PCA3 Assay is indicated for use in conjunction with other patient information to aid in the decision for repeat biopsy in men 50 years of age or older who have had one or more previous negative prostate biopsies and for whom a repeat biopsy would be recommended by a urologist based on current standard of care, before consideration of PROGENSA PCA3 assay results.”

Prolaris®, Oncotype DX® Prostate, and Decipher® gene expression profiling tests are available under the auspices of CLIA. Laboratories that offer LDTs must be licensed by CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of this test.

In November 2015, the FDA’s Office of Public Health Strategy and Analysis published a document on public health evidence for FDA oversight of LDTs.^[2] FDA argued that many tests need more FDA oversight than the regulatory requirements of CLIA. CLIA standards relate to laboratory operations, but do not address inaccuracies or unreliability of specific tests. Prolaris® is among the 20 case studies in the document cited as needing FDA oversight. The document asserted that patients are potentially receiving inappropriate prostate cancer care because there is no evidence that results from the test meaningfully improve clinical outcomes.

The other tests mentioned in this policy, if available, are offered as laboratory-developed tests under the Clinical Laboratory Improvement Amendments (CLIA) licensed laboratories.

EVIDENCE SUMMARY

Human Genome Variation Society (HGVS) nomenclature^[3] is used to describe variants found in DNA and serves as an international standard. It is being implemented for genetic testing medical evidence review updates starting in 2017. According to this nomenclature, the term “variant” is used to describe a change in a DNA or protein sequence, replacing previously-used terms, such as “mutation.” Pathogenic variants are variants associated with disease, while benign variants are not. The majority of genetic changes have unknown effects on human health, and these are referred to as variants of uncertain significance

In general, the evidence for genetic tests related to prostate cancer screening, detection, and management addresses either preliminary clinical associations between genetic tests and disease states or, in some cases, the clinical validity of these tests i.e., the association of the test result with outcomes of interest, expressed in terms of clinical performance characteristics

such as sensitivity, specificity, predictive value, and comparisons to current standards using receiver-operating curve (ROC) analysis and/or logistic regression. There is no published evidence demonstrating clinical utility (i.e., a test will change treatment decisions and improve patient important outcomes).

GENE-BASED TESTS IN GENERAL

A 2009 BlueCross BlueShield Association (BCBSA) TEC Special report of recently published studies on gene-based tests (SNVs, *PCA3*, *TMPRSS*, gene panels, and gene hypermethylation) for prostate cancer risk assessment and diagnosis concluded that, in general, research on these tests is still in a “developmental phase, currently without evidence of clinical utility.”^[4] This policy was initially based on a 2013 BlueCross BlueShield Association Technology Evaluation Center (TEC) Assessment which was updated in January 2015 with a literature review through September 30, 2014.^[5,6] Full-length publications were sought that described the analytic validity (technical performance), clinical validity (prognostic accuracy), and clinical utility (accurately identifying men experiencing improved health outcomes by avoiding treatment or undergoing more appropriate therapies) of Prolaris, Oncotype DX Prostate, and Decipher® gene expression profiling. The Blue Cross Blue Shield Association Medical Advisory Panel also reviewed the evidence in September 2017.

SINGLE-NUCLEOTIDE VARIANTS (SNVS) FOR PROSTATE CANCER RISK ASSESSMENT AND PROGNOSIS

There have been numerous large observational correlational studies focusing on the association of many different SNVs with prostate cancer, an example of which includes the study by Lindstrom of 10,501 cases of prostate cancer and 10,831 controls, which identified 36 SNVs showing association with prostate cancer risk including two (*rs2735893* and *rs266849*) showing differential association with Gleason grade. Per allele odds ratios ranged from 1.07 to 1.44.^[7]

Because the SNVs individually provide relatively modest incremental information on both the occurrence of cancer and its behavior, investigators have begun to explore use of algorithms incorporating information from multiple SNVs to increase the clinical value of testing. Several such recent studies focused on the development of testing algorithms incorporating SNVs.^[8-11]

Systematic Reviews

A systematic review of multigene panels for prostate cancer risk assessment was published by Little (2016).^[12] The authors included 21 studies that evaluated 18 individual panels. All studies were focused on clinical validity, moderate risk of bias, and had poor discriminative ability for predicting prostate cancer risk and/or distinguishing between aggressive and latent cancers. The authors noted that the current evidence is insufficient to assess analytic validity, and that “at best the panels assessed would add a small and clinically unimportant improvement” to current factors used for risk stratification, like age and family history. Additionally, they found no evidence on the clinical utility of these panels.

A 2012 AHRQ report on multigene panels in prostate cancer risk assessment, also by Little, reviewed the literature on SNV panel tests for assessing risk of prostate cancer.^[13] All of the studies included in the review had poor discriminative ability for predicting risk of prostate cancer, had moderate risk of bias, and none of the panels had been evaluated in routine clinical settings. The conclusions of the review were that the evidence on currently available

SNV panels does not permit meaningful assessment of analytic validity, the limited evidence on clinical validity is insufficient to conclude that SNV panels would perform adequately as a screening test and that there is no evidence available on the clinical utility of current panels.

Ishaak (2011) reviewed 11 replication studies involving 30 SNVs (19 in men of African descent and 10 in men with familial prostate cancer).^[14] Odds ratios were positively associated with prostate cancer, although the magnitude of association was generally small (range 1.11 to 2.63).

Amin Al Olama (2013) conducted a meta-analysis of four GWASs including 5,953 cases of aggressive prostate cancer (PCa) and 11,463 controls.^[15] Authors computed association tests for approximately 2.6 million SNVs and followed up the most significant SNVs by genotyping 49,121 samples in 29 studies through the international PRACTICAL and BPC3 consortia. The authors confirmed the association of a PCa susceptibility locus, rs11672691 on chromosome 19, but also showed an association with aggressive PCa (odds ratio [OR] 1.12, 95% confidence interval [CI] 1.03 to 1.21, $p=1.4 \times 10^{-8}$). The authors concluded their report described a genetic variant which is associated with aggressive PCa, and which is a type of PCa associated with a poorer prognosis.

Nonrandomized Studies

A pilot study by Castro (2016) tested the use of a 71-SNV panel in 100 men with a family history of prostate cancer.^[16] These men underwent a prostate biopsy regardless of PSA level, and 25 were diagnosed with prostate cancer. Age and PSA level were significantly associated with a cancer diagnosis, but the SNV risk score was not. While this study might not have been adequately powered to detect such an association, there was a clear relationship seen for age and PSA level ($p=0.00004$ and 0.00037 , respectively).

Kader (2012) evaluated a panel of 33 SNVs associated with prostate cancer in 1,654 men.^[17] Genetic score was a significant ($p<0.001$) independent predictor of prostate cancer (OR 1.72, 95% CI 1.44 to 2.09) after adjustment for clinical variables and family history. Addition of genetic markers to the classification of prostate cancer risk resulted in 33% of men reclassified into a different risk quartile. Approximately half of these ($n=267$) were downgraded to a lower risk quartile and the other half ($n=265$) were upgraded into a higher risk quartile. The net reclassification benefit was 10% ($p=0.002$). The authors concluded that with the additional information of genetic score the same number of cancers could be detected by using 15% fewer biopsies. However, this study includes a limited sample size and there is no clear indication of how clinical management changed when patients were reclassified into lower risk groups.

Ren (2013) calculated genetic scores for various combinations of 29 PCa risk-associated SNVs in 667 consecutive patients that underwent prostate biopsy.^[18] Performance of these genetic scores for discriminating prostate biopsy outcomes were compared using the area under a receiver operating characteristic curve (AUC). The discriminative performance of genetic score derived from a panel of all 29 SNVs (24 previous and five new) was similar to that derived from the 24 previously established SNVs, the AUCs of which were 0.60 and 0.61, respectively ($p=0.72$). Authors concluded that genetic score based on PCa risk-associated SNVs implicated to date is a significant predictor of biopsy outcome.

Tsuchiya (2013) identified 14 SNVs in six genes (*XRCC4*, *PMS1*, *GATA3*, *IL13*, *CASP8*, and *IGF1*) that were statistically associated with cancer-specific survival.^[19] Using a subset of six

SNVs, three subgroups of men with prostate cancer were defined by the number of SNV's present (0 to 1, 2 to 3, or 4 to 6). Median cancer-specific survival in these subgroups was 13.3, 7.0, and 3.8 years, respectively (log-rank test, $p < 0.001$).

Section Summary

Numerous studies have demonstrated the association of many gene panels and SNVs with prostate cancer. These studies, in early stages of development, have shown a modest degree of association with future risk for prostate cancer. The clinical utility of these tests is uncertain; there is no evidence that information obtained from gene panels or SNV testing can be used to change clinical management in ways that will improve outcomes.

PCA3 FOR PROSTATE CANCER DIAGNOSIS

PCA3 is overexpressed in prostate cancer and *PCA3* mRNA can be detected in urine samples collected after prostate massage. When normalized using PSA to account for the amount of prostate cells released into the urine (*PCA3* Score), the test has been proposed for use in discriminating between patients with eventual benign findings on (first or second) biopsies from those with malignant biopsy results. In particular, the test may be especially helpful at identifying patients with elevated PSA levels but negative first biopsy results who need a follow-up biopsy.

Systematic Reviews

A meta-analysis by Lee (2020) compared *PCA3* in urine or prostatic fluid to prostate biopsy results using data from 54 case-control and cohort studies.^[20] Of the 17,575 total patients, 4,034 had a diagnosis of prostate cancer. Using a *PCA3* cutoff of 35, the pooled sensitivity and specificity were 0.71 (95% CI 0.67 to 0.74) and 0.68 (95% CI 0.63 to 0.74), respectively. Study heterogeneity, I^2 , was 81.1 for sensitivity and 96.4 for specificity. The AUC was 0.75 (95% CI 0.71 to 0.79).

Qin (2020) published a meta-analysis on the diagnostic accuracy of *PCA3* measurement that included 65 retrospective case-control studies published between 2003 and 2018.^[21] There were a total of 8,139 cases and 14,116 controls. Of these, 22,255 had a *PCA3* test, including 5,065 cases. All but two of the studies used urine levels of *PCA3*. The reported pooled sensitivity and specificity were 0.68 (95% CI 0.64 to 0.72) and 0.72 (95% CI 0.68 to 0.75), respectively, and the AUC was 0.76 (95% CI 0.72 to 0.79), but it was not clear what cutoff values were used for these calculations.

A systematic review and meta-analysis by Rodríguez (2020) focused on urine *PCA3* measurement in patients that had not yet undergone prostate biopsy.^[22] Nine studies met the inclusion criteria and of these, five that used 35 as a cutoff value were included in the meta-analysis. The pooled sensitivity was 0.69 (95% CI 0.61 to 0.75, $I^2 = 0\%$), and the pooled specificity was 0.65 (95% CI 0.55 to 0.73, $I^2 = 0\%$). The AUC was 0.73 (95% CI 0.67 to 0.80). The authors noted that the bias assessment identified an unclear risk of bias related to study flow and timing for most studies.

Cui (2016) reported on results of a systematic review that searched PubMed and EMBASE for case-control or cohort studies of the PROGENSA® *PCA3* test.^[23] Quality was assessed using the QUADAS tool. Pooled estimates were calculated using random-effects models and summarized ROCs when evidence of threshold effect was detected. The review included 46 studies with over 12,000 men. The quality of the selected studies was rated as moderate to

high. The most common *PCA3* cutoff for categorizing low and high risk was 35; 25 studies had a *PCA3* cutoff of 35. Most were performed in the United States and Europe; five were conducted in Asia. The estimates of AUC were lower for studies including men having repeated (0.68, 95% CI 0.67 to 0.70) vs initial (0.80, 95% CI 0.78 to 0.82) biopsies. AUC values were 0.74 (95% CI 0.73 to 0.76) for studies with a cutoff value of 35 and 0.77 (95% CI 0.75 to 0.79) for studies with a cutoff value not equal to 35, although the group with varying cutoff ($\neq 35$) had a greater range and more variable performance estimates

Nicholson (2015) published a health technology assessment on behalf of the National Health Service in England and Wales.^[24] Publications from 2000 to May 2014 were included in a systematic review. Participants were men suspected of having prostate cancer for whom the results of an initial prostate biopsy were negative or equivocal; and a *PCA3* score or phi in combination with existing standard tests, multiparametric MRI and clinical judgement were evaluated for analytic validity and clinical validity. Overall, six studies met inclusion for the analytical validity review; and fifteen studies met inclusion for the clinical validity review. The authors found issues regarding the precision of the *PCA3* assay measurements, and insufficient evidence to identify useful clinical thresholds.

In 2013, the Agency for Healthcare Quality and Research (AHRQ) published a comparative effectiveness review entitled, "*PCA3* Testing for the Diagnosis and Management of Prostate Cancer."^[25] Literature was searched and updated through May 15, 2012. Forty-three studies were included; all were rated poor quality. In their conclusion, the authors stated, "For diagnostic accuracy, there was a low strength of evidence that *PCA3* had better diagnostic accuracy for positive biopsy results than [serum] total PSA elevations, but insufficient evidence that this led to improved intermediate or long-term health outcomes." This finding appeared to apply to both initial and repeat biopsies. Evidence was insufficient to assess the use of *PCA3* in treatment decision-making for men with positive biopsy.

In a 2012 systematic review, authors discuss the potential use of genetic markers to better define groups of men at high risk of developing prostate cancer, to improve screening techniques, discriminate indolent versus aggressive disease, and improve therapeutic strategies in patients with advanced disease.^[26] Genetic tests for *PCA3* and *TMPRSS2-ERG* genes were included. Authors concluded that most markers have not been prospectively validated for providing useful prognostic or predictive information or improvement upon clinicopathologic parameters already in use.

A meta-analysis by Ruiz-Aragon (2010) reviewed 14 studies of *PCA3* for use in predicting prostate biopsy results.^[27] Sensitivity of testing ranged from 47% to 82% and specificity from 56% to 89%. Global results provided a sensitivity of 85% (CI 84 to 87) and a specificity of 96% (CI 96 to 97). No publications on how this information affected decision making or either short- or long-term outcomes has been published.

Randomized Controlled Trials

In 2014, the National Cancer Institute conducted a prospective trial to validate the diagnostic use of *PCA3* to complement PSA-based detection of prostate cancer.^[28] The target population included men who had been screened for prostate cancer, primarily with a PSA test, some of whom had undergone a previous prostate biopsy. The study included 859 men from 11 centers in the United States. The primary study endpoint was the diagnosis of prostate cancer on biopsy and the secondary study endpoint was diagnosis of high-grade prostate cancer, defined as a Gleason score greater than 6. The primary analyses, including *PCA3* thresholds, were

determined a priori, and statistical power was based on independent analyses of prevalidation data from similar cohorts. Of the men in the study, 562 were presenting for their initial prostate biopsy. Positive predictive value was 80% (95% CI 72% to 86%), and using a *PCA3* score of more than 60, diagnostic sensitivity and specificity of *PCA3* was 0.42 (95% CI 0.36 to 0.48) and 0.91 (95% CI 0.87 to 0.94), respectively. For patients who underwent a repeat biopsy, the NPV was 88% (95% CI, 81% to 93%), and by using a *PCA3* score of less than 20, sensitivity and specificity were 0.76 (95% CI 0.64 to 0.86) and 0.52 (95% CI 0.45 to 0.58), respectively. For the detection of high-grade cancer, *PCA3* performance in combination with Prostate Cancer Prevention Trial's (PCPT) risk calculator was improved by the addition of *PCA3* to the PCPT risk calculator factors with an AUC improvement of 0.74 to 0.78 for initial biopsy and 0.74 to 0.79 on repeat biopsy ($p \leq 0.003$).

Nonrandomized Studies

Several studies published between 2013 and 2018 reported positive associations between *PCA3* levels and prostate cancer diagnosis.^[29-37] Predictive value was increased when *PCA3* testing was combined with PSA level and other clinical information.^[37-40] Other groups reported moderate diagnostic accuracy of *PCA3* testing. Among men with PSA level greater than 3 ng/mL, AUC of *PCA3* was 0.74.^[41] Conversely, in men with *PCA3* scores of 100 or greater, positive predictive value was 39%.^[42] In a multicenter study of 647 men, sensitivity and specificity were 67% and 72%, respectively; AUC was 0.742.^[43] Three studies compared *PCA3* to multiparametric MRI; MRI was more accurate than *PCA3*,^[44,45] but the combination was better than either alone.^[46] *PCA3* has also been associated with the risk of Gleason grade reclassification in a study of 90 men receiving 5 α -reductase inhibitor therapy during active surveillance.^[47]

Clinical utility studies using assay results for decision-making for initial biopsy, repeat biopsy, or treatment have not been reported. One group reported potential reductions in unnecessary biopsies of 48 to 52% with attendant increases in missed prostate cancers of 6 to 15% using either a *PCA3*-based nomogram^[48] or *PCA3* level corrected for prostate volume (*PCA3* density).^[49] Although both studies were prospective, neither assessed utility of the test for clinical decision-making because all patients underwent biopsy, and recurrence or survival outcomes were not evaluated. Another group evaluated the estimated long-term impact of using the *PCA3* score to guide the decision to recommend a repeat biopsy for men with elevated PSA levels.^[50] Their models suggested that using a *PCA3* score threshold of 25 would result in 55.4% reduction in repeat biopsies for a base-case patient, while reducing the 10-year survival by 0.93%. However, these estimates have not been validated in real patient populations.

Section Summary

Studies of *PCA3* as a diagnostic test for prostate cancer reported sensitivities and specificities in the moderate range. In general, these studies are preliminary and report on clinical performance characteristics in different populations and with various assay cutoff values, reflecting the lack of standardization in performance and interpretation of *PCA3* results. Several studies have reported a modest incremental improvement in diagnostic accuracy when *PCA3* was tested in combination with PSA level and other clinical findings. The clinical utility of this test is uncertain, as there is insufficient evidence that the use of *PCA3* can be used to change management in ways that improve outcomes.

TMPRSS FUSION GENES

TMPRSS2 fusion gene detection has been studied for prognostic value (e.g., to identify aggressive disease or to predict disease recurrence). In prostate cancer, it may be fused to an ETS family transcription factor (*ERG*, *ETV1*, *ETV4*, or *ETV5*), which modulates transcription of target genes involved in cell growth, transformation, and apoptosis (*TMPRSS2:ERG*). The result of gene fusion with an ETS transcription gene is that the androgen-responsive promoter of *TMPRSS2* upregulates expression of the ETS gene, suggesting a mechanism for neoplastic transformation. Fusion genes may be detected in tissue, serum, or urine.

Systematic Review

Yao (2014) published a systematic review with meta-analysis of *TMPRSS2:ERG* for the detection of prostate cancer.^[51] Literature was searched and 32 articles were identified. Pooled sensitivity, specificity, positive likelihood ratio, and negative likelihood ratio were 47% (95% CI 46 to 49), 93% (95% CI 92 to 94), 8.9 (95% CI 5.7 to 14.1), and 0.49 (95% CI 0.43 to 0.55), respectively. Statistical heterogeneity was high ($I^2 > 85\%$). It was unclear whether studies in screening populations were pooled with enriched patient samples, e.g., elevated PSA and/or biopsy-negative. There also was variability in the type of tissue samples analyzed (urine, prostatic secretions, biopsy or surgical specimens); the type of *TMPRSS2:ERG* assays used (fluorescence in situ hybridization [FISH], immunohistochemistry [IHC], real-time reverse transcriptase polymerase chain reaction [RT-PCR], and transcription-mediated amplification); and in *TMPRSS2:ERG* threshold cutoff values.

Nonrandomized Studies

Leyton (2014) investigated the predictive value of *PCA3* and *TMPRSS2* as individual biomarkers and as part of a panel in a prospective, multicenter study of 443 men.^[52] *TMPRSS2* was found to be highly specific (93%) for predicting clinically significant prostate cancer on biopsy. Because of this high specificity, the authors suggested that re-biopsy or magnetic resonance imaging (MRI) be performed in *TMPRSS2:ERG*-positive patients who do not have prostate cancer detected on initial biopsy. The authors stated that if *PCA3* in combination with *TMPRSS2* data had been used to select men for prostate biopsy, 35% of biopsies could have been avoided. However, the clinical utility of this test is uncertain, as there are no studies that report the test leads to changes in management that result in improved health outcomes.

Whelan (2014) compared two multivariate models to assess up-staging in 216 patients meeting National Comprehensive Cancer Network (NCCN) criteria for active surveillance.^[53] One model included *TMPRSS2:ERG* plus serum PSA; the other model included serum PSA, total RNA in expressed prostatic secretion (EPS, collected by milking the urethra after prostatic massage), and total EPS volume. AUCs were similar (0.80, 95% CI 0.75 to 0.85] and 0.79, 95% CI 0.73 to 0.84], respectively). However, the second model was more accurate for detecting patients who were up-staged, or up-staged and up-graded, by NCCN criteria. Specifically, the second model decreased the risk of up-staging in patients with a negative test approximately eight-fold (from 7% to 1%); decreased the risk of up-staging plus up-grading approximately five-fold (from 5% to 1%); and doubled the prevalence of up-staging in the positive test group. In comparison, the *TMPRSS2:ERG* model decreased up-staging 2.4-fold (from 7% to 3%) and decreased upstaging and upgrading approximately three-fold (from 5% to 2%).

A modeling study by Merdan (2015) estimated that using a *TMPRSS2:ERG* score to guide repeat biopsy decisions in men with elevated PSA could avoid 64.7% of these biopsies, but

also reduce the 10-year survival rate by 1.4%.^[50] These estimations have not been validated in real-world trials.

A number of studies have reported positive associations between *TMPRSS2* fusion gene levels and prostate cancer diagnosis.^[33,34,54,55] One study reported a lack of association between *TMPRSS2:ERG* status and biochemical relapse-free rate in 244 men treated with image-guided radiotherapy (IGRT) for prostate cancer.^[56] The authors concluded that “*TMPRSS2-ERG* is therefore unlikely to be a predictive factor for IGRT response.”

No studies of clinical utility have been published to date; the evidence consists of correlational studies (association between a fusion gene and prostate cancer).^[57-60] However, the results of available studies differ as to the accuracy of *TMPRSS2:ERG* in improving the ability to predict prostate cancer, and/or the ability to estimate prognosis for this purpose.

Section Summary

Limited evidence reports that the measurement of *TMPRSS2:ERG* may improve the ability to predict prostate cancer, and/or the ability to estimate prognosis. However, the results of available studies differ as to the accuracy of *TMPRSS2:ERG* for this purpose. In addition, the clinical utility of this test is uncertain.

TMPRSS2:ERG* IN COMBINATION WITH *PCA3

Nonrandomized Studies

Sanda (2017), from the National Cancer Institute Early Detection Research Network, reported separate developmental and validation cohorts for high-grade prostate cancer in men undergoing initial prostate biopsy.^[61] For the validation cohort, any of the following was considered a positive result: PSA level greater than 10 ng/mL, urine *TMPRSS2:ERG* score greater than 8, or urine *PCA3* score greater than 20. The sensitivity and specificity of this combination were reported to be 92.6% and 33.4%, respectively, an increase from the 17% specificity of *PSA* alone without loss of sensitivity. The difference in specificity was statistically significant, with a prespecified one-sided p-value of 0.04 (lower bound of one-sided 95% CI 0.73%).

Tomlins (2011) developed a transcription-mediated amplification assay to measure *TMPRSS2:ERG* fusion transcripts in parallel with *PCA3*.^[60] Combining results from these two tests and incorporating them into the multivariate Prostate Cancer Prevention Trial risk calculator improved the identification of patients with clinically significant cancer by Epstein criteria and high-grade cancer on biopsy. Though the study was large (1,312 men at multiple centers), results were confounded by assay modifications during the course of the study, by the use of cross-validation rather than independent validation, and the use of independent training and testing sets. A validation study by the same group evaluated this risk prediction model (termed Mi-Prostate Score or MiPS), which incorporated serum PSA with urine *TMPRSS2:ERG*, and *PCA3*, in a group of 1,244 men presenting for biopsy.^[62] They reported that it improved on *PSA* alone for predicting prostate cancer and high-grade prostate cancer, but did not assess the clinical utility of this risk score.

A study by Feibus (2016) evaluated the clinical use of *PCA3* and *TMPRSS2:ERG* in African-American men undergoing prostate biopsy.^[36] This study included 182 African-American and 139 non-African-American patients. They found that *PCA3* and *TMPRSS2:ERG* scores were associated with prostate cancer, and adding *PCA3* to a standard of care plus *PSA* model

improved concordance statistics for the detection of any prostate cancer in both groups. However, *PCA3* score was only predictive of high grade prostate cancer in African-Americans and not in non-African-Americans, while *TMPRSS2:ERG* did not improve these measures in either group.

In a pilot study, Salami (2013) evaluated 45 men using a multivariable algorithm that included serum PSA plus urine *TMPRSS2:ERG* and *PCA3* from a post-DRE sample.^[63] Samples were collected before prostate biopsy at two centers. For cancer prediction, sensitivity and specificity were 80% and 90%, respectively. AUC was 0.88. Limitations in this study included the small number of patients and authors used a quantitative RT-PCR assay to measure *PCA3* in urine sediment, and evaluation of this model with the commercially available TMA assay of whole urine would help inform broader clinical applicability. Authors concluded a larger validation study could determine whether a multiplex model combining *PCA3*, *TMPRSS2:ERG* and serum PSA can predict aggressive versus indolent prostate cancer than any of these biomarkers alone.

Robert (2013) retrospectively examined tissue levels of *TMPRSS2:ERG* and *PCA3* in 48 men with benign prostatic hypertrophy, 32 men with normal prostate tissue sampled next to prostate cancer, and 48 men with prostate cancer.^[64] Sensitivity, specificity, and positive and negative predictive values for the tests in combination were 94%, 98%, 96%, and 96%, respectively. The limitations of the study included that not all samples were confirmed with a diagnosis, there was a lack of urinary data, and the study results are not generalizable to clinical practice.

Section Summary

Concomitant detection of *TMPRSS2:ERG* and *PCA3* may accurately identify men with prostate cancer. However, estimated accuracy varies across the available studies and clinical utility has not been established.

EXODX™ PROSTATE (INTELLISCORE)

McKiernan (2016) conducted a multicenter validation study of urine exosome *PCA3*, *ERG*, and *SPDEF* RNA expression to predict high-grade (Gleason score ≥ 7) prostate cancer.^[65] The study included men age 50 or older with either a suspicious DRE or PSA levels between 2 and 10 who were scheduled for biopsy. The threshold for a positive test, 15.6, was derived from a training set (n=255) separate from the validation set (n=519). The assay improved on the standard of care alone, with an AUC of 0.73 compared with 0.63 for the standard of care (p<0.001) and 0.62 for the PCPT risk calculator. The sensitivity and specificity for the test were reported to be 97.44% and 27.68%, respectively.

GENE HYPERMETHYLATION FOR PROSTATE CANCER DIAGNOSIS AND PROGNOSIS

Nonrandomized Studies

Several studies have reported associations between DNA hypermethylation at various gene loci (*RASSF1A*, *APC*, *GSTP1*, *PTGS2*, *RAR-beta*, *TIG1*, *AOX1*, *C1orf114*, *GAS6*, *HAPLN3*, *KLF8*, and *MOB3B*) and prostate cancer.^[28,60,66-76] In contrast, some studies have not found evidence of an association.^[77,78] Further, Kachakova (2013) concluded that *HIST1H4K* hypermethylation was more likely due to aging than to prostate carcinogenesis.^[79] ConfirmMDx (MDxHealth) is a commercially available test for gene methylation intended to distinguish true- from false-negative prostate biopsies to avoid the need for repeat biopsy in cases of a true

negative and to identify men who may need a repeat biopsy. The test measures methylation of the genes *GSTP1*, *APC*, and *RASSF1*.

Three blinded multicenter validation studies of the ConfirmMDx test have been performed. Partin (2014) reported on results of the DOCUMENT study; it evaluated archived, cancer-negative prostate biopsy core tissue samples from 350 men from five U.S. urology centers.^[80] All patients underwent repeat biopsy within 24 months. Men with two consecutive negative biopsies were classified as controls and men with a negative biopsy followed by a positive biopsy were classified as cases. Thirty (9%) men were excluded from analysis because of noneligibility (n=2), insufficient DNA (n=1), insufficient biopsy cores (n=23), or detection of adenocarcinoma in the first biopsy based on central pathology review (n=4); 320 men were included in analysis (92 cases, 228 controls). Median age was 62 years (range, not given). Median PSA level was 5.3 ng/mL; 23% of men had PSA levels less than 4 ng/mL and 10% had a PSA level of 10 ng/mL or higher. Sixty percent of men had a normal DRE. Forty-two (13%) of the men were black, 232 (73%) were white, and 13 (4%) were Asian. The ConfirmMDx test, performed on the first biopsy, resulted in a NPV of 88% (95% CI 85% to 91%), sensitivity of 62% (95% CI 51% to 72%), and specificity of 64% (95% CI 57% to 70%). The study was not powered to determine accurately the performance characteristics in a subgroup of black patients, but the estimated sensitivity was 77% (95% CI 46% to 95%), specificity was 66% (95% CI, 46% to 82%), and NPV was 93% (85% CI 82% to 97%). Multivariate analysis of potential predictors of cancer on repeat biopsy, corrected for age, PSA, DRE, first biopsy histopathology characteristics, and race, showed that the ConfirmMDx test was the most significant independent predictor of patient outcome (OR 2.69, 95% CI 1.60 to 4.51).

A study by Waterhouse (2019) was specific to African American men undergoing repeat biopsy within 30 months of a negative index biopsy.^[81] Of the 211 patients in the study, 81 had a positive second biopsy (cases), while 130 had negative results (controls). In this population, the ConfirmMDx had a sensitivity of 74.1% and a specificity of 60.0%. For Gleason score ≥ 7 , the sensitivity and specificity were 78% and 53%, respectively.

The MATLOC study, reported by Stewart (2013), tested archived cancer-negative prostate biopsy needle core tissue samples from 498 men from the U.K. and Belgium.^[82] Patients underwent repeat biopsy within 30 months; cases had a positive second biopsy while controls had a negative second biopsy. A total of 483 men were included in the analysis (87 cases, 396 controls). The median PSA level was 5.9 ng/mL; 21% of men had PSA levels less than 4 ng/mL and 18% had PSA levels of 10 ng/mL or higher. Seventy-three percent of men had benign DRE. The ConfirmMDx test, performed on the first biopsy, resulted in a NPV of 90% (95% CI 87% to 93%), sensitivity of 68% (95% CI 57% to 77%), and specificity of 64% (95% CI 59% to 69%). Multivariate analysis of potential predictors of cancer on repeat biopsy, corrected for patient age, PSA, DRE, and first biopsy histopathology characteristics, showed that the ConfirmMDx test was the most significant independent predictor of patient outcome (OR 3.17, 95% CI 1.81 to 5.53).

Van Neste (2016) reported on results of combined data from the DOCUMENT and MATLOC studies to investigate whether DNA methylation intensities were associated with high-grade (Gleason score, ≥ 7) prostate cancer.^[83] DNA methylation was the most significant and important predictor of high-grade cancer, resulting in an NPV of 96% (precision not reported).

Wojno (2014) reported on a field observation study in which practicing urologists at five centers used the ConfirmMDx test to evaluate at least 40 men with previous cancer-negative biopsies

who were considered at risk for prostate cancer.^[84] Centers reported whether patients who had a negative test assay result had undergone a repeat biopsy at the time of the analysis. Median patient follow-up after the assay results were received was nine months. A total of 138 patients were included in the analysis. The median PSA level was 4.7 ng/mL. Repeat biopsies had been performed in six (4.3%) of the 138 men with a negative ConfirmMDx test, in which no cancer was identified.

Aubry (2013) analyzed the expected reduction in biopsies associated with ConfirmMDx use.^[85] Using the MATLOC estimates of performance characteristics for ConfirmMDx, the authors estimated that 1,106 biopsies per one million people would be avoided. The study did not include decision analysis comparing the tradeoff in reduction in biopsies and missed cancers.

Section Summary

Studies evaluating the diagnostic accuracy and predictive ability of gene hypermethylation report differing results regarding the accuracy of hypermethylation. These inconsistent results make it difficult to determine whether hypermethylation is a useful parameter for diagnosis and/or prognosis of prostate cancer. Two clinical validation studies have reported on the clinical validity of the ConfirmMDx score in the intended use population. The studies did not provide estimates of validity compared with a standard clinical examination with percent free PSA. No data are available on the long-term clinical outcomes or clinical utility of the test. The indirect chain of evidence is incomplete due to the limitations of evidence on the comparative clinical validity and utility.

MITOCHONDRIAL DNA VARIANT TESTING

Variants in the mitochondrial genome (mtDNA) are emerging as tools for the diagnosis of prostate cancer. A growing body of literature is reporting significant associations between both single nucleotide changes and large-scale deletions in mtDNA and prostate cancer, however, the identified studies are of small to medium sample size and do not address clinical utility. A laboratory developed assay offered by Mitomics (formerly Genesis Genomics), called the Prostate Core Mitomics Test™ (PCMT), is a proprietary test which is intended to determine whether a patient has prostate cancer, despite a negative prostate biopsy, by analyzing deletions in mitochondrial DNA by PCR to detect “tumor field effect”. The test is performed on the initial negative prostate biopsy tissue. According to the company website, a negative PCMT result confirms the results of the negative biopsy (i.e., the patient doesn’t have prostate cancer) and that the patient can avoid a second biopsy, but that a positive PCMT means that the patient is at high risk of undiagnosed prostate cancer. The company website states that the sensitivity of the test is 85% and has a negative predictive value of 92%.

Nonrandomized Studies

A study published by Giorgi (2016) examined the role of the mitochondrial genome in prostate cancer risk in 4,086 prostate cancer cases and 3,698 controls from the Multiethnic Cohort.^[86] In this study, 350 mitochondrial SNVs were tested in five racial/ethnic populations: Asian Americans, Africans, Europeans, Latinos, and Native Hawaiians. No significant associations were found.

Legisi (2016) queried a pathology services database to identify (1) men who had a negative initial prostate biopsy and a negative PCMT (n=644), and (2) men who had a negative initial prostate biopsy and a repeat biopsy (n=823).^[87] Of the 644 patients with a negative PCMT, 35

had a repeat biopsy and five (14.2%) were false-negatives who were found to have cancer on rebiopsy. The number of false-negatives among patients who did not have a repeat biopsy cannot be determined from this study. Of the second group of 823 men who had a repeat biopsy, 132 had a PCMT. Changes in physician decisionmaking led to earlier detection of prostate cancer by 2.5 months and an increase in cancer detection rates, but this was only observed when men with atypical small acinar proliferation on index biopsy were not included. Interpretation of these results is limited because testing was not random or consecutive.

Published literature from Genesis Genomics on the use of mitochondrial DNA variants in prostate includes several studies. A 2006 study retrospectively analyzed mitochondrial DNA variants from three tissue types from 24 prostatectomy specimens: prostate cancer, adjacent benign tissue and benign tissue distant to the tumor (defined as tissue from a nondiseased lobe or at least 10 cell diameters from the tumor if in the same lobe).^[88] Prostate needle biopsy tissue from 12 individuals referred for biopsy that were histologically benign were used as controls. Results from the prostatectomy tissue analysis showed that 16 of 24 (66.7%) had variants in all three tissue types, 22 of 24 (91.7%) had variants in malignant samples, 19 of 24 (79.2%) in adjacent benign samples and 22 of 24 in distant benign glands. Overall, 273 somatic variants were observed in this sample set. In the control group, seven (58.3%) patients were found to have between one and five alterations, mainly in non-coding regions. The authors concluded that the variants found in the malignant group versus the control group were significantly different and that mitochondrial DNA variants are an indicator of malignant transformation in prostate tissue.

Maki (2008) reported the discovery and characterization of a 3.4-kb mitochondrial genome deletion and its association with prostate cancer^[89]. A pilot study analyzed 38 benign biopsy specimens from 22 patients, 41 malignant biopsy specimens from 24 patients and 29 proximal to malignant (PTM) biopsy specimens from 22 patients. All of the patients with malignant biopsies had a subsequent prostatectomy, and the diagnosis of cancer was confirmed. The PTM biopsy samples were negative for cancer and were from the cohort who underwent prostatectomy. A confirmation study used 98 benign biopsy specimens from 22 patients, 75 malignant biopsy specimens from 65 patients, and 123 PTM biopsy specimens from 96 patients. In the confirmation study, patients were required to have at least two successive negative biopsies; the first negative biopsy was used for analyses. For both the pilot and confirmation studies, samples for analysis were selected based upon review of pathology reports. The levels of the variant were measured by quantitative PCR and using PCR cycle threshold data were used to calculate a score for each biopsy sample. In the pilot study, the scores were statistically significant between benign and malignant ($p < 0.0001$) and benign and proximal ($p < 0.003$) samples. The PTM samples closely resembled the malignant sample, with no statistically significant resolution between their scores ($p < 0.833$), to which the authors attributed as a field cancerization phenomenon. Results from the larger confirmation study were similar. Compared with histopathologic examination of the benign and malignant samples, the sensitivity and specificity were 80% and 71%, respectively, and the area under a ROC curve was 0.83 for the deletion. A blinded, external validation study showed a sensitivity and specificity of 83% and 79% and an AUC of 0.87.

Robinson (2010) assessed the clinical value of the 3.4-kb deletion described in the Maki (2008) study in predicting re-biopsy outcomes^[89]. Levels of the deletion were measured by quantitative PCR in prostate biopsies negative for cancer from 101 patients who underwent repeat biopsy within a year and had known outcomes. Of the 101 first biopsies, the diagnosis was normal in eight, atypical and/or had prostatic intraepithelial neoplasia (PIN) in 50 and

hyperplasia or inflammation in 43. Using an empirically established cycle threshold cutoff, the lowest cycle threshold as diagnostic of prostate cancer, and the histopathologic diagnosis on second biopsy, the clinical performance of the deletion was calculated. The final data was based on 94 patients, who on second biopsy had 20 malignant and 74 benign diagnoses. The cycle cutoff gave a sensitivity and specificity of 84% and 54%, respectively, with AUC of 0.75. Negative predictive value was 91%.

Section Summary

Studies using the PCMT test for the diagnosis of prostate cancer reported sensitivities and specificities in the moderate range. In general, these studies are preliminary and only report on clinical validity. There is a lack of standardization in methodology and the clinical utility of this test was not addressed.

GENE PANELS

A three-gene panel (*HOXC6*, *TDRD1*, *DLX1*) developed by Leyten (2015) is now commercially available as SelectMDx.^[90] Xiao (2016) reported the development of an eight-gene panel (*PMP22*, *HPN*, *LMTK2*, *FN1*, *EZH2*, *GOLM1*, *PCA3*, *GSTP1*) that distinguished high-grade prostate cancer from indolent prostate cancer with a sensitivity of 93% and NPV of 61%.^[91] Validation of this panel is needed.

GENE EXPRESSION ANALYSIS

Oncotype Dx[®] Prostate

Analytic Validity

In the only study of validity for the Oncotype DX[®] Prostate test, Knezevic (2013) reported analytical accuracy and reproducibility of Oncotype Dx[®] Prostate.^[92] Estimates of analytic precision and reproducibility were derived from analysis of RNA prepared from 10 microdissected prostate tumor samples obtained by needle biopsy. Individual Gleason scores were assigned using the 2005 International Society of Urological Pathology Consensus guidelines.^[93]

The results showed that the assay could accurately measure expression of the 12 cancer-related and five reference genes over a range of absolute RNA inputs (0.005 to 320 ng); the limit of detection in a sample was 0.5 ng/uL. The analytic accuracy showed average variation of less than 9.7% across all samples at RNA inputs typical of needle biopsy specimens. The amplification efficiency for the 17 genes in the test ranged from 88% to 100%, with a median of 93% (standard deviation [SD] 6%) for all 17 genes in the assay. Analytic precision was assessed by examining variability between replicate results obtained using the same mRNA input. Reproducibility was measured by calculating both within and between mRNA input variation. A low input level of 5 ng mRNA was used to reflect the lowest 2.5 percentile of a tumor sample of 0.023 cm³. When converted to Genomic Prostate Score (GPS) units (unit measure for reporting test results), the SD for analytic precision was 1.86 GPS units (95% CI 1.60 to 2.20) on the 100-unit scale. The SD for reproducibility was 2.11 GPS units (95% CI 1.83 to 2.50) on the 100-point scale. This study provides sufficient evidence to establish the analytic validity of Oncotype Dx[®] Prostate.

Clinical Validity

Van Den Eeden (2018) reported on a retrospective study using a stratified cohort sampling design including 279 of 6,184 men who were diagnosed with prostate cancer within a registry between 1995 and 2010 and underwent radical prostatectomy (RP) within 12 months of diagnosis, with a median followup of 9.8 years.^[94] In an analysis adjusted for NCCN risk classifications, the GPS was associated with BCR-free survival (HR 2.1, 95% CI 1.4 to 3.1), distant metastasis (HR 2.3, 95% CI 1.4 to 3.9), and prostate cancer death following RP (HR 2.7, 95% CI 1.5 to 4.8). Ten-year prostate cancer death by GPS score was displayed in a figure stratified by NCCN risk classification, which provides some information on potential reclassification. Ten-year prostate cancer death appears to be close to 0 for men who are NCCN low-risk regardless of GPS score, indicating little useful reclassification of NCCN low-risk men based on GPS. For NCCN intermediate risk, the risk of prostate cancer death ranges from approximately zero for a GPS of less than 40 to close to 40% for a GPS of 100. It is unclear how many men with GPS less than 40 were NCCN favorable intermediate risk.

Brand (2016) combined studies from Klein (2014) and Cullen (2015) using a patient-specific meta-analysis.^[95] The GPS was compared to the CAPRA score, NCCN risk group, and AUA/EAU risk group. The authors tested whether the GPS added predictive value for the likelihood of favorable pathology above the clinical risk assessment tools. The model including the GPS and CAPRA score provided the best risk discrimination; the AUC improved from 0.68 to 0.73 by adding the GPS to CAPRA score. The AUC improved from 0.64 to 0.70 by adding the GPS to the NCCN risk group. The improvements were reported to be significant but the confidence intervals for AUC were not provided.

Whalen (2016) prospectively evaluated the correlation of GPS with final pathology at RP in a clinical practice setting. Eligible men were 50 years of age and older with more than 10 years of life expectancy, PSA levels of 20 ng/mL or less, stage cT1c-cT2c newly diagnosed, untreated prostate cancer, and who met NCCN classifications as very low risk, low risk, or low-intermediate risk.^[96] Men were enrolled from May 2013 to August 2014 at an academic medical center. Genomic Health reclassified patients' cancers as "less favorable," "consistent with," or "more favorable" than what would have been predicted by their NCCN risk group. The primary outcome was adverse pathology at RP defined as any pT3 stage and primary Gleason grade of 4 or any pattern 5. Fifty patients had RP pathology and the reclassification results for these participants are discussed here; 21 (42%) met the definition of adverse pathology. The NCCN risk classification categorized two (4%) patients as very low risk, 34 (68%) as low risk, and 14 (28%) as low-intermediate risk. Twenty-three (46%) of patients were reclassified using GPS. Confidence intervals were not provided.

One publication compiled results of three cohorts, two of which were used for test development, of contemporary (1997 to 2011) patients in a prostatectomy study (n=441; Cleveland Clinic database, 1987 to 2004), a biopsy study (n=167; Cleveland Clinic database, 1998 to 2007), and an independent clinical validation study cohort (n=395; mean age, 58 years; University of California, San Francisco Urologic Oncology Data Base, 1998 to 2011).^[97]

A study by Salmasi (2018) examined the ability of the test to predict adverse pathology in 134 patients with NCCN very low, low, or intermediate risk prostate cancer, who had additionally undergone MRI testing.^[98] MRI results were reported by both UCLA score and Pi-RADSv2. In contrast to the results of other studies, the PSA and MRI scores in this group were not independent predictors of adverse pathology, while the GPS was (OR 3.3, 95% CI 1.74 to 6.62, p<0.001),

Results from the clinical validation study and prostatectomy study provide information on the potential clinical validity of this test. The cohorts had a mix of low to low-intermediate clinical risk characteristics using National Comprehensive Cancer Network (NCCN) or American Urological Association (AUA) criteria.

The clinical validation study was prospectively designed, used masked review of prostatectomy pathology results, and as such met the REMARK (Reporting Recommendations for Tumor Marker Prognostic Studies) guidelines for biomarker validation.^[99] The prostatectomy study used a case-cohort design to select a 1:3 ratio of recurrent to nonrecurrent patients. The prespecified primary endpoint of the validation study was the ability of the GPS to predict the likelihood of favorable pathology in the needle biopsy specimen. Favorable pathology was defined as freedom from high-grade or non-organ-confined disease. In the prostatectomy study, the ability of the GPS to further stratify patients within AUA groupings was related to clinical recurrence-free interval in regression-to-the-mean estimated survival curves.

The validation study results showed that the GPS could refine stratification of patients within specific NCCN criteria groupings, as summarized in Table 1. The proportions in Table 1 were estimated from a plot of GPS versus the percent likelihood of favorable pathology.^[100]

Table 1. Reclassification of Prostate Cancer Risk Categories with Oncotype Dx® Prostate^[100]

NCCN Risk Level	Estimated Mean Likelihood of Favorable Tumor Pathology	
	NCCN Criteria, %	GPS + NCCN, % Range
Very low	≈84	63-91
Low	≈76	55-86
Intermediate	≈56	29-75

GPS: Genomic Prostate Score; NCCN: National Comprehensive Cancer Network.

The actual number of patients correctly or incorrectly reclassified between all three categories cannot be ascertained from the data provided. The results suggested that the combination of GPS plus clinical criteria could reclassify patients on an individual basis within established clinical risk categories. However, whether these findings support a conclusion that the GPS could predict the disease-specific survival based solely on the level of pathology in a biopsy specimen is unclear. Moreover, extrapolation of this evidence to a true active surveillance population, for which the majority in the study would be otherwise eligible, is difficult because all patients had elective radical prostatectomy within 6 months of diagnostic biopsy.

The Klein (2014) prostatectomy study, although used to identify genes to include in the GPS, provided estimates of clinical recurrence rates stratified by AUA criteria (Epstein), compared with rates after further stratification according to the GPS from the validation study. The survival curves for clinical recurrence reached a duration of nearly 18 years based on the dates individuals in the cohort were entered into the database (1987-2004). The reclassifications are summarized in Table 2. The GPS groups are grouped by tertiles defined in the overall study. These data suggest the GPS can reclassify a patient’s risk of recurrence based on a specimen obtained at biopsy. However, the findings do not necessarily reflect a clinical scenario of predicting disease progression in untreated patients under active surveillance.

Table 2. Reclassification of Prostate Cancer 10-Year Clinical Recurrence Risk with Oncotype Dx® Prostate

Overall 10-Year Risk, % (AUA Risk Level)	10-Year Risk, % (GPS Low Group)	10-Year Risk, % (GPS Intermediate Group)	10-Year Risk, % (GPS High Group)
3.4 Low	2.0	3.4	7.0
9.6 (Intermediate)	2.8	5.1	14.3
18.2 (high)	6.2	9.2	28.6

AUA: American Urological Association; GPS: Genomic Prostate Score.

A retrospective cohort study by Cullen (2015) included men with NCCN-defined very low-through intermediate-risk prostate cancer undergoing RP within six-months of diagnosis.^[101] The sample was obtained from men enrolled in the Center for Prostate Disease Research longitudinal study at two U.S. military medical centers. A Gleason score of 4 or 5 with non-organ-confined disease was considered adverse pathology. Biopsies were available for 500 (57.9%) of 864 eligible patients; 382 (44.2% of eligible) with both adequate tissue for gene expression analysis and available RP pathology were included in the analysis. Selected patients were older (61.0 years vs. 59.7 years, $p=0.013$) and had both higher Gleason scores ($p<0.001$) and NCCN risk classification (29.8% vs 32.9% intermediate, $p=0.035$). Median follow-up was 5.2 years and biochemical recurrence (BCR) occurred in 62 (15.4%). Adverse pathology was noted in 163 (34%) men. In an analysis adjusted for baseline characteristics, the GPS was associated with BCR-free survival (HR 2.73 for each 20-point increase, 95% CI 1.84 to 3.96) and adverse pathology following RP (HR 3.23 per 20-point increase, 95% CI 2.17 to 4.97). The GPS improved the C statistic for adverse pathology over NCCN risk alone from 0.63 to 0.72 (CIs not reported). Comparisons with other predictors such as CAPRA or Gleason score alone were not reported. Study implications were limited by the low proportion of eligible men in the analysis and differences between excluded and included men.

Clinical Utility

Klein (2014) also reported a decision-curve analysis that they have proposed reflects the clinical utility of Oncotype Dx® Prostate.^[100] The analysis investigated the predictive impact of the GPS in combination with the Cancer of the Prostate Risk Assessment (CAPRA) validated tool^[102] versus the CAPRA score alone on the net benefit for the outcomes of patients with high-grade disease (Gleason >4+3), high-stage disease, and combined high-grade and high-stage disease. They reported that, over a range of threshold probabilities for implementing treatment, “incorporation of the GPS would be expected to lead to fewer treatments of patients who have favorable pathology at prostatectomy without increasing the number of patients with adverse pathology left untreated.” For example, at a threshold risk of 40% (e.g., a man weighing the harms of prostatectomy versus benefit over active surveillance at 4:6) the test could identify 2 per 100 with high-grade or high-stage disease at a fixed false positive rate compared with using the CAPRA score alone. However, no confidence intervals were presented for the decision curve analysis. Thus, an individual patient could use the findings to assess his balance of benefits and harms (net benefit) when weighing the choice to proceed immediately to curative radical prostatectomy with its attendant adverse sequelae, or to enter an active surveillance program. The latter would have an immediate benefit realized by forgoing radical prostatectomy, but perhaps would be associated with greater downstream risks of disease progression and subsequent therapies.

Section Summary

No direct evidence of clinical utility was found. Klein's decision-curve analyses suggest a potential ability of the combined GPS and CAPRA data to help patients make decisions based on relative risks associated with immediate treatment or deferred treatment (i.e., active surveillance). This would reflect the clinical utility of the test. However, it is difficult to ascribe possible clinical utility of Oncotype Dx[®] Prostate in active surveillance because all patients regardless of clinical criteria elected radical prostatectomy within six months of diagnostic biopsy. Moreover, the validity of using different degrees of tumor pathology as a surrogate for cancer-specific death is unclear. Reports from validation studies lack precision estimates for important variables such as risk estimates for recurrence.

Prolaris[®]

Analytic Validity

Although there is no reference standard for gene expression profiling tests, other measures of technical performance are relevant and include reproducibility, tissue-sample adequacy, potential batch effects, and test-set bias. Warf (2015) evaluated the precision of the Cell Cycle Progression (CCP) score using six formalin-fixed, paraffin-embedded (FFPE) biopsy (three replicate scores) and 12 FFPE RP (four to six replicate scores) specimens.^[103] Overall precision was estimated from replicate samples, intended to reflect combined variation from tissue dissection through gene expression. Across replicate samples, the standard deviation of the CCP score was 0.1 (95% CI 0.08 to 0.13). After eight weeks of sample storage, results were similar. In 2013, Myriad Genetics reported 95.3% of samples were adequate to produce a CCP score.^[104]

Clinical Validity: Needle Biopsy, Conservative Management

In 2016, results of a systematic review and meta-analysis supported by the manufacturer were reported.^[105] Published and unpublished studies of prognostic validity or clinical utility of CCP testing were eligible for inclusion. Seven published studies were identified; five were clinical validity studies. Two are reviewed in the following paragraphs and the remaining validity studies will be reviewed in a subsequent section on post-RP management. The two "utility" studies are discussed in the following section. Two validity studies reported outcomes for disease-specific mortality^[106,107] but of the two, only the Cuzick (2012)^[106] included newly diagnosed patients, so the pooled outcome is not of relevance in this section.

The more recent study by Cuzick (2015) examined three U.K. cancer registries from 1990 to 2003 to identify men with prostate cancer who were conservatively managed following needle biopsy, with follow-up through December 2012. Men were excluded if they had undergone RP or radiation therapy within six months of diagnosis.^[104] A combination of the CCP and Cancer of the Prostate Risk Assessment (CAPRA) scores was used to predict prostate cancer death. There were 989 men who fit eligibility criteria; CCP scores were calculable for 761 (77%) and combined CCP and clinical variables were available for 585 (59%). Median age at diagnosis was 70.8 years and median follow-up was 9.5 years. The prostate cancer mortality rate was 17% (n=100), with 29% (n=168) dying from competing causes. Higher CCP scores were associated with increased 10-year risk of prostate cancer mortality: 7% (CCP score <0), 15% (CCP score 0 to 1), 36% (CCP score 1 to 2), 59% (CCP score >2). A one-unit increase in CCP was associated with a crude HR for death of 2.08 (95% CI 1.76 to 2.46) and when adjusted for CAPRA score yielded a HR of 1.76 (95% CI 1.47 to 2.14). For the combined CAPRA/CCP score, the HR for 10-year prostate cancer mortality increased to 2.17 (95% CI 1.83 to 2.57). The c-statistic for the CAPRA score was 0.74; adding the CCP score increased the C statistic

to 0.78 (no confidence intervals for the AUC were reported). Treatment changes after six months were documented in only part of one of the three cohorts; at 24 months, 45% of the men in this cohort had undergone radiotherapy or prostatectomy. Therefore, the potential effect of treatment changes on prognostic estimates is uncertain.

Lin (2018) validated a CCR cutoff of 0.8 using a subset of 585 conservatively managed men from the Cuzick (2015) cohort.^[108] Of the 585 men, 60 had CCR scores of 0.8 or less. Among the 284 men who were at low- or intermediate-risk by NCCN criteria, 59 had CCR scores of 0.8 or less. The text reports that the estimated 10-year prostate cancer mortality risk was 2.7% for men with CCR scores below the threshold and 3.3% (95% CI 1.9% to 5.7%) at the threshold in the full cohort, and 2.3% below the threshold and 2.9% (95% CI 1.3% to 6.7%) at the threshold in the cohort that excluded high-risk men. However, the Kaplan-Meier curves show an estimated prostate cancer mortality at 10 years of 0% for men with CCR of 0.8 or less in both cohorts. The Kaplan-Meier curve estimated prostate cancer mortality at 10 years for men with CCR greater than 0.8 was 20% in the full cohort and 9% in the cohort excluding high-risk men.

The Cuzick (2012) study examined the prognostic value of Prolaris[®] for prostate cancer death in a conservatively managed needle biopsy cohort.^[106] Cell cycle expression data were read blind to all other data. Patients were identified from six cancer registries in Great Britain and were included if they had clinically localized prostate cancer diagnosed by needle biopsy between 1990 through 1996; were younger than 76 years at diagnosis; had a baseline PSA measurement; and were conservatively managed. Potentially eligible patients who underwent RP, died, showed evidence of metastatic disease within six months of diagnosis, or received hormone therapy before diagnostic biopsy were excluded. The original biopsy specimens were retrieved and centrally reviewed by a panel of expert urologic pathologists to confirm the diagnosis and, where necessary, to reassign Gleason scores.⁴⁴ Of 776 patients diagnosed by needle biopsy and for which a sample was available to review histology, needle biopsies were retrieved for 527 (68%), 442 (84%) of which had adequate material to assay. From the 442 samples, 349 (79%) produced a CCP score and had complete baseline and follow-up information, representing 45% of 776 patients initially identified. The median follow-up time was 11.8 years. Ninety deaths from prostate cancer occurred within 2799 person-years.

The primary, unadjusted analysis found a one-unit increase in CCP score associated with a two-fold increase (HR 2.02, 95% CI 1.62 to 2.53) in the risk of dying from prostate cancer. In a multivariate model including CCP, Gleason score, and PSA level, the adjusted HR for a one-unit increase in CCP score was 1.65 (95% CI 1.31 to 2.09). However, changes in HRs may not reflect meaningful changes in absolute risk. It appears that there might be a large change in risk for scores below 2 compared with above 2, but no CIs are reported so it is impossible to draw conclusions. Measures that would suggest improved discriminatory ability (e.g., AUC or reclassification) compared with an existing nomogram were not reported. The authors did not provide evidence that the test could correctly reclassify men initially at high risk to lower risk to avoid overtreatment, or conversely, correctly reclassify those initially at low risk to high risk to avoid undertreatment.

Table 3. Univariate and Multivariate Associations between CCP and Death from Prostate Cancer in the Cuzick 2012 and 2015 Validation Studies

Study (year)	N	Unadjusted	Multivariate
		Hazard Ratio (95% CI) for 1-unit increase in CCP	Hazard Ratio (95% CI) for 1-unit increase in CCP
Cuzick (2012) ^[106]	349	2.02 (1.62 to 2.53)	1.65 (1.31 to 2.09) ^a
Cuzick (2015) ^[104]	585	2.08 (1.76 to 2.46)	1.76 (1.47 to 2.14) ^b

CI: confidence interval

CCP: cell cycle progression

^a adjusted for Gleason score and prostate-specific cancer level

^b adjusted for Cancer of the Prostate Risk Assessment

In summary, Table 3 displays the association between CCP score adjusted for CAPRA and Table 4 shows the risk of death by groups of CCP score. The CCR score is most relevant as it appears in the sample report provided by the manufacturer. Table 3 demonstrates an association between CCP and risk of death on the relative scale but does not necessarily indicate that there is a difference in absolute risk that would be meaningful for clinical decision making. Table 4 displays the estimated absolute risk of death for the CCP score but notably absent are CIs which would help in interpretation. However, given the data provided, several concerns arise. Even the lowest risk group shown in Cuzick (2012)^[106] has a 10-year death rate of 20%, which may be explained by the population characteristics (i.e., not PSA screen-selected, a third with Gleason >7 and half with PSA >25); however, a death rate of 20% is unlikely to be low enough to forgo immediate treatment.

Table 4 does not include the death rates by CCR score; however, the predicted 10-year prostate cancer death rates by CCR score were provided in a figure in Cuzick (2015).^[104] The predicted 10-year risk for CAPRA alone compared with CCR was provided in a dot plot. The authors stated that CCR identified 11 men with a CAPRA score of 2 (indicating an estimated 10-year mortality rate of 4%) who “had a higher risk” based on CCR score. From the dot plot, it appears that for these 11 men, the 10-year mortality rate estimated by CCR score ranged from just greater than 4% to about 8%. The authors also indicated that for 31 men with CAPRA of 3 (corresponding to a 10-year risk of death rate of 5.7%) the risk as estimated by CCR was less than 4.0%, from the plot the CCR estimated risk appears to range from about 2.5% to 4% for those 31 men. It is not clear that either of these reclassifications would change estimated mortality enough to alter treatment decisions.

Table 4. Kaplan-Meier Estimates of Prostate Cancer Death at 10 Years According to CCP Score Groupings in the Cuzick Validation Studies

CCP Score	Cuzick (2012) ^[106]		Cuzick (2015) ^[104]	
	N	10-Year Death Rate (%) ^a	N	10-Year Death Rate (%) ^a
≤ 0	36	19.3	194	7
0 to ≤ 1	133	19.8	251	15
1 to ≤ 2	114	21.1	110	36
2 < CCP ≤ 3	50	48.2	30 ^b	59
> 3	16	74.9		

CCP: cell cycle progression

^a Confidence intervals not reported

^b Grouped CCP scores >2

Clinical Validity: Posttreatment (radical prostatectomy and external beam radiation therapy)

Four studies have reported clinical validity in the post-RP management setting. Three of these studies –Cuzick (2011),^[107] Cooperberg (2013),^[97] and Bishoff (2014) –reported on post-RP patients. Koch et al (2016) reported on post-RP patients with BCR. Freedland et al (2013) reported on post-RT patients but is included in this section for completeness.

Bishoff (2014) examined the prognostic ability of the CCP score in three cohorts: Martini Clinic (n=283, simulated biopsies from FFPE RP specimen), Durham Veterans Affairs Medical Center (n=176, diagnostic biopsies) and Intermountain Healthcare (n=123, diagnostic biopsies).^[109] The combined analysis included all 582 patients. Gleason scores were 7 or lower in 93% of men. In the combined cohorts, a unit increase in the CCP score increased the adjusted HR for BCR by 1.47 (95% CI 1.23 to 1.76). Metastatic events (n=12) were too few to draw conclusions. Although the CCP score was associated with increased risk of BCR, the analyses do not allow examining whether the CCP score provides improved discrimination over clinicopathologic variables.

In Myriad-funded study, Freedland (2013) evaluated the CCP score's ability to predict biochemical recurrence (BCR) in a cohort of men treated with external beam radiation therapy (EBRT).^[110] The CCP score was derived retrospectively from diagnostic biopsy specimens of men diagnosed with prostate cancer from 1991 to 2006 (n=141). The primary outcome assessed was time from EBRT to BCR. In a multivariable analysis with Gleason score, PSA, percent positive cores, and androgen deprivation therapy, the HR was 2.11 for a one-unit increase in CCP score (equivalent to a doubling of gene expression) (p=0.034), indicating that CCP provides prognostic information that is not provided by standard clinical parameters. At 10 years post-EBRT, the CCP score was associated with prostate cancer specific mortality (p=0.013). The limitations of this study include small size of the cohort, small number of treatment failures (only 19 patients [13%] had BCR), and short follow-up time. The authors conceded that “definitive conclusions regarding time dependency will require additional studies”.

Cooperberg (2013) sought to evaluate the CCP score in a RP cohort and the incremental improvement over the Cancer of the Prostate Risk Assessment Postsurgical (CAPRA-S) score for predicting BCR employing a prospective-retrospective design (conforming to a PROBE study design).^[97] A prognostic model was developed from the RP cohort described by Cuzick (2011).^[107] The validation cohort was obtained from patients identified from the University of California, San Francisco (UCSF) Urologic Oncology Database. Tissue sufficient to obtain a CCP score was available for 413 men (69% of the 600 eligible samples). Both UCSF and Myriad Genetics performed statistical analyses. In the validation cohort, 95% had Gleason scores of 7 or lower, 16% of samples had positive margins, 4% had seminal vesicle invasion, and 23% had extracapsular extension. BCR occurred in 82 men (19.9%). The unadjusted HR for BCR increased by 2.1 (95% CI 1.6 to 2.9) per unit increase in CCP score. A predictive model for the combined CCP/CAPRA-S developed in the Cuzick (2011)^[107] RP cohort applied to the UCSF cohort obtained an AUC for BCR with CAPRA-S alone of 0.73 increasing to 0.77 for the combined CCP/CAPRA-S.

Cuzick (2011) examined the potential use of the Prolaris® CCP test combined with a clinical score following RP, using a retrospective cohort and the prospective-retrospective design for archived samples.^[107] The study also included a cohort of men with localized prostate cancer detected from specimens obtained during transurethral resection of the prostate, which is not a population of interest here, and so has not been described. Men conservatively managed post RP between 1985 and 1995 were identified from a tumor registry (n=366 with CCP scores,

Scott and White Clinic, in Texas). The primary endpoint was time to biochemical recurrence (BCR) and the secondary end point was prostate cancer death. Myriad Genetics assessed CCP scores blindly. The median age of patients was 68 years and the median follow-up 9.4 years. Gleason scores were 7 or lower in 96%, but margins were positive in 68%. Cancers were clinically staged as T3 in 34%; following RP, 64% was judged pathologic stage T3. CCP score was associated with BCR (adjusted HR 1.77, 95% CI 1.40 to 2.22). Analyses of prostate cancer deaths in the RP cohort were problematic, owing to only 12 (3%) deaths. The clinical score included PSA, stage, positive surgical margins, and Gleason score. The model was optimized using stepwise variable selection (e.g., a development model). The AUC for BCR within five years in the RP cohort was 0.825 for the clinical score and 0.842 for the combined clinical/CCP score. The discriminatory ability of the clinical score is of note. Although the CCP increased the AUC by 2%, whether that improvement might be clinically useful is unclear lacking reclassification or examination of net benefit.

Clinical Utility

One large prospective registry study, funded by Myriad, was recently published that evaluated the impact of the CCP test on treatment decision making for patients newly diagnosed with prostate cancer.^[111] Patients (n=1,206) with newly diagnosed prostate adenocarcinoma had the CCP test performed on initial prostate biopsy tissue. Changes in treatment decision making was tracked using the answers provided by physicians in sequential surveys relative to initial therapy recommendations (before cell cycle progression). The CCP test caused a change in actual treatment in 47.8% of patients, 72.1% of which were reductions and 26.9% of which were increases in treatment. For each clinical risk category there was a significant change in treatment modality (intervention vs nonintervention) before vs after CCP testing (p=0.0002). This study did not report any changes in patient-important outcomes, such as biochemical recurrence, cancer-specific survival or long-term survival. Although this study reported a change of management in a modest percentage of patients, there was no evidence that these changes in management lead to clinically important improvements in health outcomes.

A prospective study by Hu (2018) tracked the usage of several genomic classification tests in patients with prostate cancer and correlated the results with treatment decisions.^[112] However, this study also did not evaluate clinical outcomes.

Two retrospective survey studies that assessed the potential impact of Prolaris® on physicians' treatment decisions.^[113,114] The authors of each study have suggested their findings support the "clinical utility" of the test, based on whether the results would lead to a change in treatment. Although this information may be useful in assessing the potential test uptake by urologists, it does not demonstrate clinical utility in clinical settings. In a decision-curve analysis, Cooperberg^[97] found the CAPRA-S score superior to CCP alone (as well as treat-none or treat-all strategies) in men postprostatectomy. A combined CCP/CAPRA-S predictor appeared only slightly better than CAPRA-S alone for thresholds of approximately 30% or more. For example, at a threshold of 30% (ie, meaning a man would value the harm-to-benefit of treatment such as radiotherapy as 3:7), the combined CCP/CAPRA-S would detect about two more men per 100 likely to experience BCR if the false-positive rate was fixed. However, the lack of confidence intervals for the decision-curve analysis, together with the small difference, is consistent with an uncertain net benefit obtained by adding CCP to the CAPRA-S score.

Decipher® Prostate Biopsy

Two retrospective cohort studies have been published reporting the clinical validity of Decipher® Biopsy in men with newly diagnosed, localized prostate cancer. Berlin (2019) compared NCCN subclassifications with GC scores in 121 patients with intermediate-risk prostate cancer treated with radiation therapy.^[115] The primary and secondary endpoints of the study were biochemical failure (PSA antigen nadir +2 ng/mL) and metastasis. GC score performed better than the NCCN classifications for biochemical failure (HR 1.36, 95% CI 1.09 to 1.71, p=0.007) and metastasis (HR 2.05, 95% CI 1.24 to 4.24, p=0.004), and a combination of the two performed better than either alone (AUC 0.89 vs. 0.86 for GC alone and 0.54 for NCCN classes alone).

A study by Nguyen (2017) included 235 patients who had either RP or first-line radiotherapy (with or without androgen deprivation therapy) between 1987 and 2024, with a median follow-up of six years.^[116] After adjusting for treatment and clinical data, the biopsy Decipher® score was associated with five-year metastasis (HR 1.39 per 0.1 unit increase, 95% CI 1.09 to 1.8). The frequency of metastasis within five years was 4.1%, 7.8%, and 21% for the Decipher® low-, intermediate-, and high-risk groups.

Decipher® Prostate RP

The Decipher Prostate RP test classifies as low-risk those patients who can delay or defer RT after prostatectomy, or as high-risk those who would potentially benefit from early radiation. The GC is a continuous risk score between 0 and 1, with higher risk scores indicating a greater probability of developing metastasis.

Clinical Validity

The clinical validity of the Decipher® genomic classifier (GC) has been reported in studies to predict metastasis, mortality or BCR after RP in patients with postoperative high-risk features like pathologic stage T2 with positive margins, pathologic stage T3 disease or a rising PSA.^[80,117-126] All studies were retrospective and used registry data or clinical records. The development study was a nested case-control design.^[122] Owing to apparent overlap in samples, the number of unique patients in the studies is difficult to ascertain. Many studies were supported by GenomeDx, which offers the Decipher® test; these studies identified multiple authors as company employees.

Four studies,^[119-122] including the test (validation) sample from the development study, examined men observed following radical prostatectomy and undergoing adjuvant or salvage radiotherapy. Median follow-up periods ranged from 6.4 to 16.9 years. The distributions of Gleason scores in the studies varied—from 24.3% to 49.3% with 8 or higher and 0.4% to 15.1% with 6 or lower. Extracapsular extension of the tumor ranged from 42.7% and 72.3% of men across of the studies.

Klein (2016) evaluated the ability of the Decipher® GC to predict metastasis from the prostate needle biopsy diagnostic tumor tissue from 56 men.^[124] Median follow-up time was eight years. In that time, eight patients metastasized and three died of PCa. Decipher® plus NCCN model had an improved c-index of 0.88 (95% CI 0.77 to 0.96) compared to NCCN alone (c-index 0.75, 95% CI 0.64 to 0.87). Using the Cox multivariable analysis, Decipher® was the only significant predictor of metastasis when adjusting for age, preoperative PSA and biopsy Gleason score (HR 1.72 per 10% increase, 95% CI 1.07 to 2.81, p= 0.02).

A study by Freedland (2016) evaluated the Decipher® GC as a predictor of metastasis in a retrospective study of 170 men who received salvage radiation therapy following cancer recurrence after RP.^[127] After a median follow-up of 5.7 years, 12% of the patients were diagnosed with metastases. The GC results were associated with metastasis (HR 1.58 for a 0.1 unit increase in GC, 95% CI 1.16 to 2.17, $p=0.002$) and the c-index for the Decipher® test was 0.85 (95% CI 0.73 to 0.88), compared with 0.63 to 0.65 for the Cancer of the Prostate Risk Assessment Score and Briganti risk models.

Ross (2015) assessed the prognostic accuracy for metastasis through 10 years, excluding men receiving any adjuvant therapy following radical prostatectomy over median follow-up periods of 7.8 and nine years.^[128] The investigators reported a 6.5% five-year cumulative incidence of metastases in men with GC scores of 0.45 or lower, compared with 30.3% in those with scores higher than 0.60. The AUCs for development of metastases was 0.76 for the GC. In addition, it was found that combining the GC with the best clinicopathologic tool improve the AUC. The study did not include a “standard” reclassification table, but did report 10-year cumulative incidence of metastases stratified by GC and CAPRA-S. The GC appeared to discriminate within CAPRA-S categories, added little to a score greater than 5.

Den (2015) reported on the use of the Decipher® genomic classifier (GC) to provide prognostic and predictive information into the development of metastases in men receiving post-RP RT (either three-dimensional conformal or IMRT).^[117] Genomic classifier scores were calculated from 188 men who were identified within the GenomeDx prostate cancer database with pathologic stage T3 or margin-positive prostate cancer and had received post-RP RT at one of two academic centers between 1990 and 2009. The primary endpoint was metastatic disease (regional or distant) documented on computed tomography or bone scan. Adjuvant versus salvage RT was defined by PSA levels of 0.2 ng/mL or less and more than 0.2 ng/mL before initiation of RT. The clinical characteristics of eligible patients included 72% of men with extraprostatic extension, 35% with seminal vesicle invasion, and 78% with positive surgical margins. Twenty-one percent of patients had a Gleason score of 8 or more. Fifty-one percent of patients received adjuvant RT (89% within 12 months of RP) and overall, patients received RT at a median of five months after RP (range 1 to 160 months). Thirty percent of patients received hormonal therapy with RT. Median follow-up after RP and RT was 10 and 8 years, respectively. Cumulative incidence of metastatic disease at five years after RT for low, average, and high GC scores was 0%, 9%, and 29% ($p=0.002$). In a multivariate analysis, GC and pre-RP PSA were independent predictors of metastasis (both $p<0.01$). In the low GC score group (score <0.4) there was no difference in cumulative incidence of metastasis compared with patients who received adjuvant or salvage RT ($p=0.79$), however, for patients with higher GC scores (≥ 0.4), the cumulative incidence of metastasis at five years was 6% for patients treated with adjuvant RT compared to 23% treated with salvage RT ($p<0.01$). The authors concluded that patients with low GC scores are best treated with salvage RT and those with high GC scores with adjuvant RT.

Klein (2014) evaluated whether use of the Decipher® GC test improved accuracy in predicting metastasis within five years following radical prostatectomy (rapid metastasis [RM]).^[80] Participants included 169 patients who underwent radical prostatectomy between 1987 and 2008, of which 15 were RM and 154 were non-RM controls. Metastasis developed between 1.7 and 3.3 years (median 2.3 years). Test performance was evaluated both individually and in combination with clinical risk factors. After adjusting for clinical factors, Decipher® was a significant predictor of RM (OR 1.48, $p=0.018$). Compared to the Stephenson model, the CAPRA-S, and previously reported biomarkers, Decipher® had the highest concordance index

(c-index), with the highest c-index achieved with integration of Decipher® into the Stephenson nomogram.

Karnes (2013) prospectively created a randomly selected subcohort from the same initial 1,010 post-prostatectomy patients in the Cooperberg study.^[121] Patients with metastasis at diagnosis or with any prior treatment for prostate cancer were excluded. A randomly-selected subcohort was created, with genomic data was available for 219 patients. Following radical prostatectomy, the rates of biochemical recurrence (BCR) at three years was 35% and metastasis at five years was 6%. Median genomic classifier scores were consistently higher in patients with metastases at last follow-up (mean 6.7 years). Median genomic classifier scores also increased with higher Gleason scores. The authors concluded that the higher net benefit of genomic-based classifiers suggested increased specificity (i.e., lower false positives) compared with clinical-only risk models. Because patients with intermediate risk tumors may progress to advanced disease, the authors recommended further study of genomic classifiers in randomized datasets to determine whether genomic classifier scores from diagnostic biopsy specimens can predict metastasis as well as postoperative specimens. A possible limitation of this study was that nearly 15% of patients were node-positive and 45% received adjuvant therapy. Whether the genomic classifier predicted benefit from local (i.e., radiation) or systemic (e.g., hormone) therapies could not be determined because patients were not randomized to these treatments.

Den (2014) reported that within a Decipher® low-risk group that was treated post-RP with RT, there was no difference in oncologic outcomes (either biochemical failure or metastasis) whether they received adjuvant or salvage RT.^[118] For the men classified as high-risk by Decipher, a median four-year PSA-free survival advantage was observed in the patients that received adjuvant versus salvage RT. Of these men classified as high-risk by GC, those who received adjuvant radiation had a 3% cumulative incidence of metastases as compared with 23% incidence of metastasis by eight years in those who delayed treatment and received salvage radiation.

Additional, smaller nonrandomized studies have found associations between GC score and metastatic lymph node involvement^[129] and in recurrence after robotic-assisted laparoscopic prostatectomy.^[130]

Ross (2016) reported results of a retrospective, comparative study of RT after RP for 422 men with pT3 disease or positive margins.^[131] The men were from four cohorts previously described (Karnes [2013], Den [2014], Ross [2016], Freedland [2016]). The four treatment groups were adjuvant RT (n=111), minimal residual disease salvage RT (n=70), salvage RT (n=83), and no RT (n=157). The primary endpoint was metastasis. Thirty-seven men developed metastasis and the median follow-up was eight years. Both CAPRA-S (HR 1.39, 95% CI 1.18 to 1.62) and Decipher® (HR 1.28, 95% CI 1.08 to 1.52) were independently associated with metastasis in multivariable analysis. There was no evidence that treatment effect was dependent on genomic risk (interaction p=0.16 for CAPRA-S, p=0.39 for Decipher), Men with low CAPRA-S or low Decipher® scores had a low risk of metastatic events regardless of treatment selection and men with high CAPRA-S or Decipher® scores benefitted from adjuvant RT compared to the other treatments.

Clinical Utility

Several studies have compared physician's treatment recommendations before and after receiving GC results.^[112,132-137] Because the studies did not include information on outcomes

and clinical validity has not been established, it is not known whether these treatment decisions represent a clinical improvement in management.

DECIPHER® BLADDER

The Decipher® Bladder genomic subtyping classifier (GSC) uses gene expression information to classify muscle-invasive bladder cancer by subtype, which may help identify patients more likely to benefit from neoadjuvant chemotherapy prior to radical cystectomy. The development of the test was described in a study by Seiler (2017), which used the results of whole transcriptome profiling in samples from 343 patients with muscle-invasive bladder cancer prior to chemotherapy to classify tumors into the following subtypes: claudin-low, basal, luminal-infiltrated and luminal.^[138] These subtypes were associated with overall survival, with the basal subtype demonstrating the most survival benefit with neoadjuvant chemotherapy compared with surgery alone.

Additional retrospective studies have linked various GSC results with outcomes in patients with bladder cancer,^[139,140] but these findings require validation, ideally in prospective studies that evaluate health outcomes in patients managed with and without GSC testing.

PRACTICE GUIDELINE SUMMARY

AMERICAN SOCIETY OF CLINICAL ONCOLOGY

The American Society of Clinical Oncology (ASCO) published guidelines on the use of molecular biomarkers in localized prostate cancer (2020), which included the following recommendations:^[141]

- Tissue-based molecular biomarkers (evaluating the sample with the highest volume of the highest Gleason pattern) may improve risk stratification when added to standard clinical parameters, but the Expert Panel endorses their use only in situations in which the assay results, when considered as a whole with routine clinical factors, are likely to affect a clinical decision.
- These assays are not recommended for routine use as they have not been prospectively tested or shown to improve long-term outcomes—for example, quality of life, need for treatment, or survival.

AMERICAN UROLOGICAL ASSOCIATION

The American Urological Association (AUA), American Society for Radiation Oncology, and the Society of Urologic Oncology (2018) published joint guidelines on the management of clinically localized prostate cancer.^[142,143] The guidelines stated that among most low-risk localized prostate cancer patients, genomic biomarkers have not demonstrated a clear role in the selection of active surveillance or in the follow-up of patients on active surveillance.

In 2013 (confirmed in 2018), the AUA published guidelines for the early detection of prostate cancer.^[144] Based on a systematic review of the literature, the guideline panel recognized that novel urinary markers, such as *PCA3* and *TMPRSS2:ERG*, may be “used as adjuncts for informing decisions about the need for a prostate biopsy – or repeat biopsy – after PSA screening,” but emphasized the lack of evidence “that these tests will increase the ratio of benefit to harm.”

The American Urological Association and the Society of Abdominal Radiology published a joint

consensus statement in 2016 on prostate magnetic resonance imaging (MRI) and MRI-targeted biopsy.^[145] The Association recommended:

“In patients with negative or low suspicion magnetic resonance imaging (PI-RADS [Prostate Imaging Reporting and Data System] assessment category of 1 or 2, respectively), other ancillary markers (ie PSA, PSAD, PSAV, PCA3, PHI, 4K) may be of value in identifying patients warranting repeat systematic biopsy, although further data are needed on this topic.”

EVALUATION OF GENOMIC APPLICATIONS IN PRACTICE AND PREVENTION (EGAPP)

In 2013, the EGAPP Working Group published the following recommendations for *PCA3* testing in prostate cancer, based on an AHRQ comparative effectiveness systematic review summarized above:^[146]

- Evidence was insufficient to recommend *PCA3* testing to inform decisions for when to re-biopsy previously biopsy-negative patients for prostate cancer, or to inform decisions to conduct initial biopsies for prostate cancer in at-risk men (e.g., previous elevated PSA or suspicious digital rectal examination).
- Evidence was insufficient to recommend *PCA3* testing in men with cancer-positive biopsies to determine if the disease is indolent or aggressive in order to develop an optimal treatment plan.
- The overall certainty of clinical validity to predict the diagnosis of prostate cancer using *PCA3* is deemed “low.” Clinical use for diagnosis is discouraged unless further evidence supports improved clinical validity.
- The overall certainty of net health benefit is deemed “low.” Clinical use is discouraged unless further evidence supports improved clinical outcomes.

NATIONAL COMPREHENSIVE CANCER NETWORK

Prostate Cancer

The National Comprehensive Cancer Network (NCCN) guidelines for Prostate Cancer Early Detection suggest considering tests that may improve risk prediction.^[147] *PCA3* is not recommended for use prior to biopsy. The guidelines note:

“Biomarkers that improve the specificity of detection are not, as yet, mandated as first-line screening tests. However, there may be some patients who meet PSA standards for consideration of prostate biopsy, but for whom the patient and/or the physician wish to further define risk. Free PSA may improve cancer detection. The probability of higher-grade cancer (Gleason score $\geq 3+4$, Grade Group 2 or higher) may be further defined utilizing the Prostate Health Index (PHI), SelectMDx, 4Kscore, and ExoDx Prostate Test. *PCA3* score is potentially informative after a negative biopsy. Extent of validation of these tests across diverse populations is variable. It is not known how such tests could be applied in optimal combination with MRI as of yet.”

The NCCN guidelines for prostate cancer provide a table of tissue-based tests for prostate cancer prognosis.^[148] The guidelines state that:

“Men with low or favorable intermediate-risk disease and life expectancy ≥ 10 y may consider the use of the following tumor-based molecular assays: Decipher, Oncotype DX Prostate,

Prolaris, and ProMark. Men with unfavorable intermediate- and high-risk disease and life expectancy ≥ 10 may consider the use of Decipher and Prolaris tumor-based molecular assays. Retrospective studies have shown that molecular assays performed on prostate biopsy or radical prostatectomy (RP) specimens provide prognostic information independent of NCCN or CAPRA risk groups. These include, but are not limited to, likelihood of death with conservative management, likelihood of biochemical progression after RP or external beam therapy, and likelihood of developing metastasis after RP or salvage radiotherapy.

The guidelines additionally recommend the Decipher assay to inform adjuvant treatment if adverse features are found after RP, noting that evidence supports adjuvant or salvage radiotherapy for most men with adverse pathological features or detectable PSA who have no evidence of metastasis.

Bladder Cancer

The NCCN guidelines for bladder cancer do not discuss the use of genomic subtyping classifiers.^[149]

U.S. PREVENTIVE SERVICES TASK FORCE RECOMMENDATIONS

The U.S. Preventive Services Task Force published recommendations for Prostate Cancer Screening on May 2012. Genetic tests addressed in this policy, including *PCA3*, were not mentioned.

SUMMARY

There is not enough research to recommend using gene-based tests for prostate or bladder cancer screening, detection and management, as many important characteristics of these tests have not yet been determined. Some research shows that they might help predict the diagnosis or prognosis of prostate cancer, but it is not yet known how much information they add to currently available tests. More research is needed to demonstrate how these tests can improve outcomes for patients. Therefore, use of gene-based testing for screening, detection, and management of prostate or bladder cancer is considered investigational.

REFERENCES

1. U.S. Food and Drug Administration. Premarket Approval (PMA): PROGENSA PCA3 ASSAY. [cited 09/14/2020]; Available from: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P100033>
2. The Public Health Evidence for FDA Oversight of Laboratory Developed Tests: 20 Case Studies. [cited 09/14/2020]; Available from: <http://wayback.archive-it.org/7993/20171114205911/https://www.fda.gov/AboutFDA/ReportsManualsForms/Reports/ucm472773.htm>
3. den Dunnen, JT, Dalgleish, R, Maglott, DR, et al. HGVS Recommendations for the Description of Sequence Variants: 2016 Update. *Human mutation*. 2016 Jun;37(6):564-9. PMID: 26931183

4. TEC Assessment 2008. "Special report: recent developments in prostate cancer genetics and genetic testing." BlueCross BlueShield Association Technology Evaluation Center, Vol. 23, Tab 7.
5. TEC Assessment 2013. "Microarray-based Gene Expression Analysis for Prostate Cancer Management." BlueCross BlueShield Association Technology Evaluation Center, Vol. 28, Tab 11.
6. TEC Assessment 2014. "Gene Expression Analysis for Prostate Cancer Management." BlueCross BlueShield Association Technology Evaluation Center, Vol. 29 Tab 9.
7. Lindstrom, S, Schumacher, F, Siddiq, A, et al. Characterizing associations and SNP-environment interactions for GWAS-identified prostate cancer risk markers--results from BPC3. *PLoS one*. 2011;6(2):e17142. PMID: 21390317
8. Gudmundsson, J, Sulem, P, Rafnar, T, et al. Common sequence variants on 2p15 and Xp11.22 confer susceptibility to prostate cancer. *Nat Genet*. 2008 Mar;40(3):281-3. PMID: 18264098
9. Zheng, SL, Sun, J, Wiklund, F, et al. Cumulative association of five genetic variants with prostate cancer. *N Engl J Med*. 2008 Feb 28;358(9):910-9. PMID: 18199855
10. Salinas, CA, Koopmeiners, JS, Kwon, EM, et al. Clinical utility of five genetic variants for predicting prostate cancer risk and mortality. *Prostate*. 2009 Mar 1;69(4):363-72. PMID: 19058137
11. Kim, ST, Cheng, Y, Hsu, FC, et al. Prostate cancer risk-associated variants reported from genome-wide association studies: meta-analysis and their contribution to genetic Variation. *Prostate*. 2010 Dec 1;70(16):1729-38. PMID: 20564319
12. Little, J, Wilson, B, Carter, R, et al. Multigene panels in prostate cancer risk assessment: a systematic review. *Genet Med*. 2016 Jun;18(6):535-44. PMID: 26426883
13. Little, J, Wilson, B, Carter, R, et al. Multigene panels in prostate cancer risk assessment. *Evidence report/technology assessment*. 2012 Jul(209):1-166. PMID: 24423032
14. Ishak, MB, Giri, VN. A systematic review of replication studies of prostate cancer susceptibility genetic variants in high-risk men originally identified from genome-wide association studies. *Cancer Epidemiol Biomarkers Prev*. 2011 Aug;20(8):1599-610. PMID: 21715604
15. Amin Al Olama, A, Kote-Jarai, Z, Schumacher, FR, et al. A meta-analysis of genome-wide association studies to identify prostate cancer susceptibility loci associated with aggressive and non-aggressive disease. *Hum Mol Genet*. 2013;22:408-15. PMID: 23065704
16. Castro, E, Mikropoulos, C, Bancroft, EK, et al. The PROFILE Feasibility Study: Targeted Screening of Men With a Family History of Prostate Cancer. *The oncologist*. 2016 Jun;21(6):716-22. PMID: 27151655
17. Kader, AK, Sun, J, Reck, BH, et al. Potential impact of adding genetic markers to clinical parameters in predicting prostate biopsy outcomes in men following an initial negative biopsy: findings from the REDUCE trial. *European urology*. 2012 Dec;62(6):953-61. PMID: 22652152
18. Ren, S, Xu, J, Zhou, T, et al. Plateau effect of prostate cancer risk-associated SNPs in discriminating prostate biopsy outcomes. *Prostate*. 2013 Dec;73(16):1824-35. PMID: 24037738
19. Tsuchiya, N, Matsui, S, Narita, S, et al. Distinct cancer-specific survival in metastatic prostate cancer patients classified by a panel of single nucleotide polymorphisms of cancer-associated genes. *Genes Cancer*. 2013;4:54-60. PMID: 23946871

20. Lee, D, Shim, SR, Ahn, ST, et al. Diagnostic Performance of the Prostate Cancer Antigen 3 Test in Prostate Cancer: Systematic Review and Meta-analysis. *Clinical genitourinary cancer*. 2020 Mar 14. PMID: 32280028
21. Qin, Z, Yao, J, Xu, L, et al. Diagnosis accuracy of PCA3 level in patients with prostate cancer: a systematic review with meta-analysis. *International braz j urol : official journal of the Brazilian Society of Urology*. 2020 Sep-Oct;46(5):691-704. PMID: 31961625
22. Rodríguez, SVM, García-Perdomo, HA. Diagnostic accuracy of prostate cancer antigen 3 (PCA3) prior to first prostate biopsy: A systematic review and meta-analysis. *Canadian Urological Association journal = Journal de l'Association des urologues du Canada*. 2020 May;14(5):E214-E9. PMID: 31793864
23. Cui, Y, Cao, W, Li, Q, et al. Evaluation of prostate cancer antigen 3 for detecting prostate cancer: a systematic review and meta-analysis. *Scientific reports*. 2016 May 10;6:25776. PMID: 27161545
24. Nicholson, A, Mahon, J, Boland, A, et al. The clinical effectiveness and cost-effectiveness of the PROGENSA(R) prostate cancer antigen 3 assay and the Prostate Health Index in the diagnosis of prostate cancer: a systematic review and economic evaluation. *Health Technol Assess*. 2015 Oct;19(87):i-xxxii, 1-191. PMID: 26507078
25. Bradley, LA, Palomaki, G, Gutman, S, Samson, DJ, Aronson, N. PCA3 Testing for the Diagnosis and Management of Prostate Cancer. *Agency for Healthcare Research and Quality (US)*. 2013; AHRQ Comparative Effectiveness Reviews. Rockville (MD). PMID: 23638486
26. Choudhury, AD, Eeles, R, Freedland, SJ, et al. The role of genetic markers in the management of prostate cancer. *European urology*. 2012 Oct;62(4):577-87. PMID: 22695242
27. Ruiz-Aragon, J, Marquez-Pelaez, S. [Assessment of the PCA3 test for prostate cancer diagnosis: a systematic review and meta-analysis]. *Actas Urol Esp*. 2010 Apr;34(4):346-55. PMID: 20470697
28. Wei, JT, Feng, Z, Partin, AW, et al. Can urinary PCA3 supplement PSA in the early detection of prostate cancer? *J Clin Oncol*. 2014;32:4066-72. PMID: 25385735
29. Leyten, GH, Wierenga, EA, Sedelaar, JP, et al. Value of PCA3 to Predict Biopsy Outcome and Its Potential Role in Selecting Patients for Multiparametric MRI. *Int J Mol Sci*. 2013;14:11347-55. PMID: 23759986
30. Rodon, N, Trias, I, Verdu, M, et al. Diagnostic and predictive value of urine PCA3 gene expression for the clinical management of patients with altered prostatic specific antigen. *Actas Urol Esp*. 2014 Apr;38(3):150-5. PMID: 24099827
31. Chevli, KK, Duff, M, Walter, P, et al. Urinary PCA3 as a Predictor of Prostate Cancer in a Cohort of 3,073 Men Undergoing Initial Prostate Biopsy. *J Urol*. 2013 Dec 11. PMID: 24333241
32. Capoluongo, E, Zambon, CF, Basso, D, et al. PCA3 score of 20 could improve prostate cancer detection: results obtained on 734 Italian individuals. *Clinica chimica acta; international journal of clinical chemistry*. 2014 Feb 15;429:46-50. PMID: 24269853
33. Hagglof, C, Hammarsten, P, Stromvall, K, et al. TMPRSS2-ERG expression predicts prostate cancer survival and associates with stromal biomarkers. *PloS one*. 2014;9:e86824. PMID: 24505269
34. Lin, DW, Newcomb, LF, Brown, EC, et al. Urinary TMPRSS2:ERG and PCA3 in an active surveillance cohort: results from a baseline analysis in the Canary Prostate Active Surveillance Study. *Clin Cancer Res*. 2013;19:2442-50. PMID: 23515404

35. Ankerst, DP, Groskopf, J, Day, JR, et al. Predicting prostate cancer risk through incorporation of prostate cancer gene 3. *J Urol.* 2008 Oct;180(4):1303-8; discussion 8. PMID: 18707724
36. Feibus, AH, Sartor, O, Moparty, K, et al. Clinical Use of PCA3 and TMPRSS2:ERG Urinary Biomarkers in African-American Men Undergoing Prostate Biopsy. *J Urol.* 2016 Oct;196(4):1053-60. PMID: 27140073
37. Fenstermaker, M, Mendhiratta, N, Bjurlin, MA, et al. Risk Stratification by Urinary PCA3 Testing Prior to MRI-US Fusion-Targeted Prostate Biopsy among Men with No Previous History of Biopsy. *Urology.* 2016 Aug 22. PMID: 27562202
38. Gittelman, MC, Hertzman, B, Bailen, J, et al. PCA3 molecular urine test as a predictor of repeat prostate biopsy outcome in men with previous negative biopsies: a prospective multicenter clinical study. *J Urol.* 2013 Jul;190(1):64-9. PMID: 23416644
39. Hansen, J, Auprich, M, Ahyai, SA, et al. Initial prostate biopsy: development and internal validation of a biopsy-specific nomogram based on the prostate cancer antigen 3 assay. *European urology.* 2013 Feb;63(2):201-9. PMID: 22854248
40. Ankerst, DP, Goros, M, Tomlins, SA, et al. Incorporation of Urinary Prostate Cancer Antigen 3 and TMPRSS2:ERG into Prostate Cancer Prevention Trial Risk Calculator. *European urology focus.* 2018 Feb 5. PMID: 29422418
41. Rubio-Briones, J, Casanova, J, Dumont, R, et al. Optimizing prostate cancer screening; prospective randomized controlled study of the role of PSA and PCA3 testing in a sequential manner in an opportunistic screening program. *Actas Urol Esp.* 2014 May;38(4):217-23. PMID: 24169211
42. Schroder, FH, Venderbos, LD, van den Bergh, RC, et al. Prostate cancer antigen 3: diagnostic outcomes in men presenting with urinary prostate cancer antigen 3 scores ≥ 100 . *Urology.* 2014 Mar;83(3):613-6. PMID: 24581524
43. Ochiai, A, Okihara, K, Kamoi, K, et al. Clinical utility of the prostate cancer gene 3 (PCA3) urine assay in Japanese men undergoing prostate biopsy. *BJU international.* 2013 May;111(6):928-33. PMID: 23331404
44. Porpiglia, F, Russo, F, Manfredi, M, et al. The Roles of Multiparametric Magnetic Resonance Imaging, PCA3 and Prostate Health Index-Which is the Best Predictor of Prostate Cancer after a Negative Biopsy? *J Urol.* 2014 Feb 8. PMID: 24518780
45. Porpiglia, F, Cantiello, F, De Luca, S, et al. In-parallel comparative evaluation between multiparametric magnetic resonance imaging, prostate cancer antigen 3 and the prostate health index in predicting pathologically confirmed significant prostate cancer in men eligible for active surveillance. *BJU international.* 2016 Oct;118(4):527-34. PMID: 26350955
46. Busetto, GM, De Berardinis, E, Sciarra, A, et al. Prostate cancer gene 3 and multiparametric magnetic resonance can reduce unnecessary biopsies: decision curve analysis to evaluate predictive models. *Urology.* 2013 Dec;82(6):1355-60. PMID: 24080222
47. Fradet, V, Toren, P, Nguile-Makao, M, et al. Prognostic value of urinary prostate cancer antigen 3 (PCA3) during active surveillance of patients with low-risk prostate cancer receiving 5alpha-reductase inhibitors. *BJU international.* 2018 Mar;121(3):399-404. PMID: 28972698
48. Ruffion, A, Devonec, M, Champetier, D, et al. PCA3 and PCA3-based nomograms improve diagnostic accuracy in patients undergoing first prostate biopsy. *Int J Mol Sci.* 2013;14:17767-80. PMID: 23994838

49. Ruffion, A, Perrin, P, Devonec, M, et al. Additional value of PCA3 density to predict initial prostate biopsy outcome. *World journal of urology*. 2014 Aug;32(4):917-23. PMID: 24500192
50. Merdan, S, Tomlins, SA, Barnett, CL, et al. Assessment of long-term outcomes associated with urinary prostate cancer antigen 3 and TMPRSS2:ERG gene fusion at repeat biopsy. *Cancer*. 2015 Nov 15;121(22):4071-9. PMID: 26280815
51. Yao, Y, Wang, H, Li, B, Tang, Y. Evaluation of the TMPRSS2:ERG fusion for the detection of prostate cancer: a systematic review and meta-analysis. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2014 Mar;35(3):2157-66. PMID: 24142545
52. Leyten, GH, Hessels, D, Jannink, SA, et al. Prospective multicentre evaluation of PCA3 and TMPRSS2-ERG gene fusions as diagnostic and prognostic urinary biomarkers for prostate cancer. *European urology*. 2014 Mar;65(3):534-42. PMID: 23201468
53. Whelan, C, Kawachi, M, Smith, DD, et al. Expressed prostatic secretion biomarkers improve stratification of NCCN active surveillance candidates: performance of secretion capacity and TMPRSS2:ERG models. *J Urol*. 2014 Jan;191(1):220-6. PMID: 23669563
54. Chan, SW, Nguyen, PN, Violette, P, et al. Early detection of clinically significant prostate cancer at diagnosis: a prospective study using a novel panel of TMPRSS2:ETS fusion gene markers. *Cancer medicine*. 2013 Feb;2(1):63-75. PMID: 24133629
55. Vaananen, RM, Lilja, H, Kauko, L, et al. Cancer-associated changes in the expression of TMPRSS2-ERG, PCA3, and SPINK1 in histologically benign tissue from cancerous vs noncancerous prostatectomy specimens. *Urology*. 2014 Feb;83(2):511 e1-7. PMID: 24468524
56. Dal Pra, A, Lalonde, E, Sykes, J, et al. TMPRSS2-ERG status is not prognostic following prostate cancer radiotherapy: implications for fusion status and DSB repair. *Clin Cancer Res*. 2013;19:5202-9. PMID: 23918607
57. Laxman, B, Tomlins, SA, Mehra, R, et al. Noninvasive detection of TMPRSS2:ERG fusion transcripts in the urine of men with prostate cancer. *Neoplasia*. 2006 Oct;8(10):885-8. PMID: 17059688
58. Laxman, B, Morris, DS, Yu, J, et al. A first-generation multiplex biomarker analysis of urine for the early detection of prostate cancer. *Cancer research*. 2008 Feb 1;68(3):645-9. PMID: 18245462
59. Tosoian, JJ, Loeb, S, Kettermann, A, et al. Accuracy of PCA3 measurement in predicting short-term biopsy progression in an active surveillance program. *J Urol*. 2010 Feb;183(2):534-8. PMID: 20006883
60. Tomlins, SA, Aubin, SM, Siddiqui, J, et al. Urine TMPRSS2:ERG fusion transcript stratifies prostate cancer risk in men with elevated serum PSA. *Sci Transl Med*. 2011 Aug 3;3(94):94ra72. PMID: 21813756
61. Sanda, MG, Feng, Z, Howard, DH, et al. Association Between Combined TMPRSS2:ERG and PCA3 RNA Urinary Testing and Detection of Aggressive Prostate Cancer. *JAMA oncology*. 2017 Aug 1;3(8):1085-93. PMID: 28520829
62. Tomlins, SA, Day, JR, Lonigro, RJ, et al. Urine TMPRSS2:ERG Plus PCA3 for Individualized Prostate Cancer Risk Assessment. *European urology*. 2016 Jul;70(1):45-53. PMID: 25985884
63. Salami, SS, Schmidt, F, Laxman, B, et al. Combining urinary detection of TMPRSS2:ERG and PCA3 with serum PSA to predict diagnosis of prostate cancer. *Urologic oncology*. 2013 Jul;31(5):566-71. PMID: 21600800

64. Robert, G, Jannink, S, Smit, F, et al. Rational basis for the combination of PCA3 and TMPRSS2:ERG gene fusion for prostate cancer diagnosis. *Prostate*. 2013 Jan;73(2):113-20. PMID: 22674214
65. McKiernan, J, Donovan, MJ, O'Neill, V, et al. A Novel Urine Exosome Gene Expression Assay to Predict High-grade Prostate Cancer at Initial Biopsy. *JAMA oncology*. 2016 Jul 1;2(7):882-9. PMID: 27032035
66. Eilers, T, Machtens, S, Tezval, H, et al. Prospective diagnostic efficiency of biopsy washing DNA GSTP1 island hypermethylation for detection of adenocarcinoma of the prostate. *Prostate*. 2007 May 15;67(7):757-63. PMID: 17373715
67. Ellinger, J, Albers, P, Perabo, FG, Muller, SC, von Ruecker, A, Bastian, PJ. CpG island hypermethylation of cell-free circulating serum DNA in patients with testicular cancer. *J Urol*. 2009 Jul;182(1):324-9. PMID: 19447423
68. Goessl, C, Muller, M, Heicappell, R, Krause, H, Miller, K. DNA-based detection of prostate cancer in blood, urine, and ejaculates. *Annals of the New York Academy of Sciences*. 2001 Sep;945:51-8. PMID: 11708494
69. Ellinger, J, Bastian, PJ, Jurgan, T, et al. CpG island hypermethylation at multiple gene sites in diagnosis and prognosis of prostate cancer. *Urology*. 2008 Jan;71(1):161-7. PMID: 18242387
70. Sunami, E, Shinozaki, M, Higano, CS, et al. Multimarker circulating DNA assay for assessing blood of prostate cancer patients. *Clinical chemistry*. 2009 Mar;55(3):559-67. PMID: 19131636
71. Van Neste, L, Herman, JG, Otto, G, Bigley, JW, Epstein, JI, Van Criekinge, W. The epigenetic promise for prostate cancer diagnosis. *Prostate*. 2012 Aug 1;72(11):1248-61. PMID: 22161815
72. Trock, BJ, Brotzman, MJ, Mangold, LA, et al. Evaluation of GSTP1 and APC methylation as indicators for repeat biopsy in a high-risk cohort of men with negative initial prostate biopsies. *BJU international*. 2012 Jul;110(1):56-62. PMID: 22077694
73. Ge, YZ, Xu, LW, Jia, RP, et al. The association between RASSF1A promoter methylation and prostate cancer: evidence from 19 published studies. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2014 Apr;35(4):3881-90. PMID: 24353088
74. Moritz, R, Ellinger, J, Nuhn, P, et al. DNA hypermethylation as a predictor of PSA recurrence in patients with low- and intermediate-grade prostate cancer. *Anticancer Res*. 2013;33:5249-54. PMID: 24324057
75. Haldrup, C, Mundbjerg, K, Vestergaard, EM, et al. DNA methylation signatures for prediction of biochemical recurrence after radical prostatectomy of clinically localized prostate cancer. *J Clin Oncol*. 2013;31:3250-8. PMID: 23918943
76. Chen, Y, Li, J, Yu, X, et al. APC gene hypermethylation and prostate cancer: a systematic review and meta-analysis. *Eur J Hum Genet*. 2013;21:929-35. PMID: 23299921
77. Henrique, R, Ribeiro, FR, Fonseca, D, et al. High promoter methylation levels of APC predict poor prognosis in sextant biopsies from prostate cancer patients. *Clin Cancer Res*. 2007 Oct 15;13(20):6122-9. PMID: 17947477
78. Woodson, K, O'Reilly, KJ, Ward, DE, et al. CD44 and PTGS2 methylation are independent prognostic markers for biochemical recurrence among prostate cancer patients with clinically localized disease. *Epigenetics*. 2006 Oct-Dec;1(4):183-6. PMID: 17998819
79. Kachakova, D, Mitkova, A, Popov, E, et al. Evaluation of the clinical value of the newly identified urine biomarker HIST1H4K for diagnosis and prognosis of prostate cancer in

- Bulgarian patients. *Journal of BUON : official journal of the Balkan Union of Oncology*. 2013 Jul-Sep;18(3):660-8. PMID: 24065480
80. Partin, AW, Van Neste, L, Klein, EA, et al. Clinical validation of an epigenetic assay to predict negative histopathological results in repeat prostate biopsies. *J Urol*. 2014 Oct;192(4):1081-7. PMID: 24747657
 81. Waterhouse, RL, Jr., Van Neste, L, Moses, KA, et al. Evaluation of an Epigenetic Assay for Predicting Repeat Prostate Biopsy Outcome in African American Men. *Urology*. 2019 Jun;128:62-5. PMID: 29660369
 82. Stewart, GD, Van Neste, L, Delvenne, P, et al. Clinical utility of an epigenetic assay to detect occult prostate cancer in histopathologically negative biopsies: results of the MATLOC study. *J Urol*. 2013 Mar;189(3):1110-6. PMID: 22999998
 83. Van Neste, L, Partin, AW, Stewart, GD, Epstein, JI, Harrison, DJ, Van Criekinge, W. Risk score predicts high-grade prostate cancer in DNA-methylation positive, histopathologically negative biopsies. *Prostate*. 2016 Sep;76(12):1078-87. PMID: 27121847
 84. Wojno, KJ, Costa, FJ, Cornell, RJ, et al. Reduced Rate of Repeated Prostate Biopsies Observed in ConfirmMDx Clinical Utility Field Study. *American health & drug benefits*. 2014 May;7(3):129-34. PMID: 24991397
 85. Aubry, W, Lieberthal, R, Willis, A, Bagley, G, Willis, SM, 3rd, Layton, A. Budget impact model: epigenetic assay can help avoid unnecessary repeated prostate biopsies and reduce healthcare spending. *American health & drug benefits*. 2013 Jan;6(1):15-24. PMID: 24991343
 86. Giorgi, EE, Li, Y, Caberto, CP, et al. No Association between the Mitochondrial Genome and Prostate Cancer Risk: The Multiethnic Cohort. *Cancer Epidemiol Biomarkers Prev*. 2016 Jun;25(6):1001-3. PMID: 27021046
 87. Legisi, L, DeSa, E, Qureshi, MN. Use of the Prostate Core Mitomic Test in Repeated Biopsy Decision-Making: Real-World Assessment of Clinical Utility in a Multicenter Patient Population. *American health & drug benefits*. 2016 Dec;9(9):497-502. PMID: 28465777
 88. Parr, RL, Dakubo, GD, Crandall, KA, et al. Somatic mitochondrial DNA mutations in prostate cancer and normal appearing adjacent glands in comparison to age-matched prostate samples without malignant histology. *The Journal of molecular diagnostics : JMD*. 2006;8:312-9. PMID: 16825503
 89. Maki, J, Robinson, K, Reguly, B, et al. Mitochondrial genome deletion aids in the identification of false- and true-negative prostate needle core biopsy specimens. *Am J Clin Pathol*. 2008;129:57-66. PMID: 18089489
 90. Leyten, GH, Hessels, D, Smit, FP, et al. Identification of a Candidate Gene Panel for the Early Diagnosis of Prostate Cancer. *Clin Cancer Res*. 2015 Jul 1;21(13):3061-70. PMID: 25788493
 91. Xiao, K, Guo, J, Zhang, X, et al. Use of two gene panels for prostate cancer diagnosis and patient risk stratification. *Tumour biology : the journal of the International Society for Oncodevelopmental Biology and Medicine*. 2016 Aug;37(8):10115-22. PMID: 26820133
 92. Knezevic, D, Goddard, AD, Natraj, N, et al. Analytical validation of the Oncotype DX prostate cancer assay - a clinical RT-PCR assay optimized for prostate needle biopsies. *BMC genomics*. 2013 Oct 8;14:690. PMID: 24103217
 93. Epstein, JI, Allsbrook, WC, Jr., Amin, MB, Egevad, LL. The 2005 International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading of Prostatic Carcinoma. *Am J Surg Pathol*. 2005;29:1228-42. PMID: 16096414

94. Van Den Eeden, SK, Lu, R, Zhang, N, et al. A Biopsy-based 17-gene Genomic Prostate Score as a Predictor of Metastases and Prostate Cancer Death in Surgically Treated Men with Clinically Localized Disease. *European urology*. 2018 Jan;73(1):129-38. PMID: 28988753
95. Brand, TC, Zhang, N, Crager, MR, et al. Patient-specific Meta-analysis of 2 Clinical Validation Studies to Predict Pathologic Outcomes in Prostate Cancer Using the 17-Gene Genomic Prostate Score. *Urology*. 2016 Mar;89:69-75. PMID: 26723180
96. Whalen, MJ, Hackert, V, Rothberg, MB, McKiernan, JM, Benson, MC, Badani, KK. Prospective correlation between likelihood of favorable pathology on the 17-Gene Genomic Prostate Score and actual pathological outcomes at radical prostatectomy. *Urol Pract*. 2016 Sep;3(5):379-86. PMID:
97. Cooperberg, MR, Simko, JP, Cowan, JE, et al. Validation of a cell-cycle progression gene panel to improve risk stratification in a contemporary prostatectomy cohort. *J Clin Oncol*. 2013;31:1428-34. PMID: 23460710
98. Salmasi, A, Said, J, Shindel, AW, et al. A 17-Gene Genomic Prostate Score Assay Provides Independent Information on Adverse Pathology in the Setting of Combined Multiparametric Magnetic Resonance Imaging Fusion Targeted and Systematic Prostate Biopsy. *J Urol*. 2018 Mar 7. PMID: 29524506
99. McShane, LM, Altman, DG, Sauerbrei, W, Taube, SE, Gion, M, Clark, GM. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol*. 2005;23:9067-72. PMID: 16172462
100. Klein, EA, Cooperberg, MR, Magi-Galluzzi, C, et al. A 17-gene assay to predict prostate cancer aggressiveness in the context of Gleason grade heterogeneity, tumor multifocality, and biopsy undersampling. *European urology*. 2014 Sep;66(3):550-60. PMID: 24836057
101. Cullen, J, Rosner, IL, Brand, TC, et al. A Biopsy-based 17-gene Genomic Prostate Score Predicts Recurrence After Radical Prostatectomy and Adverse Surgical Pathology in a Racially Diverse Population of Men with Clinically Low- and Intermediate-risk Prostate Cancer. *European urology*. 2015 Jul;68(1):123-31. PMID: 25465337
102. Cooperberg, MR, Broering, JM, Carroll, PR. Risk assessment for prostate cancer metastasis and mortality at the time of diagnosis. *Journal of the National Cancer Institute*. 2009;101:878-87. PMID: 19509351
103. Warf, MB, Reid, JE, Brown, KL, et al. Analytical validation of a cell cycle progression signature used as a prognostic marker in prostate cancer. *J Mol Biomark Diagn*. 2015;6(4). PMID:
104. Cuzick, J, Stone, S, Fisher, G, et al. Validation of an RNA cell cycle progression score for predicting death from prostate cancer in a conservatively managed needle biopsy cohort. *British journal of cancer*. 2015 Jul 28;113(3):382-9. PMID: 26103570
105. Sommariva, S, Tarricone, R, Lazzeri, M, Ricciardi, W, Montorsi, F. Prognostic value of the Cell Cycle Progression Score in patients with prostate cancer: a systematic review and meta-analysis. *European urology*. 2016 Jan;69(1):107-15. PMID: 25481455
106. Cuzick, J, Berney, DM, Fisher, G, et al. Prognostic value of a cell cycle progression signature for prostate cancer death in a conservatively managed needle biopsy cohort. *British journal of cancer*. 2012;106:1095-9. PMID: 22361632
107. Cuzick, J, Swanson, GP, Fisher, G, et al. Prognostic value of an RNA expression signature derived from cell cycle proliferation genes in patients with prostate cancer: a retrospective study. *Lancet Oncol*. 2011 Mar;12(3):245-55. PMID: 21310658

108. Lin, DW, Crawford, ED, Keane, T, et al. Identification of men with low-risk biopsy-confirmed prostate cancer as candidates for active surveillance. *Urologic oncology*. 2018 Jun;36(6):310 e7- e13. PMID: 29655620
109. Bishoff, JT, Freedland, SJ, Gerber, L, et al. Prognostic utility of the cell cycle progression score generated from biopsy in men treated with prostatectomy. *J Urol*. 2014 Aug;192(2):409-14. PMID: 24508632
110. Freedland, SJ, Gerber, L, Reid, J, et al. Prognostic utility of cell cycle progression score in men with prostate cancer after primary external beam radiation therapy. *International journal of radiation oncology, biology, physics*. 2013 Aug 1;86(5):848-53. PMID: 23755923
111. Shore, ND, Kella, N, Moran, B, et al. Impact of the Cell Cycle Progression Test on Physician and Patient Treatment Selection for Localized Prostate Cancer. *J Urol*. 2015 Sep 25. PMID: 26403586
112. Hu, JC, Tosoian, JJ, Qi, J, et al. Clinical Utility of Gene Expression Classifiers in Men With Newly Diagnosed Prostate Cancer. *JCO precision oncology*. 2018;2. PMID: 32832833
113. Crawford, ED, Scholz, MC, Kar, AJ, et al. Cell cycle progression score and treatment decisions in prostate cancer: results from an ongoing registry. *Current medical research and opinion*. 2014 Jun;30(6):1025-31. PMID: 24576172
114. Shore, N, Concepcion, R, Saltzstein, D, et al. Clinical utility of a biopsy-based cell cycle gene expression assay in localized prostate cancer. *Current medical research and opinion*. 2014 Apr;30(4):547-53. PMID: 24320750
115. Berlin, A, Murgic, J, Hosni, A, et al. Genomic Classifier for Guiding Treatment of Intermediate-Risk Prostate Cancers to Dose-Escalated Image Guided Radiation Therapy Without Hormone Therapy. *International journal of radiation oncology, biology, physics*. 2019 Jan 1;103(1):84-91. PMID: 30170099
116. Nguyen, PL, Haddad, Z, Ross, AE, et al. Ability of a Genomic Classifier to Predict Metastasis and Prostate Cancer-specific Mortality after Radiation or Surgery based on Needle Biopsy Specimens. *European urology*. 2017 Nov;72(5):845-52. PMID: 28528811
117. Den, RB, Yousefi, K, Trabulsi, EJ, et al. Genomic classifier identifies men with adverse pathology after radical prostatectomy who benefit from adjuvant radiation therapy. *J Clin Oncol*. 2015 Mar 10;33(8):944-51. PMID: 25667284
118. Den, RB, Feng, FY, Showalter, TN, et al. Genomic prostate cancer classifier predicts biochemical failure and metastases in patients after postoperative radiation therapy. *International journal of radiation oncology, biology, physics*. 2014 Aug 1;89(5):1038-46. PMID: 25035207
119. Cooperberg, MR, Davicioni, E, Crisan, A, Jenkins, RB, Ghadessi, M, Karnes, RJ. Combined value of validated clinical and genomic risk stratification tools for predicting prostate cancer mortality in a high-risk prostatectomy cohort. *European urology*. 2015 Feb;67(2):326-33. PMID: 24998118
120. Ross, AE, Feng, FY, Ghadessi, M, et al. A genomic classifier predicting metastatic disease progression in men with biochemical recurrence after prostatectomy. *Prostate cancer and prostatic diseases*. 2014 Mar;17(1):64-9. PMID: 24145624
121. Karnes, RJ, Bergstralh, EJ, Davicioni, E, et al. Validation of a genomic classifier that predicts metastasis following radical prostatectomy in an at risk patient population. *J Urol*. 2013 Dec;190(6):2047-53. PMID: 23770138

122. Erho, N, Crisan, A, Vergara, IA, et al. Discovery and validation of a prostate cancer genomic classifier that predicts early metastasis following radical prostatectomy. *PLoS one*. 2013;8(6):e66855. PMID: 23826159
123. Prensner, JR, Zhao, S, Erho, N, et al. RNA biomarkers associated with metastatic progression in prostate cancer: a multi-institutional high-throughput analysis of SChLAP1. *Lancet Oncol*. 2014 Dec;15(13):1469-80. PMID: 25456366
124. Klein, EA, Haddad, Z, Yousefi, K, et al. Decipher Genomic Classifier Measured on Prostate Biopsy Predicts Metastasis Risk. *Urology*. 2016 Jan 22. PMID: 26809071
125. Karnes, RJ, Choerung, V, Ross, AE, et al. Validation of a Genomic Risk Classifier to Predict Prostate Cancer-specific Mortality in Men with Adverse Pathologic Features. *European urology*. 2018 Feb;73(2):168-75. PMID: 28400167
126. Spratt, DE, Dai, DLY, Den, RB, et al. Performance of a Prostate Cancer Genomic Classifier in Predicting Metastasis in Men with Prostate-specific Antigen Persistence Postprostatectomy. *European urology*. 2018 Jul;74(1):107-14. PMID: 29233664
127. Freedland, SJ, Choerung, V, Howard, L, et al. Utilization of a Genomic Classifier for Prediction of Metastasis Following Salvage Radiation Therapy after Radical Prostatectomy. *European urology*. 2016 Oct;70(4):588-96. PMID: 26806658
128. Ross, AE, Johnson, MH, Yousefi, K, et al. Tissue-based Genomics Augments Post-prostatectomy Risk Stratification in a Natural History Cohort of Intermediate- and High-Risk Men. *European urology*. 2015 Jun 6. PMID: 26058959
129. Xu, MJ, Kornberg, Z, Gadzinski, AJ, et al. Genomic Risk Predicts Molecular Imaging-detected Metastatic Nodal Disease in Prostate Cancer. *European urology oncology*. 2019 Nov;2(6):685-90. PMID: 31411984
130. Jambor, I, Falagario, U, Ratnani, P, et al. Prediction of biochemical recurrence in prostate cancer patients who underwent prostatectomy using routine clinical prostate multiparametric MRI and decipher genomic score. *Journal of magnetic resonance imaging : JMRI*. 2019 Sep 30. PMID: 31566845
131. Ross, AE, Den, RB, Yousefi, K, et al. Efficacy of post-operative radiation in a prostatectomy cohort adjusted for clinical and genomic risk. *Prostate cancer and prostatic diseases*. 2016 Sep;19(3):277-82. PMID: 27136742
132. Badani, KK, Thompson, DJ, Brown, G, et al. Effect of a genomic classifier test on clinical practice decisions for patients with high-risk prostate cancer after surgery. *BJU international*. 2015 Mar;115(3):419-29. PMID: 24784420
133. Michalopoulos, SN, Kella, N, Payne, R, et al. Influence of a genomic classifier on post-operative treatment decisions in high-risk prostate cancer patients: results from the PRO-ACT study. *Current medical research and opinion*. 2014 Aug;30(8):1547-56. PMID: 24803160
134. Badani, K, Thompson, DJ, Buerki, C, et al. Impact of a genomic classifier of metastatic risk on postoperative treatment recommendations for prostate cancer patients: a report from the DECIDE study group. *Oncotarget*. 2013 Apr;4(4):600-9. PMID: 23592338
135. Nguyen, PL, Shin, H, Yousefi, K, et al. Impact of a Genomic Classifier of Metastatic Risk on Postprostatectomy Treatment Recommendations by Radiation Oncologists and Urologists. *Urology*. 2015 Jul;86(1):35-40. PMID: 26142578
136. Lobo, JM, Dicker, AP, Buerki, C, et al. Evaluating the clinical impact of a genomic classifier in prostate cancer using individualized decision analysis. *PLoS one*. 2015;10(3):e0116866. PMID: 25837660
137. Gore, JL, du Plessis, M, Zhang, J, et al. Clinical Utility of a Genomic Classifier in Men Undergoing Radical Prostatectomy: The PRO-IMPACT Trial. *Practical radiation oncology*. 2020 Mar-Apr;10(2):e82-e90. PMID: 31761540

138. Seiler, R, Ashab, HAD, Erho, N, et al. Impact of Molecular Subtypes in Muscle-invasive Bladder Cancer on Predicting Response and Survival after Neoadjuvant Chemotherapy. *European urology*. 2017 Oct;72(4):544-54. PMID: 28390739
139. Grivas, P, Bismar, TA, Alva, AS, et al. Validation of a neuroendocrine-like classifier confirms poor outcomes in patients with bladder cancer treated with cisplatin-based neoadjuvant chemotherapy. *Urologic oncology*. 2020 Apr;38(4):262-8. PMID: 31812633
140. Necchi, A, Raggi, D, Gallina, A, et al. Impact of Molecular Subtyping and Immune Infiltration on Pathological Response and Outcome Following Neoadjuvant Pembrolizumab in Muscle-invasive Bladder Cancer. *European urology*. 2020 Jun;77(6):701-10. PMID: 32165065
141. Eggener, SE, Rumble, RB, Armstrong, AJ, et al. Molecular Biomarkers in Localized Prostate Cancer: ASCO Guideline. *J Clin Oncol*. 2020 May 1;38(13):1474-94. PMID: 31829902
142. Sanda, MG, Cadeddu, JA, Kirkby, E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part II: Recommended Approaches and Details of Specific Care Options. *J Urol*. 2018 Apr;199(4):990-7. PMID: 29331546
143. Sanda, MG, Cadeddu, JA, Kirkby, E, et al. Clinically Localized Prostate Cancer: AUA/ASTRO/SUO Guideline. Part I: Risk Stratification, Shared Decision Making, and Care Options. *J Urol*. 2018 Mar;199(3):683-90. PMID: 29203269
144. American Urological Association. Early Detection of Prostate Cancer, 2018. [cited 09/14/2020]; Available from: <http://www.auanet.org/education/guidelines/prostate-cancer-detection.cfm>
145. Rosenkrantz, AB, Verma, S, Choyke, P, et al. Prostate Magnetic Resonance Imaging and Magnetic Resonance Imaging Targeted Biopsy in Patients with a Prior Negative Biopsy: A Consensus Statement by AUA and SAR. *J Urol*. 2016 Dec;196(6):1613-8. PMID: 27320841
146. Recommendations from the EGAPP Working Group: does PCA3 testing for the diagnosis and management of prostate cancer improve patient health outcomes? *Genet Med*. 2014;16:338-46. PMID: 24071797
147. National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology™. Prostate Cancer Early Detection. v.2.2020. [cited 09/14/2020]; Available from: https://www.nccn.org/professionals/physician_gls/pdf/prostate_detection.pdf
148. National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology™. Prostate Cancer. v.2.2020. [cited 09/14/2020]; Available from: https://www.nccn.org/professionals/physician_gls/pdf/prostate.pdf
149. National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology™. Bladder Cancer v.6.2020 [cited 9/14/2020]; Available from: https://www.nccn.org/professionals/physician_gls/pdf/bladder_blocks.pdf
150. BlueCross BlueShield Association Medical Policy Reference Manual "Genetic and Protein Biomarkers for the Diagnosis and Cancer Risk Assessment of Prostate Cancer." Policy No. 2.04.33

CODES

Codes	Number	Description
CPT	0011M	Oncology, prostate cancer, mRNA expression assay of 12 genes (10 content and 2 housekeeping), RT-PCR test utilizing blood plasma and urine, algorithms

Codes	Number	Description
		to predict high-grade prostate cancer risk
	0016M	Oncology (bladder), mRNA, microarray gene expression profiling of 209 genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as molecular subtype (luminal, luminal infiltrated, basal, basal claudin-low, neuroendocrine-like)
	0005U	Oncology (prostate) gene expression profile by real-time RT-PCR of 3 genes (ERG, PCA3, and SPDEF), urine, algorithm reported as risk score
	0047U	Oncology (prostate), mRNA, gene expression profiling by real-time RT-PCR of 17 genes (12 content and 5 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a risk score
	0053U	Oncology (prostate cancer), FISH analysis of 4 genes (ASAP1, HDAC9, CHD1 and PTEN), needle biopsy specimen, algorithm reported as probability of higher tumor grade
	81313	PCA3/KLK3 (prostate cancer antigen 3 [non-protein coding]/kallikrein-related peptidase 3 [prostate specific antigen]) ratio (eg, prostate cancer)
	81479	Unlisted molecular pathology procedure
	81541	Oncology (prostate), mRNA gene expression profiling by real-time RT-PCR of 46 genes (31 content and 15 housekeeping), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a disease-specific mortality risk score
	81542	Oncology (prostate), mRNA, microarray gene expression profiling of 22 content genes, utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as metastasis risk score
	81551	Oncology (prostate), promoter methylation profiling by real-time PCR of 3 genes (GSTP1, APC, RASSF1), utilizing formalin-fixed paraffin-embedded tissue, algorithm reported as a likelihood of prostate cancer detection on repeat biopsy
	81599	Unlisted multianalyte assay with algorithmic analysis
HCPCS	None	

Date of Origin: October 2012