

## ***Genotyping for 9p21 Single Nucleotide Polymorphisms to Predict Risk of Cardiovascular Disease or Aneurysm***

**Effective:** July 1, 2018

**Next Review:** May 2019

**Last Review:** May 2018

### **IMPORTANT REMINDER**

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

### **DESCRIPTION**

Single nucleotide polymorphisms (SNPs) found in the 9p21 locus have been associated with myocardial infarction (MI) and other manifestations of cardiovascular disease (CVD). Genotyping for 9p21 SNPs may be offered as an approach to identify patients who may be at increased risk of some of these outcomes.

### **MEDICAL POLICY CRITERIA**

The use of genotyping for 9p21 single nucleotide polymorphisms is considered **investigational** for all indications, including but not limited to identification of:

- A. Patients who may be at increased risk of cardiovascular disease or its manifestations (e.g., myocardial infarction, ischemic stroke, peripheral arterial disease, coronary artery calcification); or
- B. Patients who may be at increased risk for aneurysmal disease (e.g., abdominal aortic aneurysms, intracranial aneurysms, polypoidal choroidal vasculopathy).

*NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.*

## CROSS REFERENCES

1. [Genetic and Molecular Diagnostic Testing](#), Genetic Testing, Policy No. 20

## BACKGROUND

SNPs occur normally throughout a person's DNA. They occur once in every 300 nucleotides on average, which means there are roughly 10 million SNPs in the human genome. Most commonly, these variations are found in the DNA between genes. They can act as biological markers, helping scientists locate genes that are associated with disease. When SNPs occur within a gene or in a regulatory region near a gene, they may play a more direct role in disease by affecting the gene's function.

SNPs are not absolute indicators of disease development. Most SNPs have no effect on health or development. SNPs do not cause disease, but they can help determine the likelihood that someone will develop a particular illness. Some of these genetic differences, however, have proven to be very important in the study of human health. Researchers have found SNPs that may help predict an individual's response to certain drugs, susceptibility to environmental factors such as toxins, and risk of developing specific diseases. SNPs can also be used to track the inheritance of disease genes within families. Future studies will work to identify SNPs associated with complex diseases such as heart disease, diabetes, and cancer.

### 9P21 AND CARDIOVASCULAR DISEASE

In 2007, genome-wide association studies using single nucleotide polymorphism (SNP) arrays resulted in the near simultaneous reporting of the first common genetic variant that affects the risk of coronary heart disease (CHD) in Caucasians at chromosome 9p21.3 (also known as 9p21). CHD is defined as inadequate circulation to cardiac muscle and surrounding tissue resulting in myocardial infarction (MI), unstable angina pectoris, coronary revascularization, or death.<sup>[1-4]</sup> Estimates of CHD risk were confirmed in case-control replication studies in a variety of study populations, showing that the identified SNPs were associated with CHD and even more specifically with MI.<sup>[5]</sup> In all studies, the association of any identified SNP with CHD risk was shown to be independent of traditional risk factors.<sup>[5]</sup>

Several studies have extended the 9p21 association to other vascular diseases including ischemic stroke; thus, 9p21 may be reported as associated with cardiovascular disease (CVD) outcomes, defined as including CHD outcomes plus ischemic stroke. Associations have also been reported with abdominal aortic aneurysm and with intracranial arterial aneurysm.<sup>[6]</sup>

Several genes are found at the 9p21 locus, including ANRIL, which encodes a large noncoding RNA that may have regulatory functions, and CDKN2A and CDKN2B, which encode cyclin-dependent kinase inhibitors.<sup>[6]</sup> The mechanisms by which the SNPs lead to increased CHD risk have been largely unknown. Harismendy (2011) identified several potential enhancer regulatory DNA sequences in the 9p21 region.<sup>[7]</sup> They reported that the SNP rs10747278, consistently associated with increased risk of CHD, occurs in one of these enhancer sequences and that the risk allele disrupts a transcription factor binding site involved in the inflammatory response (STAT1). The interaction of STAT1 with part of the inflammatory signaling pathway, interferon-gamma, is impaired in 9p21 risk carriers. Congrains (2012) genotyped 18 SNPs across the CVD-associated region and determined the impact of 9p21 variants on gene expression.<sup>[8]</sup> The authors reported that, "several SNPs in 9p21 locus affect the expression of ANRIL, which is further in control of the regulation of CDKN2A/B and cell

growth. Cell proliferation mediates the progression of atherosclerosis and is also directly or indirectly involved in the pathogenesis of diseases associated with this locus.”

## COMMERCIAL TESTING

Quest Diagnostics offers the CardioIQ 9p21 Genotype test, which detects the rs10757278 A>G and rs1333049 G>C SNPs within the 9p21 locus of chromosome. The information on the website indicates that the SNPs are associated with the risk of coronary heart disease.<sup>[9]</sup> It is suggested that the test is suitable for individuals with an intermediate risk of CHD, but is not appropriate for use in African Americans.

Cardiac risk genotyping panels offered by other laboratories may include and individually report 9p21 SNP results. For example, the deCODE MI™ (deCODE Genetics) test genotypes 9p21.3 rs10757278 in addition to seven other SNPs from other chromosomal loci to estimate the risk of coronary heart disease and MI. Quest Diagnostics offers the Cardio IQ® 9p21 Genotype test for rs10757278 (A>G) and rs1333049 (G>C)

## REGULATORY STATUS

There is no manufactured test kit for 9p21 genotyping that has been reviewed by the U.S. Food and Drug Administration (FDA). Clinical laboratories may develop and validate tests in-house for 9p21 (“home-brew”) and market them as a laboratory service; such tests must meet the general regulatory standards of the Clinical Laboratory Improvement Act (CLIA). The laboratory offering the service must be licensed by CLIA for high-complexity testing.

## EVIDENCE SUMMARY

Human Genome Variation Society (HGVS) nomenclature<sup>[10]</sup> is used to describe variants found in DNA and serves as an international standard. It is being implemented for genetic testing medical evidence review updates starting in 2017. According to this nomenclature, the term “variant” is used to describe a change in a DNA or protein sequence, replacing previously-used terms, such as “mutation.” Pathogenic variants are variants associated with disease, while benign variants are not. The majority of genetic changes have unknown effects on human health, and these are referred to as variants of uncertain significance.

Validation of the clinical use of any genetic test focuses on three main principles:

1. Analytic validity, which refers to the technical accuracy of the test in detecting a variant that is present or in excluding a variant that is absent;
2. Clinical validity, which refers to the diagnostic performance of the test (i.e., sensitivity, specificity, positive and negative predictive values) in detecting clinical disease; and
3. Clinical utility, which refers to how the results of the diagnostic test will be used to change disease management and whether these changes in management lead to clinically important improvements in health outcomes.

This evidence review evaluates well-designed studies regarding the ability of test results to:

- Guide decisions in the clinical setting related to either treatment, management, or prevention; and
- Improve health outcomes as a result of those decisions.

## 9P21 POLYMORPHISMS AND CORONARY HEART DISEASE (CHD)

A number of studies address analytical validity, clinical validity, and clinical utility as described below. Clinical utility is satisfied when the evidence shows that using a test to change medical management for at least some patients significantly improves outcomes. Most of the evidence regarding the clinical utility of 9p21 testing is related to its role in risk-stratifying patients for coronary heart disease; a smaller body of evidence exists for its utility in other conditions. Direct evidence for how patient management is changed with 9p21 testing is not addressed in the literature.

## Systematic Reviews

Chen (2015) performed a meta-analysis to evaluate the association between the rs10757278 polymorphism (or its proxy rs1333049) on chromosome 9p21 with myocardial infarction (MI) susceptibility in individuals of Caucasian and Chinese ancestry, pooling data from 17 large case-control studies (n=14,924 cases and 28,039 controls).<sup>[11]</sup> The investigators reported a significant association between rs10757278 and MI with a modest odds ratio (OR) of 1.29 (95% confidence interval [CI] 1.22 to 1.36,  $p=6.09 \times 10^{-22}$ ) in the pooled population. Further subgroup analysis found significant association between rs10757278 polymorphism and MI in Asian (OR 1.21, 95% CI 1.16 to 1.27,  $p=1.82 \times 10^{-17}$ ) and Caucasian populations (OR 1.34, 95% CI 1.28 to 1.40,  $p=8.51 \times 10^{-39}$ ). Since this is the first report of a strong association between rs10757278 and MI, directly conflicting with previously published reports, additional studies of this size are needed.

In a systematic review and meta-analysis, Patel (2014) compared the association between variants at the chromosome 9p21 locus and risk of first versus subsequent CHD events.<sup>[12]</sup> The authors identified 31 cohorts reporting on 193,372 individuals. Among the 16 cohorts of individuals without prior CHD (n=168,209), there were 15,664 first CHD events. Variants in 9p21 were associated with first event with a pooled hazard ratio (HR) of 1.19 (95% CI 1.17 to 1.22) per risk allele. In individuals with established CHD (n=25,163) there were 4,436 subsequent events providing >99% and 91% power to detect a per-allele HR of 1.19 or 1.10, respectively. The pooled HR for subsequent events was 1.01 (95% CI 0.97 to 1.06) per risk allele. There was strong evidence of heterogeneity between the effect estimates for first and subsequent events (p-value for heterogeneity= $5.6 \times 10^{-11}$ ). The authors concluded that 9p21 shows differential association with risk of first versus subsequent CHD events.

In a meta-analysis of 21 studies that included patients with information on CAD, MI status and 9p21 genotype (n=33,673), Chan (2013) also found associations with CAD and the 9p21 locus.<sup>[13]</sup> The authors suggest that the 9p21 has a stronger association with CAD compared to MI.

Another meta-analysis, by Dong (2013) of 21 case-control studies evaluated the association between 9p21 SNPs and CHD in an East Asian population, including 25,945 cases and 31,777 controls, and found a significant association between the rs1333049 and CHD (OR 1.30, 95% CI 1.25 to 1.35,  $p<0.001$ ).<sup>[14]</sup>

Zhou (2012) conducted a meta-analysis of seven case-control studies (n=7,123 total). Authors suggest the genetic variation on the 9p21 chromosome may contribute to early-onset CAD, however the effect size was small.<sup>[15]</sup> Another 2014 meta-analysis by different authors evaluating case-control studies had similar findings.<sup>[16]</sup>

Schunkert (2011) conducted a meta-analysis of 14 genome-wide association studies of coronary artery disease (CAD).<sup>[17]</sup> Authors concluded that their large-scale meta-analysis

identified the association of CAD with 13 novel chromosomal loci. Authors suggested these newly identified loci affect CAD risk carriers and may improve treatment of this common disease.

The Coronary Artery Disease Genetics Consortium meta-analyzed four large genome-wide association studies of CAD and identified five loci newly associated with CAD.<sup>[18]</sup> Authors suggested their findings may implicate new pathways for CAD susceptibility. These results compare well with those of Palomaki (2010), who conducted the first formal systematic review of the 9p21 literature to estimate the strength of the association between established 9p21 SNP variants and coronary heart disease and to examine clinical utility.<sup>[19]</sup> Authors reviewed the published literature for effect size, heterogeneity, publication bias, strength of evidence, and evidence of clinical utility of the test. Authors analyzed 47 data sets from 22 articles were analyzed including 35,872 cases and 95,837 controls. The authors concluded that the association between 9p21 SNPs and heart disease, which varied by age at disease onset, was statistically significant; however, the magnitude of the association was small.

This group also addressed clinical utility with a reclassification analysis, evaluating whether genotyping helped reclassify individuals more accurately than traditional risk factors according to their known outcomes, which was measured by calculating the net reclassification index (NRI) with data from three studies (four data sets).<sup>[19]</sup> None of the NRIs were statistically significant. In addition, the study showing the largest NRI achieved most of the risk reclassification because of reduced risk in individuals without events, which would have less chance of improving outcomes. Moreover, in two individual studies the NRI actually worsened when 9p21 risk alleles were added to algorithms that also included family history as a CAD risk factor.<sup>[20,21]</sup> Therefore based on this meta-analysis, evidence for clinical utility of 9p21 testing is insufficient.

## **Randomized Controlled Trials**

While there have been no randomized trials focused on genetic testing for 9p21 SNPs, Anand (2016) published the results of a single-blind, randomized trial of a digital health intervention designed to reduce the burden of MI in South Asian people living in Ontario and British Columbia, a group with a relatively high burden of premature MI.<sup>[22]</sup> The main outcome of the study was a change in MI risk score, which included clinical risk measures as well as genetic risk as determined by 9p21 risk alleles. The intervention used emails or text messages to promote diet and physical activity changes, but found no significant differences between groups after one year. Additionally, the follow-up results did not appear to be influenced by participants' knowledge of their genetic risk.

## **Nonrandomized Studies**

Several studies analyzing individual patient cohorts or case-control populations for association of 9p21 and CHD/CAD have been published since the systematic reviews described above.<sup>[23-30]</sup> Most have found statistically significant, but small associations, with substantial heterogeneity between subgroups.

Evidence for the clinical utility of 9p21 variant testing is not addressed in the literature. Risk assessment may influence patient and provider decisions about preventive interventions and behavioral change. However, as Palomaki (2010)<sup>[19]</sup> noted, only 37% of U.S. physicians reported regular use of a heart disease risk score,<sup>[31]</sup> and the evidence that such risk scores translate into net clinical benefits is minimal.<sup>[32]</sup> Thus, the clinical utility of 9p21 genotyping

cannot be assumed even if risk assessment is improved. As noted, the evidence related to the clinical utility of 9p21 testing is related to its role in risk-stratifying patients for coronary heart disease; a smaller body of evidence exists for its utility in other conditions.<sup>[20,21,33-37]</sup>

## Section Summary

The clinical utility of 9p21 testing, or how patient management may change as a result of this testing, has not been established. The contribution of 9p21 to overall cardiovascular risk, above that of traditional risk factors, is small and not likely to be clinically important. Studies of risk reclassification do not report that 9p21 testing results in substantial numbers of patients being reclassified to clinically relevant categories. Additionally, a study testing a digital health intervention found no improvements in MI risk associated with knowledge of 9p21 genetic risk.

## 9P21 ASSOCIATION WITH ISCHEMIC STROKE

Several analytical and clinical validity studies have reported, with mixed results, on the association of 9p21 with ischemic stroke. There are no clinical utility studies which address how 9p21 test results are used to improved health outcomes in patients at risk for ischemic stroke.

### Systematic Reviews

Anderson (2010) conducted a meta-analysis of eight studies, focusing on two 9p21 SNPs, s1537378 and rs10757278.<sup>[38]</sup> Authors concluded that the variants on 9p21 were associated with ischemic stroke.

In a meta-analysis by Traylor (2012) of 15 studies that included 12,389 individuals with ischemic stroke and 62,004 controls, the 9p21 locus was only associated with large-vessel stroke.<sup>[39]</sup>

Ni (2014) reported results of a meta-analysis of genetic association studies between 9p21 polymorphisms and ischemic stroke, which included 21 studies with 34,128 patients and 153,428 controls.<sup>[40]</sup> The rs10757278 polymorphism was significantly associated with increased overall ischemic stroke risk (per allele OR for ischemic stroke: 1.11, 95% CI 1.07 to 1.15,  $p < 10^{-5}$ ) and increased large-vessel stroke risk (per-allele OR for large vessel stroke, 1.15, 95% CI 1.10 to 1.19), but not with small vessel, cardioembolic, or other types of stroke.

### Randomized Controlled Trials

There have been no randomized trials of genetic testing 9p21 SNPs for stroke risk.

### Nonrandomized Studies

Since publication of the recent meta-analyses, several nonrandomized studies have evaluated the association between 9p21 polymorphisms and ischemic stroke.

Lu (2015) published a case-control analysis of six SNPs on 9p21 and their association with ischemic stroke and carotid plaque in 528 patients with ischemic stroke (375 with carotid plaque and 153 without carotid plaque) and 258 controls.<sup>[41]</sup> The SNPs rs2383206 (OR 1.472,  $p = 0.021$ ) and rs4977574 (OR 1.519,  $p = 0.013$ ) were significantly associated with ischemic stroke without carotid plaque compared to controls. SNPs rs2383206 and rs4977574 were also modestly associated with a risk of carotid plaque among patients with ischemic stroke above age 65 (OR 2.329,  $p = 0.018$  and OR 1.997,  $p = 0.049$ , respectively). However, all six SNPs

tested were in linkage disequilibrium with each other, and were no longer significantly associated after correction for multiple testing.

Bi (2015) published a case-control analysis of four SNPs on 9p21 and their association with ischemic stroke in 116 cases and 118 controls.<sup>[42]</sup> The authors reported that the SNPs rs10757278, rs1537378 and rs1333047 conveyed anywhere between a 1.64- to 2.0-fold increased risk of ischemic stroke. However, these results were marginally significant and once corrected for multiple testing, would not have fallen within the limit considered significant. In addition, there was also a significant association between rs10757278 and lipid levels.

Yue (2015) reported that the significant statistical associations between the SNPs rs2383207, rs3731245, and rs1537378 were significantly associated with cerebral infarction in a Chinese Han population (OR 1.18, 95% CI 1.01 to 1.37; OR 1.29, 95% CI 1.06 to 1.56; OR 1.30, 95% CI 1.05 to 1.60, respectively).<sup>[43]</sup>

Dichgans (2014) analyzed data from the CARDIOGRAM/C4D consortium study described above<sup>[17,44]</sup> in conjunction with data from the METASTROKE consortium<sup>[39]</sup> to evaluate whether CAD and ischemic stroke share genetic risk in respect to common genetic variants.<sup>[45]</sup> The authors found that the 9p21 locus was significantly associated with both CAD and the phenotype of large artery stroke (PLAS =  $3.85 \times 10^{-6}$ ; Spearman's rho coefficient for large artery stroke/CAS = 0.85,  $p=2.9E^{-35}$ ).

## Section Summary

Studies on the clinical and analytical validity of 9p21 association with ischemic stroke reported mixed results. Further, the clinical utility of 9p21 variant testing for ischemic stroke has not been established.

## 9P21 ASSOCIATION WITH ANEURYSM

The 9p21 locus has been associated with risk of both intracranial and abdominal aortic aneurysms. There are no clinical utility studies on the association of 9p21 with aneurysm.

## Systematic Review

Alg (2013) reported results from a systematic review and meta-analysis of all genetic association studies of sporadic intracranial aneurysm to identify genetic risk factors for intracranial aneurysm<sup>[46]</sup>. The authors included 66 cohort or case-control studies of intracranial aneurysms that examined a total of 41 SNPs, not limited to the 9p21 locus, in 29 genes. Among polymorphisms with the strongest associations with intracranial aneurysm were the 9p21 SNPs rs10757278 (OR 1.29, 95% CI 1.21 to 1.38) and rs1333040 (OR 1.24, 95% CI 1.20 to 1.29).

## Randomized Controlled Trials

There have been no randomized trials of 9p21 SNP testing for aneurysm risk.

## Nonrandomized Studies

Research on the association of 9p21 with abdominal aortic aneurysm (AAA) includes several studies that reported 9p21 allele-specific estimates of risk in the range of 1.2-1.8.<sup>[23,47-50]</sup> Biros (2010) combined the results of their study with the results of previous studies and reported a combined estimate of about 1.3 for both 9p21 SNPs (rs10757278 and rs1333049).<sup>[50]</sup> This

estimate is lower than other well-characterized risk factor estimates for AAA such as age, family history, and smoking.<sup>[51]</sup> Wei (2014) reported slightly higher risk of AAA associated with homozygosity for the rs10757278 and rs1333040 risk alleles in a Chinese Han population, after controlling for other AAA risk factors (OR 2.31, 95% CI 1.22 to 4.36; OR 2.14, 95% CI 1.13 to 4.05, respectively).<sup>[52]</sup>

### **Section Summary**

The evidence that addresses the clinical validity of 9p21 for both intracranial and abdominal aortic aneurysms is limited. Further, the clinical utility of 9p21 variant testing for aneurysm has not been established.

### **9P21 ASSOCIATION WITH OTHER CONDITIONS**

Analytical and clinical validity studies have been reported, with mixed results, on the association of 9p21 with other conditions. There are no clinical utility studies on the association of 9p21 with the other conditions described below.

A few studies have explored the association of 9p21 variants with a variety of other conditions such as peripheral arterial disease,<sup>[53]</sup> coronary artery calcification,<sup>[54-56]</sup> aortic calcification,<sup>[55]</sup> polypoidal choroidal vasculopathy (characterized by aneurismal dilations at the border of the choroidal vascular network)<sup>[57]</sup>, cardiovascular mortality in the absence of coronary lesions<sup>[58]</sup>, and arterial stiffness in hypertensive individuals.<sup>[59]</sup> The strength of the associations was modest and none suggested clinical use.

In contrast, Folsom (2013) found no association between SNPs at the 9p21 locus with arterial elasticity and retinal microvascular diameter.<sup>[60]</sup>

Downing (2014) evaluated the impact of adding 9p21 polymorphism (rs10757269) in a risk-factor-based model predicting peripheral artery disease.<sup>[61]</sup> Among 393 subjects in the prospective Genetic Determinants of Peripheral Artery Disease study who met study inclusion criteria, the rs10757269 allele was associated with the presence of peripheral artery disease (defined as ankle-brachial index < 0.9) after controlling for traditional cardiovascular risk factors and other biomarkers (OR 1.92, 95% CI 1.29 to 2.85). The addition of 9p21 genotype to a previously-validated peripheral artery disease risk model (including age, sex, race, smoking history, body mass index, hypertension stage, diabetes status, and history of cardiovascular disease, congestion heart failure, and CAD) led to improved risk classification (net reclassification index 33.5%, p=0.001).

### **Section Summary**

Evidence that addresses the analytical and clinical validity, as well as the clinical utility of 9p21 association with other conditions, including but not limited to peripheral artery disease, coronary artery calcification, aortic calcification, polypoidal choroidal vasculopathy, and arterial stiffness is insufficient.

## **PRACTICE GUIDELINE SUMMARY**

### **AMERICAN COLLEGE OF CARDIOLOGY (ACC)/AMERICAN HEART ASSOCIATION (AHA)**

In 2013, the American College of Cardiology Foundation and the American Heart Association



Task Force on Practice Guidelines issued guidelines on the assessment of cardiovascular risk, which did not address assessment of 9p21 polymorphisms.<sup>[62]</sup>

## EGAPP WORKING GROUP (EWG)

The EGAPP Working Group (EWG) published a recommendation on “genomic profiling to assess cardiovascular risk to improve cardiovascular health” which included a recommendation on 9p21 profiling alone based on Palomaki (2010).<sup>[19]</sup> In general, the EWG found “... insufficient evidence to recommend testing for the 9p21 genetic variant or 57 other variants in 28 genes . . . to assess risk for cardiovascular disease (CVD) in the general population, specifically heart disease and stroke.” The EWG found that the magnitude of net health benefit from use of any of these tests alone or in combination is negligible. The EWG discouraged clinical use unless further evidence supports improved clinical outcomes. Based on the available evidence, the overall certainty of net health benefit was deemed “Low.”<sup>[63]</sup>

## SUMMARY

There is not enough research to show that testing for 9p21 SNP alleles can improve health outcomes for patients with any conditions. Also, there are no practice guidelines based on research that recommend this testing for any purpose. Therefore, 9p21 genotyping for all applications is considered investigational.

## REFERENCES

1. Wellcome Trust Case Control Consortium. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007 Jun 7;447(7145):661-78. PMID: 17554300
2. McPherson, R, Pertsemlidis, A, Kavaslar, N, et al. A common allele on chromosome 9 associated with coronary heart disease. *Science*. 2007 Jun 8;316(5830):1488-91. PMID: 17478681
3. Helgadottir, A, Thorleifsson, G, Manolescu, A, et al. A common variant on chromosome 9p21 affects the risk of myocardial infarction. *Science*. 2007 Jun 8;316(5830):1491-3. PMID: 17478679
4. Samani, NJ, Erdmann, J, Hall, AS, et al. Genomewide association analysis of coronary artery disease. *N Engl J Med*. 2007 Aug 2;357(5):443-53. PMID: 17634449
5. Scheffold, T, Kullmann, S, Hüge, A, et al. Six sequence variants on chromosome 9p21.3 are associated with a positive family history of myocardial infarction: a multicenter registry. *BMC cardiovascular disorders*. 2011;11:9. PMID: 21385355
6. Johansen, CT, Lanktree, MB, Hegele, RA. Translating genomic analyses into improved management of coronary artery disease. *Future cardiology*. 2010 Jul;6(4):507-21. PMID: 20608823
7. Harismendy, O, Notani, D, Song, X, et al. 9p21 DNA variants associated with coronary artery disease impair interferon-gamma signalling response. *Nature*. 2011 Feb 10;470(7333):264-8. PMID: 21307941
8. Congrains, A, Kamide, K, Oguro, R, et al. Genetic variants at the 9p21 locus contribute to atherosclerosis through modulation of ANRIL and CDKN2A/B. *Atherosclerosis*. 2012 Feb;220(2):449-55. PMID: 22178423

9. Quest Diagnostics Cardio IQ 9p21 Genotype. [cited; Available from: [http://www.questdiagnostics.com/testcenter/testguide.action?dc=TS\\_9p21Genotype\\_CardioIQ&tabview=true](http://www.questdiagnostics.com/testcenter/testguide.action?dc=TS_9p21Genotype_CardioIQ&tabview=true)
10. den Dunnen, JT, Dalgleish, R, Maglott, DR, et al. HGVS Recommendations for the Description of Sequence Variants: 2016 Update. *Human mutation*. 2016 Jun;37(6):564-9. PMID: 26931183
11. Chen, G, Fu, X, Wang, G, Liu, G, Bai, X. Genetic Variant rs10757278 on Chromosome 9p21 Contributes to Myocardial Infarction Susceptibility. *Int J Mol Sci*. 2015;16:11678-88. PMID: 26006241
12. Patel, RS, Asselbergs, FW, Quyyumi, AA, et al. 5Genetic variants at chromosome 9p21 and risk of first versus subsequent coronary heart disease events: A systematic review and meta-analysis. *Journal of the American College of Cardiology*. 2014 Mar 5. PMID: 24607648
13. Chan, K, Patel, RS, Newcombe, P, et al. Association between the chromosome 9p21 locus and angiographic coronary artery disease burden: a collaborative meta-analysis. *Journal of the American College of Cardiology*. 2013 Mar 5;61(9):957-70. PMID: 23352782
14. Dong, L, Wang, H, Wang, DW, Ding, H. Association of chromosome 9p21 genetic variants with risk of coronary heart disease in the East Asian population: a meta-analysis. *Annals of human genetics*. 2013 May;77(3):183-90. PMID: 23347249
15. Zhou, LT, Qin, L, Zheng, DC, Song, ZK, Ye, L. Meta-analysis of genetic association of chromosome 9p21 with early-onset coronary artery disease. *Gene*. 2012 Dec 1;510(2):185-8. PMID: 22975211
16. Lian, J, Ba, Y, Dai, D, et al. A replication study and a meta-analysis of the association between the CDKN2A rs1333049 polymorphism and coronary heart disease. *Journal of atherosclerosis and thrombosis*. 2014;21(11):1109-20. PMID: 24930384
17. Schunkert, H, König, IR, Kathiresan, S, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nature genetics*. 2011 Apr;43(4):333-8. PMID: 21378990
18. A genome-wide association study in Europeans and South Asians identifies five new loci for coronary artery disease. *Nature genetics*. 2011 Apr;43(4):339-44. PMID: 21378988
19. Palomaki, GE, Melillo, S, Bradley, LA. Association between 9p21 genomic markers and heart disease: a meta-analysis. *JAMA : the journal of the American Medical Association*. 2010 Feb 17;303(7):648-56. PMID: 20159873
20. Paynter, NP, Chasman, DI, Buring, JE, Shiffman, D, Cook, NR, Ridker, PM. Cardiovascular disease risk prediction with and without knowledge of genetic variation at chromosome 9p21.3. *Annals of internal medicine*. 2009 Jan 20;150(2):65-72. PMID: 19153409
21. Talmud, PJ, Cooper, JA, Palmen, J, et al. Chromosome 9p21.3 coronary heart disease locus genotype and prospective risk of CHD in healthy middle-aged men. *Clinical chemistry*. 2008 Mar;54(3):467-74. PMID: 18250146
22. Anand, SS, Samaan, Z, Middleton, C, et al. A Digital Health Intervention to Lower Cardiovascular Risk: A Randomized Clinical Trial. *JAMA cardiology*. 2016 Aug 01;1(5):601-6. PMID: 27438754
23. Bendjilali, N, Nelson, J, Weinsheimer, S, et al. Common variants on 9p21.3 are associated with brain arteriovenous malformations with accompanying arterial aneurysms. *Journal of neurology, neurosurgery, and psychiatry*. 2014 Nov;85(11):1280-3. PMID: 24777168

24. Cakmak, HA, Bayoglu, B, Durmaz, E, et al. Evaluation of association between common genetic variants on chromosome 9p21 and coronary artery disease in Turkish population. *Anatolian journal of cardiology*. 2015 Mar;15(3):196-203. PMID: 25333979
25. Szpakowicz, A, Kiliszek, M, Pepinski, W, et al. Polymorphism of 9p21.3 locus is associated with 5-year survival in high-risk patients with myocardial infarction. *PloS one*. 2014;9(8):e104635. PMID: 25105296
26. Tang, O, Lv, J, Cheng, Y, Qin, F. The Correlation Between 9p21 Chromosome rs4977574 Polymorphism Genotypes and the Development of Coronary Artery Heart Disease. *Cardiovascular toxicology*. 2017 Apr;17(2):185-9. PMID: 27240780
27. Matoo, S, Fallah, MS, Daneshpour, MS, et al. Increased Risk of CHD in the Presence of rs7865618 (A allele): Tehran Lipid and Glucose Study. *Archives of Iranian medicine*. 2017 Mar;20(3):153-7. PMID: 28287809
28. Paquette, M, Chong, M, Saavedra, YGL, Pare, G, Dufour, R, Baass, A. The 9p21.3 locus and cardiovascular risk in familial hypercholesterolemia. *Journal of clinical lipidology*. 2017 Mar - Apr;11(2):406-12. PMID: 28502497
29. Aleyasin, SA, Navidi, T, Davoudi, S. Association between rs10757274 and rs2383206 SNPs as Genetic Risk Factors in Iranian Patients with Coronary Artery Disease. *J Tehran Heart Cent*. 2017;12(3):114-8. PMID:
30. Kashyap, S, Kumar, S, Agarwal, V, Misra, DP, Rai, MK, Kapoor, A. The association of polymorphic variants, rs2267788, rs1333049 and rs2383207 with coronary artery disease, its severity and presentation in North Indian population. *Gene*. 2018;648:89-96. PMID:
31. Sposito, AC, Ramires, JA, Jukema, JW, et al. Physicians' attitudes and adherence to use of risk scores for primary prevention of cardiovascular disease: cross-sectional survey in three world regions. *Current medical research and opinion*. 2009 May;25(5):1171-8. PMID: 19323611
32. Sheridan, SL, Crespo, E. Does the routine use of global coronary heart disease risk scores translate into clinical benefits or harms? A systematic review of the literature. *BMC health services research*. 2008;8:60. PMID: 18366711
33. Davies, RW, Dandona, S, Stewart, AF, et al. Improved prediction of cardiovascular disease based on a panel of single nucleotide polymorphisms identified through genome-wide association studies. *Circulation Cardiovascular genetics*. 2010 Oct;3(5):468-74. PMID: 20729558
34. Dutta, A, Henley, W, Lang, IA, et al. The coronary artery disease-associated 9p21 variant and later life 20-year survival to cohort extinction. *Circulation Cardiovascular genetics*. 2011 Oct;4(5):542-8. PMID: 21852414
35. Shiffman, D, O'Meara, ES, Rowland, CM, et al. The contribution of a 9p21.3 variant, a KIF6 variant, and C-reactive protein to predicting risk of myocardial infarction in a prospective study. *BMC cardiovascular disorders*. 2011;11:10. PMID: 21406102
36. Do, R, Xie, C, Zhang, X, et al. The effect of chromosome 9p21 variants on cardiovascular disease may be modified by dietary intake: evidence from a case/control and a prospective study. *PLoS medicine*. 2011 Oct;8(10):e1001106. PMID: 22022235
37. Brautbar, A, Ballantyne, CM, Lawson, K, et al. Impact of adding a single allele in the 9p21 locus to traditional risk factors on reclassification of coronary heart disease risk and implications for lipid-modifying therapy in the Atherosclerosis Risk in Communities study. *Circulation Cardiovascular genetics*. 2009 Jun;2(3):279-85. PMID: 20031596
38. Anderson, CD, Biffi, A, Rost, NS, Cortellini, L, Furie, KL, Rosand, J. Chromosome 9p21 in ischemic stroke: population structure and meta-analysis. *Stroke*. 2010 Jun;41(6):1123-31. PMID: 20395606

39. Traylor, M, Farrall, M, Holliday, EG, et al. Genetic risk factors for ischaemic stroke and its subtypes (the METASTROKE collaboration): a meta-analysis of genome-wide association studies. *Lancet neurology*. 2012 Nov;11(11):951-62. PMID: 23041239
40. Ni, X, Zhang, J. Association between 9p21 genomic markers and ischemic stroke risk: evidence based on 21 studies. *PloS one*. 2014;9(3):e90255. PMID: 24625579
41. Lu, Z, Zhang, Y, Maimaiti, Y, et al. Variants on Chromosome 9p21 Confer Risks of Noncardioembolic Cerebral Infarction and Carotid Plaque in the Chinese Han Population. *Journal of atherosclerosis and thrombosis*. 2015;22(10):1061-70. PMID: 25958930
42. Bi, J, Yang, L, Liu, D, et al. Sequence variants on chromosome 9p21 are associated with ischemic stroke and the lipids level in Chinese Han population. *Journal of stroke and cerebrovascular diseases : the official journal of National Stroke Association*. 2015 Apr;24(4):894-900. PMID: 25724239
43. Yue, X, Tian, L, Fan, X, Xu, G, Shi, FD, Liu, X. Chromosome 9p21.3 Variants Are Associated with Cerebral Infarction in Chinese Population. *Journal of molecular neuroscience : MN*. 2015 Feb 11. PMID: 25665551
44. Preuss, M, Konig, IR, Thompson, JR, et al. Design of the Coronary ARtery Disease Genome-Wide Replication And Meta-Analysis (CARDIoGRAM) Study: A Genome-wide association meta-analysis involving more than 22 000 cases and 60 000 controls. *Circulation Cardiovascular genetics*. 2010 Oct;3(5):475-83. PMID: 20923989
45. Dichgans, M, Malik, R, Konig, IR, et al. Shared genetic susceptibility to ischemic stroke and coronary artery disease: a genome-wide analysis of common variants. *Stroke*. 2014;45:24-36. PMID: 24262325
46. Alg, VS, Sofat, R, Houlden, H, Werring, DJ. Genetic risk factors for intracranial aneurysms: a meta-analysis in more than 116,000 individuals. *Neurology*. 2013;80:2154-65. PMID: 23733552
47. Thompson, AR, Golledge, J, Cooper, JA, Hafez, H, Norman, PE, Humphries, SE. Sequence variant on 9p21 is associated with the presence of abdominal aortic aneurysm disease but does not have an impact on aneurysmal expansion. *European journal of human genetics : EJHG*. 2009 Mar;17(3):391-4. PMID: 18854858
48. Helgadottir, A, Thorleifsson, G, Magnusson, KP, et al. The same sequence variant on 9p21 associates with myocardial infarction, abdominal aortic aneurysm and intracranial aneurysm. *Nature genetics*. 2008 Feb;40(2):217-24. PMID: 18176561
49. Bown, MJ, Braund, PS, Thompson, J, London, NJ, Samani, NJ, Sayers, RD. Association between the coronary artery disease risk locus on chromosome 9p21.3 and abdominal aortic aneurysm. *Circulation Cardiovascular genetics*. 2008 Oct;1(1):39-42. PMID: 20031540
50. Biros, E, Cooper, M, Palmer, LJ, Walker, PJ, Norman, PE, Golledge, J. Association of an allele on chromosome 9 and abdominal aortic aneurysm. *Atherosclerosis*. 2010 Oct;212(2):539-42. PMID: 20605023
51. Lederle, FA, Johnson, GR, Wilson, SE, et al. The aneurysm detection and management study screening program: validation cohort and final results. Aneurysm Detection and Management Veterans Affairs Cooperative Study Investigators. *Archives of internal medicine*. 2000 May 22;160(10):1425-30. PMID: 10826454
52. Wei, Y, Xiong, J, Zuo, S, et al. Association of polymorphisms on chromosome 9p21.3 region with increased susceptibility of abdominal aortic aneurysm in a Chinese Han population. *Journal of vascular surgery*. 2014 Apr;59(4):879-85. PMID: 24365123
53. Murabito, JM, White, CC, Kavousi, M, et al. Association between chromosome 9p21 variants and the ankle-brachial index identified by a meta-analysis of 21 genome-wide

- association studies. *Circulation Cardiovascular genetics*. 2012 Feb 1;5(1):100-12. PMID: 22199011
54. O'Donnell, CJ, Kavousi, M, Smith, AV, et al. Genome-wide association study for coronary artery calcification with follow-up in myocardial infarction. *Circulation*. 2011 Dec 20;124(25):2855-64. PMID: 22144573
  55. van Setten, J, Isgum, I, Smolonska, J, et al. Genome-wide association study of coronary and aortic calcification implicates risk loci for coronary artery disease and myocardial infarction. *Atherosclerosis*. 2013 Mar 13. PMID: 23561647
  56. Pechlivanis, S, Muhleisen, TW, Mohlenkamp, S, et al. Risk loci for coronary artery calcification replicated at 9p21 and 6q24 in the Heinz Nixdorf Recall Study. *BMC medical genetics*. 2013;14:23. PMID: 23394302
  57. Zhang, X, Wen, F, Zuo, C, Li, M, Chen, H, Wu, K. Association of genetic variation on chromosome 9p21 with polypoidal choroidal vasculopathy and neovascular age-related macular degeneration. *Investigative ophthalmology & visual science*. 2011 Oct;52(11):8063-7. PMID: 21896860
  58. Lee, IT, Liang, KW, Wang, JS, et al. Value of Chromosome 9p21 Polymorphism for Prediction of Cardiovascular Mortality in Han Chinese Without Coronary Lesions: An Observational Study. *Medicine (Baltimore)*. 2015;94:e1538. PMID: 26426617
  59. Cesana, F, Nava, S, Menni, C, et al. Does the 9p region affect arterial stiffness? Results from a cohort of hypertensive individuals. *Blood pressure*. 2013 Oct;22(5):302-6. PMID: 23445356
  60. Folsom, AR, Pankow, JS, Li, X, et al. No association of 9p21 with arterial elasticity and retinal microvascular findings. *Atherosclerosis*. 2013 Oct;230(2):301-3. PMID: 24075760
  61. Downing, KP, Nead, KT, Kojima, Y, et al. The combination of 9p21.3 genotype and biomarker profile improves a peripheral artery disease risk prediction model. *Vasc Med*. 2014;19:3-8. PMID: 24323119
  62. Goff, DC, Jr., Lloyd-Jones, DM, Bennett, G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. 2014 Jul 1;63(25 Pt B):2935-59. PMID: 24239921
  63. Recommendations from the EGAPP Working Group: genomic profiling to assess cardiovascular risk to improve cardiovascular health. *Genetics in medicine : official journal of the American College of Medical Genetics*. 2010 Dec;12(12):839-43. PMID: 21042222

## CODES

Codes	Number	Description
CPT	81479	Unlisted molecular pathology procedure
HCPCS	None	

**Date of Origin:** May 2013