

Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease (GERD)

Effective: March 1, 2019

Next Review: November 2019

Last Review: February 2019

IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

DESCRIPTION

Transesophageal endoscopic therapies are a group of minimally invasive antireflux procedures being investigated as alternatives to medical management or fundoplication surgery in the treatment of GERD.

MEDICAL POLICY CRITERIA

Transesophageal endoscopic therapies are considered **investigational** for the treatment of gastroesophageal reflux disease (GERD). These procedures include but are not limited to the following:

- I. Transesophageal endoscopic gastroplasty procedure (i.e., MUSE)
- II. Transoral incisionless fundoplication (TIF) procedure, (i.e., EsophyX)
- III. Transesophageal radiofrequency energy procedure (i.e., Stretta)
- IV. Endoscopic submucosal implantation of a prosthesis or injection of a bulking agent (i.e., Durasphere, polymethylmethacrylate [PMMA] beads, the Gatekeeper Reflux Repair system)

NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.

CROSS REFERENCES

1. [Bariatric Surgery](#), Surgery, Policy No. 58
2. [Gastric Reflux Surgery](#), Surgery, Policy No. 186
3. [Magnetic Esophageal Ring to Treat Gastroesophageal Reflux Disease \(GERD\)](#), Surgery, Policy No. 190

BACKGROUND

Gastroesophageal reflux disease (GERD) is a common disorder characterized by heartburn and other symptoms related to reflux of stomach acid into the esophagus. Nearly all individuals experience such symptoms at some point in their lives; a smaller number have chronic symptoms and are at risk for complications of GERD. The prevalence of GERD has been estimated to be 10% to 20% in the Western world, with a lower prevalence in Asia.^[1]

The pathophysiology of GERD involves excessive exposure to stomach acid, which occurs for several reasons. There can be an incompetent barrier between the esophagus and stomach, either due to dysfunction of the lower esophageal sphincter (LES) or incompetence of the diaphragm. Another mechanism is abnormally slow clearance of stomach acid by the esophagus. In this situation, delayed clearance leads to an increased reservoir of stomach acid and a greater tendency to reflux.

In addition to troubling symptoms, some patients will have more serious disease, which results in complications such as erosive esophagitis, dysphagia, Barrett esophagus, and esophageal carcinoma. Pulmonary complications may result from aspiration of stomach acid into the lungs and can include asthma, pulmonary fibrosis and bronchitis, or symptoms of chronic hoarseness, cough, and sore throat.

Guidelines on the management of GERD emphasize initial medical management. Weight loss, smoking cessation, head of bed elevation, and elimination of food triggers are all recommended in recent practice guidelines.^[1] Proton pump inhibitors (PPIs) have been shown to be the most effective medical treatment. In a Cochrane systematic review, PPIs demonstrated superiority to H₂-receptor agonists and prokinetics in both network meta-analyses and direct comparisons.^[2]

The most common surgical procedure used for GERD is laparoscopic Nissen fundoplication. Fundoplication involves wrapping a portion of the gastric fundus around the distal esophagus to increase LES pressure. If a hiatal hernia is present, the procedure also restores the position of the LES to the correct location. Laparoscopic fundoplication was introduced in 1991 and has been rapidly adopted because it avoids complications associated with an open procedure.

Although fundoplication results in a high proportion of patients reporting symptom relief, complications can occur, and sometimes require conversion to an open procedure. Patients who have relief of symptoms of GERD after fundoplication may have dysphagia or gas-bloat syndrome (excessive gastrointestinal gas).

Due in part to the high prevalence of gastroesophageal reflux disease, there has been interest in creating a minimally invasive transesophageal therapeutic alternative to open or laparoscopic fundoplication or chronic medical therapy. This type of procedure may be considered natural orifice transluminal surgery. Three types of procedures have been investigated.

1. Transesophageal endoscopic gastroplasty (gastroplication, transoral incisionless fundoplication) can be performed as an outpatient procedure. During this procedure, the fundus of the stomach is folded, and then held in place with staples or fasteners that are deployed by the device. The endoscopic procedure is designed to recreate a valve and barrier to reflux.
2. Radiofrequency (RF) energy has been used to produce submucosal thermal lesions at the gastroesophageal junction. (This technique has also been referred to as the Stretta procedure). Specifically, RF energy is applied through 4 electrodes inserted into the esophageal wall at multiple sites both above and below the squamocolumnar junction. The mechanism of action of the thermal lesions is not precisely known but may be related to ablation of the nerve pathways responsible for sphincter relaxation or may induce a tissue-tightening effect related to heat-induced collagen contraction and fibrosis.
3. Submucosal injection or implantation of a prosthetic or bulking agent to enhance the volume of the lower esophageal sphincter has also been investigated.

One bulking agent, pyrolytic carbon-coated zirconium oxide spheres (Durasphere®), is being evaluated.

The Gatekeeper™ Reflux Repair System (Medtronic) utilizes a soft, pliable, expandable prosthesis made of a polyacrylonitrile-based hydrogel. The prosthesis is implanted into the esophageal submucosa, and with time, the prosthesis absorbs water and expands, creating bulk in the region of implantation.

Endoscopic submucosal implantation of polymethylmethacrylate (PMMA) beads in to the lower esophageal folds has also been investigated.

REGULATORY STATUS

In 2007, EsophyX® (EndoGastric Solutions, Redmond, WA) was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for full-thickness plication. In 2016, EsophyX® Z Device with SerosaFuse Fasteners was cleared for marketing (K160960) by FDA through the 510(k) process for use in transoral tissue approximation, full thickness plication, ligation in the gastrointestinal tract, narrowing the gastroesophageal junction, and reduction of hiatal hernia of 2 cm or less in patients with symptomatic chronic gastroesophageal reflux disease (GERD).^[3] In June 2017, EsophyX2 HD and the third-generation EsophyX Z Devices with SerosaFuse fasteners and accessories were cleared for marketing by FDA through the 510(k) process (K171307) for expanded indications, including patients who require and respond to pharmacologic therapy and in patients with hiatal hernias larger than 2 cm when a laparoscopic hiatal hernia repair reduces the hernia to 2 cm or less.^[4] FDA product code: ODE.

The Medigus SRS Endoscopic Stapling System (MUSE, Medigus) was cleared for marketing by FDA through the 510(k) process in 2012 (K120299) and 2014 (K132151). MUSE is intended for endoscopic placement of surgical staples in the soft tissue of the esophagus and stomach to create anterior partial fundoplication for treatment of symptomatic chronic GERD in patients who require and respond to pharmacologic therapy. FDA product code: ODE.

In 2000, the CSM Stretta® System was cleared for marketing by FDA through the 510(k) process for general use in the electrosurgical coagulation of tissue and is specifically intended

for use in the treatment of GERD. Stretta® is currently manufactured by Mederi Therapeutics (Greenwich, CT). FDA product code: GEI.

Durasphere® is a bulking agent approved for treatment of urinary and fecal incontinence. Use of this product for esophageal reflux would be considered off-label use. The website of Carbon Medical Technologies states that Durasphere GR is an investigational device in the United States “intended to treat problems associated with GERD.”

EVIDENCE SUMMARY

MULTIPLE ENDOSCOPIC PROCEDURES

Systematic Reviews

A 2005 report of the Agency for Healthcare Research and Quality (AHRQ), on “Comparative Effectiveness of Management Strategies for Gastroesophageal Reflux Disease,” indicated additional efficacy and safety data on new endoscopic approaches were needed.^[5] A 2011 update of the AHRQ report excluded Enteryx and the NDO Plicator, since they were no longer available in the U.S., and added the EsophyX procedure (endoscopic fundoplication), which was commercialized after the 2005 review.^[6] The 2011 update reported the following:

The AHRQ report concluded that for the 3 available endoscopic procedures (EndoCinch, Stretta, EsophyX), effectiveness remains substantially uncertain for the long-term management of GERD. While some clinical benefits were observed in patients who had these procedures, the studies were generally small, of variable quality, and of short duration. In addition, all of these procedures have been associated with complications, including dysphagia, infection/fever, and bloating; complications which are also side effects associated with laparoscopic fundoplication^[7] Higher quality studies are needed to determine the role and value of endoscopic procedures in the treatment of patients with GERD. A 2015 review of endoscopic treatment of GERD noted that EndoCinch is no longer manufactured.^[8]

A systematic review was conducted in 2009 to examine 7 endoscopic treatments for GERD that included 33 studies, only 2 of which were RCTs.^[9] The remainder were case series. The authors concluded, “...despite the potential benefits of these procedures, there is insufficient evidence at present to establish their safety and efficacy, particularly in the long term.”

TRANSESOPHAGEAL ENDOSCOPIC GASTROPLASTY AND TRANSORAL INCISIONLESS FUNDOPLICATION (TIF)

Systematic Reviews

McCarty (2018) published a systematic review of RCTs and nonrandomized studies that showed significant improvement in a number of clinical outcomes for patients treated with TIF.^[10] For example, 89% of TIF patients discontinued PPI therapy after the procedure, and the Gastroesophageal Reflux Disease Health-Related Quality of Life (GERD-HRQL) questionnaire, Gastroesophageal Reflux Symptom Score, and Reflux Symptom Index (RSI) measures showed significant improvement. The review had several limitations, including the risk of heterogeneity bias, due to the inclusion of studies of first- and second-generation TIF devices and protocols.

Richter (2018) published a network meta-analysis of RCTs comparing TIF or laparoscopic Nissen fundoplication (LNF) with sham or PPIs.^[11] The meta-analysis was limited by low-quality studies (one did not report randomization method, others lacked data on allocation concealment, blinding of outcome assessors, or other aspects of study protocol). It should be noted that a reason behind for scarcity of direct comparisons between TIF and LNF is the discrepancy in populations requiring the respective treatments: consequently, TIF studies included patients with mild esophagitis and small hiatal hernias (<2 cm), while LNF studies included patients with Los Angeles grade A, B, C, or D esophagitis and all sizes of hiatal hernias.

Randomized Controlled Trials

In 2018, Trad reported five-year outcomes on the manufacturer-sponsored TEMPO randomized controlled trial (RCT).^[12] Three-year results were reported in 2016^[13], other interim results were previously reported as well.^[14,15] Below are highlights from each publication:

- Participants with small or absent hiatal hernias (<2cm) and GERD symptoms while on PPI therapy for at least six months who also had abnormal esophageal acid exposure (EAE) were randomized to either EsophyX® (n=40) treatment or PPI therapy (n=23). After six months of evaluation, 21 remaining PPI therapy participants elected to crossover to EsophyX.
- At three years follow-up, 52 participants were assessed for (1) GERD symptom resolution, (2) healing of esophagitis using endoscopy, (3) EAE, and (4) discontinuation of PPI use. Two participants required revision surgery. As assessed by questionnaire (the Reflux Disease Questionnaire [RDQ], and the Reflux Symptom Index [RSI]), primary outcomes of GERD resolution and elimination of all troublesome atypical symptoms was observed in 37/40 participants, and 42/48 participants, respectively.
- At five years follow-up, data were available for 44 patients, of whom 37 (86%) showed elimination of troublesome regurgitation at 5 years. Twenty (43%) patients were completely off PPIs at the 5-year follow-up, and 31 (70%) patients expressed satisfaction with the procedure, as assessed by the GERD-HRQL scores. While data on pH normalization were available for 24 patients at the 3-year follow-up, at 5 years, 22% (n=5) of these patients could not be assessed for pH normalization.
- Although mean symptom scores were reportedly improved, standard deviations for primary (and secondary) outcomes suggest a wide range of responses and further well-designed studies may be warranted.

In 2015, four RCTs that compared the EsophyX® device to proton pump inhibitor (PPI) treatment or to a sham control were identified, 2 of which were industry sponsored. The studies differed in whether patients' symptoms were or were not controlled on PPI therapy, in the control used (i.e., sham, sham plus PPI, PPI alone), whether patients were blinded to treatment, and in outcome measures. Included in the studies were patients on daily PPI therapy for moderate-to-severe GERD symptoms. Exclusion criteria common to the RCTs are body mass index (BMI) over 35 kg/m², hiatal hernia greater than 2 cm; esophagitis grade C or D; Barrett esophagus greater than 2 cm, and esophageal ulcer. Most studies allowed crossover to the other intervention with continued follow-up after the randomized portion of the study.

The largest RCT with the lowest risk of bias was an industry-sponsored, double-blind, sham-controlled multicenter study (RESPECT) that evaluated TIF in patients whose symptoms were not well controlled on PPIs.^[16] Of 696 patients screened, 129 met inclusion and exclusion criteria and were randomized in a 2:1 ratio; 87 patients received TIF with EsophyX®-2 combined with 6 months of placebo (TIF/placebo) and 42 patients received sham surgery with 6 months of daily PPI therapy (sham/PPI). The primary outcome measure was elimination of troublesome regurgitation, defined as mild symptoms for 2 or more days per week or moderate-to-severe symptoms for more than 1 day per week. Crossover was allowed at 3 months in the case of treatment failure or at 6 months when the blind was broken. Lack of response at 3 months was observed in 36% of patients in the sham/PPI group compared with 11% in the TIF/placebo group ($p=0.002$). Self-reported regurgitation was eliminated in 22% more patients following TIF compared to continued PPI therapy patients (67% vs 45%, $p=0.023$), while reductions in GERD symptoms scores were similar in the 2 groups. The objective measure of control of esophageal pH was significantly reduced after TIF (mean percent time esophageal pH <4 decreased from 9.3% to 6.3%, $p<0.001$), but not after sham surgery (from 8.6% to 8.9%). By the 18-month follow-up, 71% of patients in the sham/PPI group had crossed over to TIF, compared with 28% of patients in the TIF/placebo group who resumed PPI therapy ($p<0.001$). There were 5 moderate-to-severe complications in the TIF group compared to one in the sham group. Strengths of this study include the use of both sham surgery and placebo control to maintain double-blinding, adequate power, objective as well as subjective outcome measures, and use of intention-to-treat analysis. A limitation is the relatively short duration of follow-up for most outcome measures.

Several other RCTs from 2015 have evaluated TIF in patients whose symptoms are at least partially controlled by PPI therapy.

Hakonsson reported a double-blind, sham-controlled randomized trial with 44 patients who had moderate-to-severe GERD symptoms without PPI therapy.^[17] Controls received a sham procedure, and the primary outcome was the time in remission, which was longer following TIF than sham (197 days vs 107 days, $p<0.0001$). Secondary outcomes measuring GERD symptoms showed results consistent with more favorable outcomes in the TIF group, however, no statistical between-group analysis was reported for these outcomes. Dysphagia, bloating, and flatulence were reported in twice as many patients undergoing TIF (4, 4, and 2 respectively) compared with sham (2, 2, and 1, respectively). These were reported as not statistically different, however, it is unlikely that the study was powered to detect differences in these outcomes.

Witteman reported an unplanned interim analysis of an RCT of 60 patients randomized to TIF using EsophyX®-2 or continued PPI therapy.^[18] Sixty of the planned 120 patients had been recruited at the time of analysis. The patients' symptoms were adequately controlled by PPIs but they wanted to avoid lifelong PPI therapy. At 6 months, subjective GERD symptoms improved to a greater extent in the TIF group ($p<0.001$), and satisfaction scores were higher (50% satisfied vs 0%), but there was no significant difference in esophageal acid exposure ($p=0.228$) or pH normalization (50% vs 63%) between the TIF and PPI groups, respectively. At 12 months after TIF, normalization of pH was achieved in only 29% of patients and there was deteriorated valve appearance at endoscopy; 61% of TIF patients had resumed use of PPIs.

Trad reported 6- and 12-month results of an industry-funded, multicenter RCT (TEMPO) that compared TIF using EsophyX®-2 ($n=40$) versus maximal dose PPI therapy ($n=23$) in partial responders to PPI therapy.^[14,15] At the 6-month follow-up, the subjective measure of

troublesome regurgitation was eliminated in 97% of TIF patients versus 50% of PPI patients (relative risk, 1.9; $p=0.006$). At 6 months, 90% of patients in the TIF group had completely stopped PPI therapy. However, the objective measure of normalized esophageal acid exposure did not differ significantly between groups (TIF=54% vs PPI=52%, $p=0.914$). At 12 months after TIF, 77% of patients had symptom control, 82% had stopped PPI therapy, 100% had healed esophagitis, and 45% had normalized esophageal acid exposure.

Additional controlled trials (RCTs) comparing transesophageal endoscopic gastroplasty or plication procedures to sham or other endoscopic procedures have been identified.^[15,19-24] Though these studies showed a promising decrease in PPI use and symptom control at 3 to 12 months, they do not allow conclusions regarding long-term health outcomes, safety or durability of the procedure in patients with GERD for one or more of the following reasons:

Insufficient study durations – Only short-term follow-up of 3 to 12 months is available, which does not address the long-term safety and durability of the procedures.^[15,20-25] For example, there may be suture loss over time. One study reported up to 29 % of study subjects required a second procedure at 12-month follow-up.^[20] Of these patients, 72% of sutures were still present but only 19% were judged functional. A second study noted marked loss of sutures with 67% remaining at 12 months.^[22]

Small sample size – Given the prevalence of GERD in the general population, available randomized trials include very small sample sizes. The largest study of 159 patients had an almost 10% loss in reported data with an intention to treat analysis that did not include these patients. All other studies include sample sizes of 60 or fewer patients. It is unclear if these studies are adequately powered.^[15,20,22-26]

Unreliable endpoints – The use of subjective, point in time GERD questionnaires as a primary endpoint may give variable results depending upon symptoms present at the time the subject completes the questionnaire.^[15,20,21]

Improvement over the gold standard procedures was not demonstrated. In order to establish the efficacy of transoral procedures, an improvement in symptoms of gastric reflux over the current open or laparoscopic anti-reflux procedures, must be shown.^[15,24,26]

There is a single randomized trial of the TIF procedure, which compares TIF to Nissen laparoscopic fundoplication.^[25] Although the authors reported comparable results at 12 months, conclusions based upon this trial are limited by the small sample size ($n=52$) and the different methods used for TIF (both the Plicator® and the EsophyX).

Nonrandomized Studies

Observational studies^[27-58], registry data^[59,60] nonrandomized comparative studies^[61] of gastroplication and fundoplication (specifically, transoral incisionless fundoplication) procedures do not allow conclusions about their long-term effectiveness and durability.

Harms

Of note, although harms are not systematically reported across observational studies, Furnee reported an increased risk of gastric injury with laparoscopic Nissen fundoplication after failed EsophyX fundoplication.^[62] Of 88 patients in their database who underwent EsophyX fundoplication, 11 (12.5%) subsequently underwent Nissen fundoplication for persistent or recurrent symptoms at a mean 8.1 months after the primary procedure. Endoscopy showed

partial or total disruption of fasteners in 8 of the 11 patients (72.7%). Nissen fundoplication after EsophyX resulted in gastric perforation (n=2), conversion to laparotomy (n=1), subphrenic abscess requiring surgical exploration (n=1) and symptom-worsening in 4 patients.

In 2017, Huang conducted a systematic review with meta-analysis of TIF for the treatment of GERD.^[63] Authors included 5 RCTs and 13 prospective observational studies, of which 14 were performed with the TIF 2 procedure. Efficacy results from the RCTs were combined for patients whose symptoms were controlled by PPIs and for those whose symptoms were not controlled by PPIs, and are not further discussed here. Follow-up out to 6 years in prospective observational studies indicated a decrease in efficacy over time. The reported incidence of severe adverse events, consisting of gastrointestinal perforation and bleeding, was 19 (2.4%) out of 781 patients. This included 7 perforations, 5 cases of post-TIF bleeding, 4 cases of pneumothorax, 1 case requiring intravenous antibiotics, and 1 case of severe epigastric pain.

TRANSESOPHAGEAL RADIOFREQUENCY ENERGY (I.E., THE STRETTA PROCEDURE)

Systematic Reviews

Fass (2017) published a meta-analysis of cohort studies and RCTs evaluating the Stretta procedure for patients with GERD (N=2468 total, 9-558 per study).^[64] The meta-analysis included 4 RCTs, 23 cohort studies, and 1 registry. Follow-up time varied from 3-120 months. When RCT and cohort results were pooled, there were clinically significant treatment effects for several of end points; however, the analysis was limited by the lack of control groups in many studies. Also, only 1 end point was shared between the four included RCTs.

A meta-analysis of four RCTs (total N=165 patients) was published by Lipka in 2015.^[65] Three trials compared Stretta with sham, and one trial compared Stretta with PPI therapy. Results of the individual sham-controlled trials were inconsistent, generally supporting some improvement in symptoms, but not in objective measures of esophageal acid exposure. For example, Corley (2008) reported improvement in heartburn symptoms, quality of life, and general physical quality of life in the active treatment group compared with the sham group, but there were no significant differences in medication use and esophageal acid exposure.^[66] Aziz (2010) found statistically significant improvements in GERD-HRQL in all treatment groups.^[67] Arts (2012) reported that the symptom score and quality-of-life score for bodily pain improved, but no changes were observed in PPI use, esophageal acid exposure, or lower esophageal sphincter pressure after RF.^[68] Pooled results of the meta-analysis showed no significant difference between Stretta and either sham treatment or PPI management for the measured outcomes, including the ability to stop PPI therapy. The overall quality of evidence was considered to be very low with a high risk of bias, and the meta-analysis was limited by heterogeneity in the included studies, which may be due to small sample sizes, differences in measures, and differences in follow-up time.

A 2014 systematic review and meta-analysis of four randomized trials; three reviewed previously^[66-68] and one trial which compared Stretta with PPI therapy,^[69] included a total of 165 patients. The overall quality of the evidence was considered to be very low with a high risk of bias. The pooled results showed no significant difference between Stretta and sham or PPI management for the measured outcomes. The meta-analysis was limited by heterogeneity in the included studies, which may be due to small sample sizes, differences in measures, and differences in follow-up time. The author also identified significant risks associated with Stretta, including pneumonia, gastroparesis, esophageal perforation, cardiac arrest, and at least 4 deaths from review of the Manufacturer and User Facility Device Experience database.

A meta-analysis completed by Perry, included 20 studies, only 2 of which were RCTs. This meta-analysis was limited by the inclusion of lower quality studies and by the analysis, which only examined within-subject differences and did not include between-subject differences, as reported in the RCTs.^[70]

Randomized Controlled Trials

There are 4 randomized trials comparing transesophageal radiofrequency (RF) energy with a sham procedure that involved balloon inflation but no needle deployment or RF energy delivery.^[66-68]

Results of the first study failed to include 20% of the randomized patients in analysis of primary endpoints, and no intention to treat analysis was provided. Therefore, reported results of improved heartburn symptoms and GERD quality of life scores are not reliable.

Results of the second, third and fourth studies were flawed due to a small patient population and inadequate timeframe for follow up.

Other small RCT's have been published. Two compared RF to PPI therapy. One trial showed promising short-term (6 months) results but does not permit conclusions about mid- to long-term effectiveness and durability.^[71] Another compared RF with PPI therapy to PPI therapy alone.^[72] Results at 3 months appeared favorable to the Stretta group, however, the study sample was small (N=20) and power calculations were not conducted.

Nonrandomized Studies

Other clinical studies concerning transesophageal radiofrequency are limited to observational case series that do not allow conclusions about long-term effectiveness and durability.^[19,73-84] Though several case series report up to 4-10 year outcomes, there was a significant loss to follow-up in these studies such that conclusions on durability and health outcomes cannot be made.^[85]

INJECTION OR IMPLANTATION OF BIOCOMPATIBLE POLYMERS

Randomized Controlled Trials

The available evidence for the Gatekeeper Reflux Repair System consists of one RCT. An industry-funded sham-controlled single-blind multicenter study randomized 118 patients into Gatekeeper (n=75) or sham (n=43) treatment.^[86] An additional 25 patients were treated as lead-ins during the initial training of investigators and included only in the safety analysis. The patients were implanted initially with 4 Gatekeeper prostheses. At three months, 44% of implanted patients received retreatment with up to four additional prostheses due to unsatisfactory symptom control. The primary safety end point was reduction in serious device- and procedure-related adverse device effects, compared with a surgical procedure composite complication rate of 15%. Four serious adverse events were reported (2 perforations, 1 pulmonary infiltrate related to a perforation, 1 severe chest pain). The primary efficacy end point was reduction in heartburn symptoms using the GERD-HRQL questionnaire. Planned interim analysis after 143 patients were enrolled found that heartburn symptoms and esophageal acid exposure had improved significantly in both the Gatekeeper and sham groups at 6 months, but there was no significant difference between the 2 groups. The study was terminated early due to a lack of efficacy.

There is one randomized sham-controlled trial which reports results of patients randomized to receive either injection of Enteryx biopolymer or a sham procedure.^[87] At 3- and 6-months follow-up, patients in the Enteryx group had greater reductions in PPI use and more improvement in GERD health-related quality of life heartburn scores. However, the small size and short duration of the study limit interpretation of findings.

Nonrandomized Studies

Other data on injectable or implantable polymers consists of very small case series.^[19,88] The small number of patients and lack of long-term follow-up precludes scientific analysis.

PRACTICE GUIDELINE SUMMARY

Several clinical practice guidelines consider the use of transoral fundoplication or other endoscopic procedures, although none were able to recommend this treatment based upon high level evidence.

AMERICAN SOCIETY OF GENERAL SURGEONS

The American Society of General Surgeons (ASGS) consensus-based position statement on transoral fundoplication states, “the ASGS supports the use of transoral fundoplication by trained General Surgeons for the treatment of symptomatic chronic gastroesophageal reflux disease (GERD) in patients who fail to achieve satisfactory response to a standard dose of Proton Pump Inhibitor (PPI) therapy or for those who wish to avoid the need for a lifetime of medication dependence.”^[89]

AMERICAN GASTROENTEROLOGICAL ASSOCIATION

The 2008 Medical Position Statement of the American Gastroenterological Association (AGA), makes no recommendation for or against “the use of currently commercially available endoluminal antireflux procedures in the management of patients with an esophageal syndrome” based on insufficient evidence (Grade Insufficient).^[90]

AMERICAN COLLEGE OF GASTROENTEROLOGY

In 2013, the ACG released updated guidelines stating that the usage of current endoscopic therapy or transoral incisionless fundoplication cannot be recommended as an alternative to medical or traditional surgical therapy.^[1]

SOCIETY OF AMERICAN GASTROINTESTINAL ENDOSCOPIC SURGEONS

In 2017, the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) updated its evidence-based guidelines on endoluminal treatments for GERD.^[91] SAGES gave a strong recommendation based on moderate quality evidence that TIF with EsoPHYX can be performed with an acceptable safety risk in selected patients. SAGES concluded that EsoPHYX results in better control of GERD symptoms compared with proton pump inhibitor (PPI) treatment in the short term (6 months), but leads to similar improvement in objective GERD measures compared with PPIs. TIF appears to lose effectiveness during longer term follow-up and is associated with moderate patient satisfaction scores. SAGES found no comparative, controlled trials between TIF and surgical fundoplication, but preliminary evidence suggested that the surgical fundoplication can be used safely after TIF failure. SAGES gave a strong recommendation based on moderate quality evidence that Stretta is safe for adults and

significantly improves health-related quality of life score, heartburn scores, the incidence of esophagitis, and esophageal acid exposure in patients with GERD. Stretta is more effective than PPI, but less so than fundoplication.

SUMMARY

There is not enough research to show that transesophageal endoscopic therapies for the treatment of gastroesophageal reflux disease (GERD) improves health outcomes. Although clinical guidelines based on research may recommend treating GERD with one or more of the therapies mentioned, there is not enough research to know if or how well these procedures work to treat people with GERD. This does not mean that it does not work, but more research is needed to know. Therefore, the use of any of these procedures is considered investigational for the treatment of GERD.

REFERENCES

1. Katz, PO, Gerson, LB, Vela, MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. 2013 Mar;108(3):308-28; quiz 29. PMID: 23419381
2. van Pinxteren, B, Sigterman, KE, Bonis, P, Lau, J, Numans, ME. Short-term treatment with proton pump inhibitors, H2-receptor antagonists and prokinetics for gastro-oesophageal reflux disease-like symptoms and endoscopy negative reflux disease. *The Cochrane database of systematic reviews*. 2010(11):CD002095. PMID: 21069670
3. U.S. Food and Drug Administration (FDA), 510(k) Summary: EsophyX K160960. May 1, 2016. [cited 12/13/2017]; Available from: https://www.accessdata.fda.gov/cdrh_docs/pdf16/K160960.pdf
4. U.S. Food and Drug Administration (FDA), 510(k) Summary: EsophyX K171307. June 22, 2016. [cited 12/13/2017]; Available from: https://www.accessdata.fda.gov/cdrh_docs/pdf17/K171307.pdf
5. Ip, S, Bonis, P, Tatsioni, A, et al. Comparative Effectiveness of Management Strategies For Gastroesophageal Reflux Disease. *AHRQ Comparative Effectiveness Reviews*. 2005. PMID: 21348043
6. Ip, S, Chung, M, Moorthy, D, et al. Comparative Effectiveness of Management Strategies for Gastroesophageal Reflux Disease: Update. *AHRQ Comparative Effectiveness Reviews*. 2011 Sep Disease: Update;Report No.: 11-EHC049-EF. PMID: 22091471
7. Humphries, LA, Hernandez, JM, Clark, W, Luberic, K, Ross, SB, Rosemurgy, AS. Causes of dissatisfaction after laparoscopic fundoplication: the impact of new symptoms, recurrent symptoms, and the patient experience. *Surg Endosc*. 2013 May;27(5):1537-45. PMID: 23508812
8. Hummel, K, Richards, W. Endoscopic treatment of gastroesophageal reflux disease. *The Surgical clinics of North America*. 2015 Jun;95(3):653-67. PMID: 25965137
9. Chen, D, Barber, C, McLoughlin, P, Thavaneswaran, P, Jamieson, GG, Maddern, GJ. Systematic review of endoscopic treatments for gastro-oesophageal reflux disease. *Br J Surg*. 2009 Feb;96(2):128-36. PMID: 19160349
10. McCarty, TR, Itidiare, M, Njei, B, Rustagi, T. Efficacy of transoral incisionless fundoplication for refractory gastroesophageal reflux disease: a systematic review and meta-analysis. *Endoscopy*. 2018 Jul;50(7):708-25. PMID: 29625507

11. Richter, JE, Kumar, A, Lipka, S, Miladinovic, B, Velanovich, V. Efficacy of Laparoscopic Nissen Fundoplication vs Transoral Incisionless Fundoplication or Proton Pump Inhibitors in Patients With Gastroesophageal Reflux Disease: A Systematic Review and Network Meta-analysis. *Gastroenterology*. 2018 Apr;154(5):1298-308 e7. PMID: 29305934
12. Trad, KS, Barnes, WE, Prevou, ER, et al. The TEMPO Trial at 5 Years: Transoral Fundoplication (TIF 2.0) Is Safe, Durable, and Cost-effective. *Surg Innov*. 2018 Apr;25(2):149-57. PMID: 29405886
13. Trad, KS, Fox, MA, Simoni, G, et al. Transoral fundoplication offers durable symptom control for chronic GERD: 3-year report from the TEMPO randomized trial with a crossover arm. *Surg Endosc*. 2016 Sep 21. PMID: 27655380
14. Trad, KS, Barnes, WE, Simoni, G, et al. Transoral incisionless fundoplication effective in eliminating GERD symptoms in partial responders to proton pump inhibitor therapy at 6 months: the TEMPO Randomized Clinical Trial. *Surg Innov*. 2015 Feb;22(1):26-40. PMID: 24756976
15. Trad, KS, Simoni, G, Barnes, WE, et al. Efficacy of transoral fundoplication for treatment of chronic gastroesophageal reflux disease incompletely controlled with high-dose proton-pump inhibitors therapy: a randomized, multicenter, open label, crossover study. *BMC gastroenterology*. 2014;14:174. PMID: 25284142
16. Hunter, JG, Kahrilas, PJ, Bell, RC, et al. Efficacy of transoral fundoplication vs omeprazole for treatment of regurgitation in a randomized controlled trial. *Gastroenterology*. 2015 Feb;148(2):324-33 e5. PMID: 25448925
17. Hakansson, B, Montgomery, M, Cadiere, GB, et al. Randomised clinical trial: transoral incisionless fundoplication vs. sham intervention to control chronic GERD. *Aliment Pharmacol Ther*. 2015 Dec;42(11-12):1261-70. PMID: 26463242
18. Witteman, BP, Conchillo, JM, Rinsma, NF, et al. Randomized controlled trial of transoral incisionless fundoplication vs. proton pump inhibitors for treatment of gastroesophageal reflux disease. *Am J Gastroenterol*. 2015 Apr;110(4):531-42. PMID: 25823768
19. TEC Assessment 2003. "Transesophageal Endoscopic Treatment for Gastroesophageal Reflux Disease." BlueCross BlueShield Association Technology Evaluation Center, Vol. 18, Tab 20.
20. Schwartz, MP, Wellink, H, Gooszen, HG, Conchillo, JM, Samsom, M, Smout, AJ. Endoscopic gastroplication for the treatment of gastro-oesophageal reflux disease: a randomised, sham-controlled trial. *Gut*. 2007 Jan;56(1):20-8. PMID: 16763053
21. Rothstein, R, Filipi, C, Caca, K, et al. Endoscopic full-thickness plication for the treatment of gastroesophageal reflux disease: A randomized, sham-controlled trial. *Gastroenterology*. 2006 Sep;131(3):704-12. PMID: 16952539
22. Montgomery, M, Hakanson, B, Ljungqvist, O, Ahlman, B, Thorell, A. Twelve months' follow-up after treatment with the EndoCinch endoscopic technique for gastro-oesophageal reflux disease: a randomized, placebo-controlled study. *Scand J Gastroenterol*. 2006 Dec;41(12):1382-9. PMID: 17101568
23. Domagk, D, Menzel, J, Seidel, M, et al. Endoluminal gastroplasty (EndoCinch) versus endoscopic polymer implantation (Enteryx) for treatment of gastroesophageal reflux disease: 6-month results of a prospective, randomized trial. *Am J Gastroenterol*. 2006 Mar;101(3):422-30. PMID: 16542275
24. Hunter, JG, Kahrilas, PJ, Bell, RC, et al. Efficacy of Transoral Fundoplication vs Omeprazole for Treatment of Regurgitation in a Randomized Controlled Trial. *Gastroenterology*. 2014 Oct 13. PMID: 25448925

25. Svoboda, P, Kantorova, I, Kozumplik, L, et al. Our experience with transoral incisionless plication of gastroesophageal reflux disease: NOTES procedure. *Hepato-gastroenterology*. 2011 Jul-Aug;58(109):1208-13. PMID: 21937380
26. Kaindlstorfer, A, Koch, OO, Antoniou, SA, Asche, KU, Granderath, FA, Pointner, R. A randomized trial on endoscopic full-thickness gastroplication versus laparoscopic antireflux surgery in GERD patients without hiatal hernias. *Surg Laparosc Endosc Percutan Tech*. 2013;23:212-22. PMID: 23579521
27. Repici, A, Fumagalli, U, Malesci, A, Barbera, R, Gambaro, C, Rosati, R. Endoluminal fundoplication (ELF) for GERD using EsophyX: a 12-month follow-up in a single-center experience. *J Gastrointest Surg*. 2010 Jan;14(1):1-6. PMID: 19902310
28. Testoni, PA, Corsetti, M, Di Pietro, S, et al. Effect of transoral incisionless fundoplication on symptoms, PPI use, and ph-impedance refluxes of GERD patients. *World J Surg*. 2010 Apr;34(4):750-7. PMID: 20091308
29. Cadiere, GB, Rajan, A, Gernay, O, Himpens, J. Endoluminal fundoplication by a transoral device for the treatment of GERD: A feasibility study. *Surg Endosc*. 2008 Feb;22(2):333-42. PMID: 18071818
30. Filipi, CJ, Lehman, GA, Rothstein, RI, et al. Transoral, flexible endoscopic suturing for treatment of GERD: a multicenter trial. *Gastrointest Endosc*. 2001 Apr;53(4):416-22. PMID: 11275879
31. Swain, CP. Endoscopic suturing. *Baillieres Best Pract Res Clin Gastroenterol*. 1999 Apr;13(1):97-108. PMID: 11030637
32. Mahmood, Z, McMahon, BP, Arfin, Q, et al. Endocinch therapy for gastro-oesophageal reflux disease: a one year prospective follow up. *Gut*. 2003 Jan;52(1):34-9. PMID: 12477756
33. Thomson, M, Fritscher-Ravens, A, Hall, S, Afzal, N, Ashwood, P, Swain, CP. Endoluminal gastroplication in children with significant gastro-oesophageal reflux disease. *Gut*. 2004 Dec;53(12):1745-50. PMID: 15542508
34. Chadalavada, R, Lin, E, Swafford, V, Sedghi, S, Smith, CD. Comparative results of endoluminal gastroplication and laparoscopic antireflux surgery for the treatment of GERD. *Surg Endosc*. 2004 Feb;18(2):261-5. PMID: 14691698
35. Abou-Rebyeh, H, Hoepffner, N, Rosch, T, et al. Long-term failure of endoscopic suturing in the treatment of gastroesophageal reflux: a prospective follow-up study. *Endoscopy*. 2005 Mar;37(3):213-6. PMID: 15731936
36. Schiefke, I, Zabel-Langhennig, A, Neumann, S, Feisthammel, J, Moessner, J, Caca, K. Long term failure of endoscopic gastroplication (EndoCinch). *Gut*. 2005 Jun;54(6):752-8. PMID: 15888777
37. Pleskow, D, Rothstein, R, Kozarek, R, Haber, G, Gostout, C, Lembo, A. Endoscopic full-thickness plication for the treatment of GERD: long-term multicenter results. *Surg Endosc*. 2007 Mar;21(3):439-44. PMID: 17180259
38. Mahmood, Z, Byrne, PJ, McMahon, BP, et al. Comparison of transesophageal endoscopic plication (TEP) with laparoscopic Nissen fundoplication (LNF) in the treatment of uncomplicated reflux disease. *Am J Gastroenterol*. 2006 Mar;101(3):431-6. PMID: 16542276
39. Rothstein, RI. Endoscopic therapy of gastroesophageal reflux disease: outcomes of the randomized-controlled trials done to date. *J Clin Gastroenterol*. 2008 May-Jun;42(5):594-602. PMID: 18364577
40. Cadiere, GB, Buset, M, Muls, V, et al. Antireflux transoral incisionless fundoplication using EsophyX: 12-month results of a prospective multicenter study. *World J Surg*. 2008 Aug;32(8):1676-88. PMID: 18443855

41. Pleskow, D, Rothstein, R, Kozarek, R, et al. Endoscopic full-thickness plication for the treatment of GERD: Five-year long-term multicenter results. *Surg Endosc.* 2008 Feb;22(2):326-32. PMID: 18027032
42. Cadiere, GB, Van Sante, N, Graves, JE, Gawlicka, AK, Rajan, A. Two-year results of a feasibility study on antireflux transoral incisionless fundoplication using EsophyX. *Surg Endosc.* 2009 May;23(5):957-64. PMID: 19288158
43. von Renteln, D, Schiefke, I, Fuchs, KH, et al. Endoscopic full-thickness plication for the treatment of gastroesophageal reflux disease using multiple Plicator implants: 12-month multicenter study results. *Surg Endosc.* 2009 Aug;23(8):1866-75. PMID: 19440792
44. Velanovich, V. Endoscopic, endoluminal fundoplication for gastroesophageal reflux disease: initial experience and lessons learned. *Surgery.* 2010 Oct;148(4):646-51; discussion 51-3. PMID: 20708763
45. Bell, RC, Freeman, KD. Clinical and pH-metric outcomes of transoral esophagogastric fundoplication for the treatment of gastroesophageal reflux disease. *Surg Endosc.* 2011 Jun;25(6):1975-84. PMID: 21140170
46. Barnes, WE, Hoddinott, KM, Mundy, S, Williams, M. Transoral incisionless fundoplication offers high patient satisfaction and relief of therapy-resistant typical and atypical symptoms of GERD in community practice. *Surg Innov.* 2011 Jun;18(2):119-29. PMID: 21307014
47. Trad, KS, Turgeon, DG, Deljkich, E. Long-term outcomes after transoral incisionless fundoplication in patients with GERD and LPR symptoms. *Surg Endosc.* 2012 Mar;26(3):650-60. PMID: 21959689
48. Ihde, GM, Besancon, K, Deljkich, E. Short-term safety and symptomatic outcomes of transoral incisionless fundoplication with or without hiatal hernia repair in patients with chronic gastroesophageal reflux disease. *Am J Surg.* 2011 Dec;202(6):740-6; discussion 6-7. PMID: 22014853
49. Petersen, RP, Filippa, L, Wassenaar, EB, Martin, AV, Tatum, R, Oelschlager, BK. Comprehensive evaluation of endoscopic fundoplication using the EsophyX(TM) device. *Surg Endosc.* 2011 Nov 1. PMID: 22042587
50. Chen, S, Jarboe, MD, Teitelbaum, DH. Effectiveness of a transluminal endoscopic fundoplication for the treatment of pediatric gastroesophageal reflux disease. *Pediatr Surg Int.* 2012 Mar;28(3):229-34. PMID: 22124618
51. Testoni, PA, Vailati, C, Testoni, S, Corsetti, M. Transoral incisionless fundoplication (TIF 2.0) with EsophyX for gastroesophageal reflux disease: long-term results and findings affecting outcome. *Surg Endosc.* 2011 Dec 15. PMID: 22170317
52. Narsule, CK, Burch, MA, Ebright, MI, et al. Endoscopic fundoplication for the treatment of gastroesophageal reflux disease: initial experience. *J Thorac Cardiovasc Surg.* 2012 Jan;143(1):228-34. PMID: 22070927
53. Nguyen, A, Vo, T, Nguyen, XM, Smith, BR, Reavis, KM. Transoral incisionless fundoplication: initial experience in patients referred to an integrated academic institution. *Am Surg.* 2011 Oct;77(10):1386-9. PMID: 22127095
54. Muls, V, Eckardt, AJ, Marchese, M, et al. Three-year results of a multicenter prospective study of transoral incisionless fundoplication. *Surg Innov.* 2013;20:321-30. PMID: 22968006
55. Wendling, MR, Melvin, WS, Perry, KA. Impact of transoral incisionless fundoplication (TIF) on subjective and objective GERD indices: a systematic review of the published literature. *Surg Endosc.* 2013 Oct;27(10):3754-61. PMID: 23644835
56. Testoni, PA, Testoni, S, Mazzoleni, G, Vailati, C, Passaretti, S. Long-term efficacy of transoral incisionless fundoplication with Esophyx (Tif 2.0) and factors affecting

- outcomes in GERD patients followed for up to 6 years: a prospective single-center study. *Surg Endosc.* 2014 Dec 6. PMID: 25480624
57. Bell, RC, Barnes, WE, Carter, BJ, et al. Transoral incisionless fundoplication: 2-year results from the prospective multicenter U.S. study. *Am Surg.* 2014 Nov;80(11):1093-105. PMID: 25347499
 58. Stefanidis, G, Viazis, N, Kotsikoros, N, et al. Long-term benefit of transoral incisionless fundoplication using the esophyx device for the management of gastroesophageal reflux disease responsive to medical therapy. *Diseases of the esophagus : official journal of the International Society for Diseases of the Esophagus.* 2017 Feb 1;30(3):1-8. PMID: 27868281
 59. Bell, RC, Mavrelis, PG, Barnes, WE, et al. A Prospective Multicenter Registry of Patients with Chronic Gastroesophageal Reflux Disease Receiving Transoral Incisionless Fundoplication. *Journal of the American College of Surgeons.* 2012 Aug 29. PMID: 22939637
 60. Wilson, EB, Barnes, WE, Mavrelis, PG, et al. The effects of transoral incisionless fundoplication on chronic GERD patients: 12-month prospective multicenter experience. *Surg Laparosc Endosc Percutan Tech.* 2014 Feb;24(1):36-46. PMID: 24487156
 61. Frazzoni, M, Conigliaro, R, Manta, R, Melotti, G. Reflux parameters as modified by EsophyX or laparoscopic fundoplication in refractory GERD. *Aliment Pharmacol Ther.* 2011 Jul;34(1):67-75. PMID: 21539587
 62. Furnee, EJ, Broeders, JA, Draaisma, WA, et al. Laparoscopic Nissen fundoplication after failed EsophyX fundoplication. *Br J Surg.* 2010 Jul;97(7):1051-5. PMID: 20632271
 63. Huang, X, Chen, S, Zhao, H, et al. Efficacy of transoral incisionless fundoplication (TIF) for the treatment of GERD: a systematic review with meta-analysis. *Surg Endosc.* 2017 Mar;31(3):1032-44. PMID: 27495332
 64. Fass, R, Cahn, F, Scotti, DJ, Gregory, DA. Systematic review and meta-analysis of controlled and prospective cohort efficacy studies of endoscopic radiofrequency for treatment of gastroesophageal reflux disease. *Surg Endosc.* 2017 Dec;31(12):4865-82. PMID: 28233093
 65. Lipka, S, Kumar, A, Richter, JE. No evidence for efficacy of radiofrequency ablation for treatment of gastroesophageal reflux disease: a systematic review and meta-analysis. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association.* 2015 Jun;13(6):1058-67 e1. PMID: 25459556
 66. Corley, DA, Katz, P, Wo, JM, et al. Improvement of gastroesophageal reflux symptoms after radiofrequency energy: a randomized, sham-controlled trial. *Gastroenterology.* 2003 Sep;125(3):668-76. PMID: 12949712
 67. Aziz, AM, El-Khayat, HR, Sadek, A, et al. A prospective randomized trial of sham, single-dose Stretta, and double-dose Stretta for the treatment of gastroesophageal reflux disease. *Surg Endosc.* 2010 Apr;24(4):818-25. PMID: 19730952
 68. Arts, J, Bisschops, R, Blondeau, K, et al. A Double-Blind Sham-Controlled Study of the Effect of Radiofrequency Energy on Symptoms and Distensibility of the Gastro-Esophageal Junction in GERD. *Am J Gastroenterol.* 2012 Feb;107(2):222-30. PMID: 22108449
 69. BlueCross BlueShield Association Medical Policy Reference Manual "Transesophageal Endoscopic Therapies for Gastroesophageal Reflux Disease." Policy No. 2.01.38
 70. Perry, KA, Banerjee, A, Melvin, WS. Radiofrequency energy delivery to the lower esophageal sphincter reduces esophageal acid exposure and improves GERD

- symptoms: a systematic review and meta-analysis. *Surg Laparosc Endosc Percutan Tech*. 2012 Aug;22(4):283-8. PMID: 22874675
71. Coron, E, Sebillé, V, Cadiot, G, et al. Clinical trial: Radiofrequency energy delivery in proton pump inhibitor-dependent gastro-oesophageal reflux disease patients. *Aliment Pharmacol Ther*. 2008 Nov 1;28(9):1147-58. PMID: 18616516
 72. Kalapala, R, Shah, H, Nabi, Z, Darisetty, S, Talukdar, R, Nageshwar Reddy, D. Treatment of gastroesophageal reflux disease using radiofrequency ablation (Stretta procedure): An interim analysis of a randomized trial. *Indian journal of gastroenterology : official journal of the Indian Society of Gastroenterology*. 2017 Sep;36(5):337-42. PMID: 29030802
 73. Triadafilopoulos, G, DiBaise, JK, Nostrant, TT, et al. The Stretta procedure for the treatment of GERD: 6 and 12 month follow-up of the U.S. open label trial. *Gastrointest Endosc*. 2002 Feb;55(2):149-56. PMID: 11818914
 74. Richards, WO, Scholz, S, Khaitan, L, Sharp, KW, Holzman, MD. Initial experience with the Stretta procedure for the treatment of gastroesophageal reflux disease. *J Laparoendosc Adv Surg Tech A*. 2001 Oct;11(5):267-73. PMID: 11642661
 75. DiBaise, JK, Brand, RE, Quigley, EM. Endoluminal delivery of radiofrequency energy to the gastroesophageal junction in uncomplicated GERD: efficacy and potential mechanism of action. *Am J Gastroenterol*. 2002 Apr;97(4):833-42. PMID: 12003416
 76. Triadafilopoulos, G. Stretta: an effective, minimally invasive treatment for gastroesophageal reflux disease. *Am J Med*. 2003 Aug 18;115 Suppl 3A:192S-200S. PMID: 12928101
 77. Vakil, N, Sharma, P. Review article: endoscopic treatments for gastro-oesophageal reflux disease. *Aliment Pharmacol Ther*. 2003 Jun 15;17(12):1427-34. PMID: 12823144
 78. Houston, H, Khaitan, L, Holzman, M, Richards, WO. First year experience of patients undergoing the Stretta procedure. *Surg Endosc*. 2003 Mar;17(3):401-4. PMID: 12436238
 79. Noar, MD, Lotfi-Emran, S. Sustained improvement in symptoms of GERD and antisecretory drug use: 4-year follow-up of the Stretta procedure. *Gastrointest Endosc*. 2007 Mar;65(3):367-72. PMID: 17321232
 80. Reymunde, A, Santiago, N. Long-term results of radiofrequency energy delivery for the treatment of GERD: sustained improvements in symptoms, quality of life, and drug use at 4-year follow-up. *Gastrointest Endosc*. 2007 Mar;65(3):361-6. PMID: 17321231
 81. McClusky, DA, 3rd, Khaitan, L, Swafford, VA, Smith, CD. Radiofrequency energy delivery to the lower esophageal sphincter (Stretta procedure) in patients with recurrent reflux after antireflux surgery: can surgery be avoided? *Surg Endosc*. 2007 Jul;21(7):1207-11. PMID: 17308947
 82. Dundon, JM, Davis, SS, Hazey, JW, Narula, V, Muscarella, P, Melvin, WS. Radiofrequency energy delivery to the lower esophageal sphincter (Stretta procedure) does not provide long-term symptom control. *Surg Innov*. 2008 Dec;15(4):297-301. PMID: 18829607
 83. Liu, HF, Zhang, JG, Li, J, Chen, XG, Wang, WA. Improvement of clinical parameters in patients with gastroesophageal reflux disease after radiofrequency energy delivery. *World J Gastroenterol*. 2011 Oct 21;17(39):4429-33. PMID: 22110270
 84. Dughera, L, Navino, M, Cassolino, P, et al. Long-Term Results of Radiofrequency Energy Delivery for the Treatment of GERD: Results of a Prospective 48-Month Study. *Diagn Ther Endosc*. 2011;2011:507157. PMID: 22110288

85. Noar, M, Squires, P, Noar, E, Lee, M. Long-term maintenance effect of radiofrequency energy delivery for refractory GERD: a decade later. *Surg Endosc.* 2014 Aug;28(8):2323-33. PMID: 24562599
86. Fockens, P, Cohen, L, Edmundowicz, SA, et al. Prospective randomized controlled trial of an injectable esophageal prosthesis versus a sham procedure for endoscopic treatment of gastroesophageal reflux disease. *Surg Endosc.* 2010 Jun;24(6):1387-97. PMID: 20198491
87. Deviere, J, Costamagna, G, Neuhaus, H, et al. Nonresorbable copolymer implantation for gastroesophageal reflux disease: a randomized sham-controlled multicenter trial. *Gastroenterology.* 2005 Mar;128(3):532-40. PMID: 15765387
88. Feretis, C, Benakis, P, Dimopoulos, C, et al. Endoscopic implantation of Plexiglas (PMMA) microspheres for the treatment of GERD. *Gastrointest Endosc.* 2001 Apr;53(4):423-6. PMID: 11275880
89. American Society of General Surgeons (ASGS). Position statement: Transoral fundoplication. [cited 12/13/2017]; Available from: <https://theasgs.org/position-statements/american-society-of-general-surgeons-asgs-position-statement-transoral-fundoplication/>
90. Kahrilas, PJ, Shaheen, NJ, Vaezi, MF, et al. American Gastroenterological Association Medical Position Statement on the management of gastroesophageal reflux disease. *Gastroenterology.* 2008 Oct;135(4):1383-91, 91 e1-5. PMID: 18789939
91. Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) Guidelines. May 2017 Clinical Spotlight Review: Endoluminal Treatments for Gastroesophageal Reflux Disease (GERD). May 2017. [cited 12/13/2017]; Available from: <https://www.sages.org/publications/guidelines/endoluminal-treatments-for-gastroesophageal-reflux-disease-gerd/>

CODES

Codes	Number	Description
CPT	43192	Esophagoscopy; rigid, transoral; with directed submucosal injection(s), any substance
	43201	Esophagoscopy; flexible, transoral; with directed submucosal injection(s), any substance
	43210	Esophagogastroduodenoscopy, flexible, transoral; with esophagogastric fundoplasty, partial or complete, includes duodenoscopy when performed
	43236	Esophagogastroduodenoscopy, flexible, transoral, with direct submucosal injections, any substance
	43257	Esophagogastroduodenoscopy, flexible transoral; with deliver of thermal esophageal sphincter and/or gastric cardia, for treatment of gastroesophageal reflux disease
	43499	Unlisted procedure, esophagus
HCPCS	None	

Date of Origin: February 2001