

Regence

Medical Policy Manual

Surgery, Policy No. 107

Percutaneous Vertebroplasty, Kyphoplasty, Sacroplasty, and Coccygeoplasty

Effective: August 1, 2024

Next Review: April 2025

Last Review: June 2024

IMPORTANT REMINDER

Medical Policies are developed to provide guidance for members and providers regarding coverage in accordance with contract terms. Benefit determinations are based in all cases on the applicable contract language. To the extent there may be any conflict between the Medical Policy and contract language, the contract language takes precedence.

PLEASE NOTE: Contracts exclude from coverage, among other things, services or procedures that are considered investigational or cosmetic. Providers may bill members for services or procedures that are considered investigational or cosmetic. Providers are encouraged to inform members before rendering such services that the members are likely to be financially responsible for the cost of these services.

DESCRIPTION

These procedures involve the injection of a polymethylmethacrylate (PMMA), a cement, into a fractured or weakened vertebral body to provide stabilization as an alternative to spinal fusion.

MEDICAL POLICY CRITERIA

- I. Mechanical vertebral augmentation using SpineJack or Kiva[®], percutaneous vertebroplasty, or balloon kyphoplasty may be considered **medically necessary** when one or more of the following criteria (A or B) are met:
 - A. Treatment of *no more than* three symptomatic vertebral fractures of the T5-L5 spine, on any single date of service, when **all** of the following (1 – 5) criteria are met:
 1. Appropriate imaging (plain film x-ray, MRI, CT, or bone scan) has been performed preoperatively and the findings of such imaging correlate with the patient's pain; and
 2. Patient's pain is more likely than not, related to the demonstrated fracture(s); and

3. Functional impairment attributed to vertebral fracture is documented in the clinical record as limiting performance of instrumental activities of daily living (ADLs). Instrumental ADLs are defined as feeding, bathing, dressing, grooming, meal preparation, household chores, and occupational tasks that are required as a daily part of job functioning.

Clinical records must specifically document the following:

- a. The specific instrumental ADL(s) that is(are) impaired; and
 - b. A description of how performance of the instrumental ADL is limited; and
4. The patient has failed to respond to conservative treatment (e.g., analgesics, physical therapy, rest) for at least six weeks; and
5. A pre-procedure assessment has documented the absence of the following contraindications:
 - a. Untreated symptomatic foraminal or canal stenosis, facet arthropathy, or other significant coexistent spinal or bony pain generators at the planned treatment level; and
 - b. Bone fragment retropulsion; and
 - c. Symptoms that cannot be related to a fracture; and
 - d. Unstable fracture or requirement for stabilization procedure in same or adjacent spinal region; and
 - e. Active osteomyelitis whether fungal, bacterial or mycobacterial, or any other active infection, including urinary tract infection (UTI); and
 - f. Presence of painful metastases to areas other than the spine, spinal cord compression, primary bone and osteoblastic tumors, solitary plasmacytomas; and
 - g. Uncorrected coagulation disorders; and
 - h. Known allergy to any of the materials used in these procedures; and
 - i. Chronic fracture at the same vertebral level (defined as greater than six months); or

- B. Treatment of symptomatic osteoporotic vertebral fractures that are less than 6 weeks in duration that have led to hospitalization or symptoms documented to persist at a level that prevents ambulation or transfers without assistance.

- II. Mechanical vertebral augmentation using SpineJack or Kiva[®], percutaneous vertebroplasty, or balloon kyphoplasty is considered **investigational** for all other indications, including but not limited to vertebral hemangioma, acute vertebral fractures due to trauma, vertebrae of the cervical spine and thoracic levels T1-5, stabilization of insufficiency fractures or lesions of the sacrum (sacroplasty) or coccyx (coccygeoplasty), and prophylactic treatment for osteoporosis of the spine or for chronic back pain of long-standing duration, even if associated with old compression fracture(s).

- III. Percutaneous mechanical vertebral augmentation using any device other than a balloon device is considered **investigational**, including but not limited to radiofrequency-assisted vertebral augmentation with ultrahigh viscosity cement (e.g.

Radiofrequency-Targeted Vertebral Augmentation™ with the StabiliT® System) and vertebral body stenting.

NOTE: A summary of the supporting rationale for the policy criteria is at the end of the policy.

LIST OF INFORMATION NEEDED FOR REVIEW

It is critical that the list of information below is submitted for review to determine if the policy criteria are met. If any of these items are not submitted, it could impact our review and decision outcome.

- History and Physical/Chart Notes
- Current Symptomology
- Imaging completed (plain film x-ray, MRI, CT or bone scan)
- Description of functional impairment and how it limits instrumental ADLs
- Conservative treatments (e.g., analgesics, physical therapy, rest) attempted for at least six weeks and documented response
- Pre-procedure Assessment documenting absence of contraindications outlined in criteria I.A.5. a-i

CROSS REFERENCES

1. [Sacroiliac Joint Fusion](#), Surgery, Policy No. 193

BACKGROUND

Percutaneous vertebroplasty, vertebral balloon kyphoplasty, and mechanical augmentation have been proposed as options to provide mechanical support and symptomatic pain relief in patients with osteoporotic vertebral compression, insufficiency fractures, vertebral hemangioma, or osteolytic lesions of the spine (i.e., multiple myeloma or metastatic malignancies). These procedures, sometimes referred to as vertebral augmentation, have been used in all levels of the spinal column including the sacrum and coccyx. When vertebroplasty or kyphoplasty is used to treat insufficiency fractures of the sacrum or coccyx they may be referred to as sacroplasty or coccygeoplasty, respectively.

- Percutaneous vertebroplasty is an interventional radiology technique involving the fluoroscopic- or CT-guided injection of polymethylmethacrylate (PMMA) through a needle inserted into a weakened vertebral body.
- Percutaneous kyphoplasty is a variant of vertebroplasty that is intended to restore the vertebral body height and alignment along with stabilizing the fracture, using one of the following techniques:
 - Balloon kyphoplasty involves the use of a specialized bone tamp with an inflatable balloon to expand a collapsed vertebral body. PMMA is then injected into the created cavity to stabilize the vertebral body.
 - Mechanical kyphoplasty describes techniques that do not involve a balloon device.
 - In radiofrequency kyphoplasty, ultrahigh viscosity cement is injected into the fractured vertebral body. Radiofrequency energy is used to achieve the desired cement consistency. The ultrahigh viscosity cement is intended to restore height and alignment to the fractured vertebra, along with stabilizing the fracture.

- The Kiva[®] procedure uses shaped memory coil and a Kiva implant inserted into the vertebral body for structural support and to provide a reservoir for injection of bone cement. The proposed benefit of this technique is the adjustable height of the implant, which is made from a biocompatible polymer (e.g., PEEK-OPTIMA[®]), and a potential reduction in cement leakage.
- SpineJack is a mechanical vertebral augmentation technique that utilizes bipedicular 4.2 mm to 5.0 mm self-expanding jacks to restore vertebral height. Placement of the titanium devices are verified in AP and lateral view prior to expansion. Once the devices are expanded, then a proprietary bone cement is injected. The proposed benefit is greater control over expansion and greater restoration of vertebral height compared to balloon kyphoplasty. The procedure requires good bone quality.
- Vertebral body stenting utilizes an expandable scaffold instead of a balloon to restore vertebral height. The proposed advantages of vertebral body stenting are to reduce the risk of cement leakage by formation of a cavity for cement application and to prevent the loss of correction that is seen following removal of the balloon used for balloon kyphoplasty.

Although the mechanism is unknown, percutaneous vertebroplasty and kyphoplasty are intended to provide analgesic effect either through mechanical stabilization of a fractured or otherwise weakened vertebral body or through thermal damage to intraosseous nerve fibers, since PMMA undergoes a heat-releasing (exothermic) reaction during its hardening process.

REGULATORY STATUS

Percutaneous vertebroplasty and kyphoplasty are surgical procedures and therefore, they are not subject to U.S. Food and Drug Administration (FDA) approval. However, the instruments and materials used within these procedures are subject to FDA approval. The Kiva[®] VCF Treatment System (Benvenue Medical) received FDA 510(k) marketing clearance in 2014. The SpineJack Expansion Kit (Vexim) was cleared for marketing by the FDA through the 510(k) process. These systems, or rather the instruments and materials, are approved under FDA product codes: NDN, OAR, and HXG.

Instruments

Various systems for percutaneous vertebral body access and delivery of bone cement for vertebroplasty have received FDA approval. The Parallax[®] Contour[®] Vertebral Augmentation device one such example. This device creates a void in cancellous bone that can then be filled with bone cement. The void is created by removal of bone fragments; unlike balloon kyphoplasty, this procedure does not attempt to restore vertebral body height.

Percutaneous kyphoplasty requires the use of an inflatable bone tamp. One such tamp, the KyphX[®] inflatable bone tamp, received 510(k) marketing clearance from the FDA in July 1998. Other devices with FDA 510(k) marketing clearance include AVAmax[®] Vertebral Balloon system (Carefusion), NeuroTherm Parallax[®] Balloon Inflatable Bone Tamp (NeuroTherm, Inc.), Stryker iVAS[®] Balloon catheter, and Synthes Synflate[™] Vertebral Balloon System (Synthes).

Bone cement

PMMA bone cement was available as a drug product prior to enactment of the FDA's device regulation and was at first considered what the FDA terms a "transitional device." It was

transitioned to a class III device requiring premarketing applications. Several orthopedic companies have received approval of their bone cement products since 1976. In October 1999, PMMA was reclassified from class III to class II, which requires future 510(k) submissions to meet “special controls” instead of “general controls” to assure safety and effectiveness. PMMA bone cements such as KyphX[®] HV-RTM, Kyphon[®] HV-R Bone Cement, Cortoss[®] Bone Augmentation Material, Spine-Fix[®] Biomimetic Bone Cement, StabiliT[®], and Osteopal[®] V were issued 510(k) marketing clearance for the fixation of pathological fractures of the vertebral body using vertebroplasty or kyphoplasty procedures.

Vesselplasty using Vessel-X[®], (MAXXSPINE) and a similar procedure from A-Spine, are variations of vertebroplasty that are reported to reduce leakage of bone cement by containing the filler in an inflatable vessel. FDA clearance these products have not been identified.

EVIDENCE SUMMARY

For treatment of vertebral body fractures related to osteoporosis or malignancy with percutaneous vertebroplasty (VP), kyphoplasty (KP), SpineJack, or Kiva, the primary beneficial outcomes of interest are relief of pain and improvement in ability to function. Data from large, blinded, randomized controlled trials (RCTs) of sufficient long-term follow-up are required to control for the placebo effect, determine its magnitude, and determine whether any treatment effect from vertebroplasty or kyphoplasty provides improves net health outcomes compared to sham or nonsurgical treatment. Further, adverse events, such as risk of additional fractures or cement leakage, must be considered in evaluating the benefits compared to the potential harms of these procedures. Therefore, the focus of the evidence summary below is on systematic reviews, technology assessments, RCTs, and clinical practice guidelines.

PERCUTANEOUS VERTEBROPLASTY

SYSTEMATIC REVIEWS

Zhu (2022) conducted a systematic review of six studies including data from 644 patients, 330 who received VP and 284 who received KP.^[1] There was no significant difference in either group in terms of visual analog scale (VAS) scores (MD = 0.17; 95% CI, -0.39 to 0.73; P = .56), risk of cement leakage (odds ratio [OR] = 1.31; 95% CI, 0.62 to 2.74; P = .47) or Oswestry Disability Index (ODI) scores (MD = 0.51; 95% CI, -1.87 to 2.88; P = .68). The injected cement volume (MD = -0.52; 95% CI, -0.88 to -0.15; P = .005) in the VP group was linked to a markedly lower trend compared with the KP group.

Halvachizadeh (2021) conducted a systematic review and meta-analysis comparing vertebroplasty, kyphoplasty, and nonoperative management in patients with osteoporotic vertebral compression fractures.^[2] A total of 16 RCTs (N=2731 patients) were included with 11 trials comparing vertebroplasty to nonoperative management, 1 trial comparing kyphoplasty to nonoperative management, and 4 comparing kyphoplasty and vertebroplasty. Surgical intervention was associated with greater improvement of pain as compared to nonoperative management and was unrelated to the development of adjacent level fractures or quality of life. Of the trials comparing kyphoplasty and vertebroplasty, no significant differences in outcome measures were observed.

Xie (2017), in a meta-analysis of RCTs, evaluated efficacy and safety in percutaneous vertebroplasty and conservative treatment for patients with osteoporotic vertebral compression fractures.^[3] Thirteen studies were selected (total N=1231 patients; 623 to vertebroplasty, 608

to conservative treatment); among them were the two sham-controlled trials described below. Outcomes included pain relief (from one week to six months), quality of life assessments, and the rate of adjacent-level vertebral fracture. Vertebroplasty was superior for pain relief at 1 week (mean difference [MD], 1.36; 95% CI, 0.55 to 2.17) and one month (MD=1.56; 95% CI, 0.43 to 2.70); it was inferior to conservative treatment for pain relief at six months (MD = -1.59; 95% CI, -2.9 to -0.27; $p < 0.05$). Vertebroplasty showed improvement over conservative treatment for quality of life, as measured using the Quality of Life Questionnaire of the European Foundation for Osteoporosis (MD = -5.03; 95% CI, 7.94 to -2.12). No statistically significant differences were found between treatments for the rate of adjacent-level vertebral fractures (relative risk: 0.59; 95% CI, 0.43 to 0.81). Limitations included inclusion of several studies with inadequate blinding and heterogenous reporting of patient characteristics outcomes.

A 2015 Cochrane review by Buchbinder evaluated the evidence on vertebroplasty for the treatment of vertebral compression fractures.^[4] Eleven RCTs and one quasi-RCT were included in the systematic review. Two trials identified compared vertebroplasty with a sham procedure (n=209 patients; Buchbinder [2009]^[5] and Kallmes [2009]^[6], detailed below), six compared vertebroplasty with usual care (n=566), and four compared vertebroplasty with kyphoplasty (n=545). The sham-controlled trials were at low risk of bias. All other trials were judged to be at high risk of bias due to lack of blinding. Evidence was rated as moderate quality based on the low number of subjects in the sham controlled trials. Meta-analysis of the two sham-controlled trials indicated that vertebroplasty does not result in clinically significant improvements in pain, disability, quality of life, or treatment success. Results did not differ for patients with pain durations of six weeks or less compared to pain lasting more than six weeks. Sensitivity analysis indicated that studies comparing vertebroplasty to conservative management were likely to have overestimated the treatment effect. The rate of serious adverse events did not differ significantly between the vertebroplasty and control groups, but serious adverse events related specifically to the vertebroplasty procedure included osteomyelitis, cord compression, thecal sac injury, and respiratory failure.

In 2011, Staples published a patient-level meta-analysis of the two sham-controlled trials to determine whether vertebroplasty is more effective than sham in specific subsets of patients.^[7] This subset analysis focused on duration of pain (< 6 weeks vs. > 6 weeks) and severity of pain (score < 8 or >8 on an 11-point numerical rating scale). Included in the analysis were 209 participants, 27% with pain of recent onset and 47% with severe pain at baseline. The primary outcome measures, pain scores and function on the RMDQ at one month, were not significantly different between groups. Responders' analyses were also conducted based on a 3-unit improvement in pain scores, a 3-unit improvement on the RMDQ, and a 30% improvement in each of the pain and disability outcomes. The only difference observed between groups was not statistically significant. This difference was a higher proportion of the vertebroplasty group achieving at least 30% improvement in pain scores, a result that may have been confounded by the greater use of opioid medications in that group. Overall, this analysis did not support the hypothesis that selected subgroups of patients, including those with pain of six weeks duration or less or those with severe pain, would benefit from vertebroplasty.

A systematic review of the safety and efficacy of vertebroplasty in malignancy was reported by Chew in 2011.^[8] Thirty relevant studies were identified, totaling 987 patients. Included in the review were a single randomized controlled trial and seven prospective studies. Most centers reported treating no more than four vertebrae per session. Pain reduction ranged from 20% to

79%. Five deaths were attributable to vertebroplasty, two from chest infections following general anesthesia, one from a cement pulmonary embolus, and two from sepsis after emergency spinal decompression. Another 19 patients suffered a serious complication related to the procedure, with 13 requiring emergency spinal decompression. Reports of complications occurred most in studies with a mean cement volume of more than 4 ml, suggesting a possible association between the volume of cement injected and increased risk of adverse events.

RANDOMIZED CONTROLLED TRIALS

Select randomized controlled trials not already included in the systematic reviews above are detailed in this section. Two other RCTs by Buchbinder (2015) and Kallmes (2009) were included in the systematic reviews above.^[4, 6]

Randomized Controlled Trials with Sham Controls

Firanesu (2018) published the results of a randomized, double-blind, sham-controlled clinical trial performed in four community hospitals in the Netherlands from 2011 to 2015.^[9] Participants included 180 patients requiring treatment for acute osteoporotic vertebral compression fractures that were randomized to either vertebroplasty (n=91) or a sham procedure (n=89). The main outcome measured was mean reduction in VAS scores at day one, one week, and one, three, six, and 12 months. The mean reduction in VAS score was statistically significant in the vertebroplasty and sham procedure groups at all follow-up points after the procedure compared with baseline. These changes in VAS scores did not, however, differ statistically significantly between the groups during 12 months' follow-up.

In 2016, Clark reported results from the VAPOUR trial.^[10] VAPOUR was a multicenter double-blind trial of vertebroplasty in 120 patients with vertebral fractures of less than six weeks in duration and back pain of at least 7 out of 10 on an NRS. Two authors had participated in the 2009 study published by Kallmes and the trial followed a similar protocol. Both outcomes assessors and patients were masked to treatment allocation, and independent statisticians unmasked the data and prepared the trial report. The sham-vertebroplasty procedure included subcutaneous lidocaine but no periosteal numbing. Manual skin pressure and tapping on the needle was performed to simulate the needle advance, and the investigators discussed polymethylmethacrylate (PMMA) mixing and injection during the procedure. The primary outcome, the percentage of patients with an NRS score less than 4 out of 10 at 14 days after the procedure, was met in a greater percentage of patients in the vertebroplasty group (44%) than in the sham control group (21%). This between-group difference was maintained through six months. Other outcome measures were significantly improved in the vertebroplasty group at one or both time points. The benefit of vertebroplasty was found predominantly in the thoracolumbar subgroup, with 48% (95% CI, 27% to 68%) more patients meeting the primary endpoint (61% in the vertebroplasty group vs 13% in the control group). The investigators commented that the thoracolumbar junction is subject to increased dynamic load, and fractures at this junction have the highest incidence of mobility. No benefit from vertebroplasty was found in the non-thoracolumbar subgroup. Postprocedural hospital stay was reduced from a mean of 14 days in the control group to 8.5 days after vertebroplasty, even though physicians who determined the discharge date remained blinded to treatment. In the vertebroplasty group, there were two serious adverse events due to sedation and transfer to the radiology table. In the control group, two patients developed spinal cord compression; one underwent decompressive surgery and the other, not a surgical candidate, became paraplegic.

Kroon (2014) reported the 12- and 24-month outcomes of a double-blind RCT comparing VP (n=38) with sham procedure (n=40) for acute osteoporotic vertebral fractures.^[11] The initial report of 6-month outcomes (Buchbinder 2009^[5]) found no benefit of VP over sham procedure. Complete data were available for 67 (86%) and 57 (73%) of participants at 12 and 24 months, respectively. VP patients had significantly higher overall pain reduction at both time points compared to sham. There were no significant between-group differences in disability, quality of life, perceived recovery, or adverse event including subsequent vertebral fractures. The authors concluded that these outcomes provide evidence that use of VP as routine care for vertebral fractures is unsupported. Methodological limitations in the RCT included small sample size that may have had inadequate power to show significant differences in subsequent vertebral fractures.

Randomized Controlled Trials without Sham Controls

In 2016, Leali published a short report of a multicenter RCT with 400 patients with osteoporotic thoracic or lumbar vertebral compression fractures who were treated with vertebroplasty or conservative therapy.^[12] Fractures were treated within two weeks of onset of pain. Details of randomization and rate of follow-up were not reported. At one day after treatment, the vertebroplasty group had a reduction in pain scores and improvement in physical function, with VAS pain scores decreasing from 4.8 (5.0 max) to 2.3 (p=0.023.) and the Oswestry Disability Index (ODI) improving from 53.6% to 31.7% (p=0.012). Sixty-five percent of patients treated with vertebroplasty had stopped all analgesics within 48 hours. The conservatively group showed no benefit in the first 48 hours, but by six weeks VAS and ODI scores were described as similar in the two groups (specific data was not reported). Evaluation of this study is limited by the incomplete reporting.

A 2016 RCT by Yang compared vertebroplasty to conservative therapy in 135 patients over 70 years of age with severe back pain due to an osteoporotic vertebral fracture after minor or mild trauma.^[13] Vertebroplasty was performed at a mean of 8.4 days after pain onset. Patients in the conservative therapy group were placed in bedrest for at least two weeks after diagnosis with analgesics, followed by bracing and assistive devices. All patients receiving vertebroplasty could stand and walk with a brace at one day posttreatment while only 12 (23.5%) patients could stand up and walk after 2 weeks of bedrest. The average duration of bedrest from pain onset was 7.8 ± 4.7 days (range, 2-15 days) in the vertebroplasty group compared to 32.5 ± 14.3 days (range, 14-60 days) in the conservative therapy group. At 1-year follow-up, there was a similar percentage of additional compression fractures, but a significantly higher complication rate in the conservative therapy group (35.3%) compared to the vertebroplasty group (16.1%; p<0.001). Complications included pneumonia, urinary tract infection, deep vein thrombosis, depression, and sleep disorders.

In 2014, Chen reported a nonblinded RCT comparing VP with conservative management.^[14] Included patients (n=89) had MRI-confirmed chronic compression fractures with persistent severe pain for 3 months or longer. At 12 months follow-up, pain scores decreased from 6.5 to 2.5 in the VP group and from 6.4 to 4.1 in the control group (p<0.001). Complete pain relief was reported by 84.8% of VP patients and 34.9% of controls. The final Oswestry Disability Index (ODI) score in the VP group and the conservative management group was 15 and 32.1, respectively (p<0.001) and the final Roland Morris Disability Score was 8.1 and 10.7, respectively (p<0.001).

In 2011, Farrokhi reported blinded RCT that compared vertebroplasty with optimal medical management in 82 patients.^[15] Patients had painful osteoporotic vertebral compression fractures that were refractory to analgesic therapy for at least four weeks and less than one year. The patients and the physicians involved in the treatment of the patients were not aware of the treatment that the other group was receiving. Control of pain and improvement in quality of life were measured by independent raters before treatment and at one week and 2, 6, 12, 24, and 36 months after the beginning of treatment. Radiologic evaluation to measure vertebral body height and correction of deformity was performed before and after treatment and after 36 months of follow-up. At one week, the mean VAS score decreased from 8.4 to 3.3 in the vertebroplasty group and from 7.2 to 6.4 in the conservative management group, with between-group differences that remained significant through 6 months of follow-up. Group differences on the Oswestry Disability Index lower back pain score were significantly lower in the vertebroplasty group throughout the 36 months of the study. New symptomatic adjacent fractures developed in one patient (2.6%) in the vertebroplasty group and six patients (15.4%) in the conservative management group. In one patient, epidural cement leakage caused severe lower-extremity pain and weakness that was treated with bilateral laminectomy and evacuation of bone cement.

VERTOS II, reported by Klazen in 2010, was an open-label randomized trial of 202 patients at 6 hospitals in the Netherlands and Belgium.^[16] Out of 431 patients who were eligible for randomization, 229 (53%) had spontaneous pain relief during assessment. Participants with at least one painful osteoporotic vertebral fracture of six weeks or less in duration were assigned to undergo vertebroplasty or conservative management (ie, bedrest, analgesia, cast, physical support). The primary outcome was pain relief of 3 points measured on a 10-point VAS at one month and one year. A sample size of 100 per group was calculated to provide sufficient power to show a 25% difference in pain relief. All analyses were performed according to ITT principles. Clinically significant pain relief was defined as 30% change on the VAS (0-10 scale).

One hundred one subjects were enrolled into the treatment group and 101 into the control arm; 81% completed 12-month follow-up. There were no significant differences in the primary outcome (pain relief of 3 points) measured at one month and one year. Vertebroplasty resulted in greater pain relief than did medical management through 12 months (<0.001); there were significant between group differences in mean VAS score at one month (2.6; 1.74 to 3.37; $p<0.001$) and at one year (2.0; 1.13 to 2.80; $p<0.001$). Survival analysis showed significant pain relief was quicker (29.7 vs 115.6 days) and was achieved in more patients after vertebroplasty than after conservative management.

Yi (2014) assessed the occurrence of new vertebral compression fractures after treatment with cement augmenting procedures (vertebroplasty or kyphoplasty) versus conservative treatment in an RCT with 290 patients (363 affected vertebrae).^[17] Surgically treated patients were discharged the next day. Patients treated conservatively (pain medication, bedrest, body brace, physical therapy) had a mean length of stay of 13.7 days. Return to usual activity occurred at one week for 87.6% of operatively treated patients and at two months for 59.2% of conservatively treated patients. All patients were evaluated with radiographs and MRI at six months and then at yearly intervals until the last follow-up session. At a mean follow-up of 49.4 months (range, 36-80 months), 10.7% of patients had experienced 42 new symptomatic vertebral compression fractures. There was no significant difference in the incidence of new vertebral fractures between the operative (18 total; 9 adjacent, 9 nonadjacent) and conservative (24 total; 5 adjacent, 16 nonadjacent, 3 same level) groups, but the mean time to

a new fracture was significantly shorter in the operative compared with nonoperative group (9.7 months vs 22.4 months).

SECTION SUMMARY

It remains unclear whether vertebroplasty provides additional beneficial effects compared to sham procedure for painful osteoporotic compression fractures. Findings from the most recent RCTs concur with earlier nonrandomized reports that vertebroplasty may be associated with significant improvements in pain and/or function. However, the lack of significant improvements in pain or function in the sham-controlled trials and in some other RCTs is suggestive that this treatment effect may not be universal. In addition, it remains unclear whether the potential benefits of the procedure outweigh harms (such as risk of additional vertebral fractures). Despite these concerns, use of VP has become increasingly widespread as a treatment of refractory vertebral fracture.

BALLOON KYPHOPLASTY

SYSTEMATIC REVIEWS

Sun (2020) performed a meta-analysis of 32 studies (n=945) in patients with osteoporotic vertebral compression fracture treated with vertebral augmentation or conservative treatment.^[18] No significant differences were observed in the risk of clinical fracture (RR, 1.22; 95% CI, 0.70 to 2.12) or radiological fracture (RR, 0.91; 95% CI, 0.71 to 2.12). Overall, 10 studies were rated as high quality, and the remainder were rated as low quality. Results remained consistent when stratified by RCTs and non-RCTs.

Hinde (2020) performed a meta-analysis of seven studies on the effect of vertebral augmentation (either vertebroplasty and/or balloon kyphoplasty) compared with nonsurgical management in over 1.5 million patients with osteoporotic vertebral compression fractures.^[19] Compared with nonsurgical management, vertebral augmentation reduced risk of mortality (HR, 0.78; 95% CI, 0.66 to 0.92). These benefits remained significant in stratified analyses of mortality over periods of 2 years (HR, 0.70; 95% CI, 0.69 to 0.71) and 5 years (HR, 0.79; 95% CI, 0.62 to 1.00).

Wang (2018) published a meta-analysis and systematic review aimed at exploring the overall safety and efficacy of balloon kyphoplasty versus percutaneous vertebroplasty for osteoporotic vertebral compression fracture (OVCF) based on qualified studies using a search of multiple databases up to January 2018.^[20] Sixteen studies were included in the meta-analysis with 647 subjects in the kyphoplasty group and 758 subjects in the vertebroplasty group. The age of the patients in both groups was over 60 years. The results indicated that kyphoplasty significantly decreased the kyphoid wedge angle (standard mean difference: 0.98; 95% CI 0.40-1.57), increased the postoperative vertebral body height (standard mean difference, -1.27; 95% CI -1.86 to -0.67), and decreased the risk of cement leakage (relative risk, 0.62; 95% CI 0.47-0.80) in comparison with vertebroplasty. However, there was no statistical difference in

visual analog scale (VAS) scores and Oswestry Disability Index (ODI) scores between the two groups. The study is limited in that there are differences in the inclusion and exclusion criteria for patients between studies. Additionally, the operating techniques in the various studies differed and the low quality of included studies and the number of included studies is limited. Lastly, pooled data were used for analysis and individual patients' data were not available which limited a more comprehensive analysis.

Xiang (2018) reviewed the literature through April 2017, evaluating the role of unilateral balloon kyphoplasty and conducted a meta-analysis to compare the efficacy and safety of unilateral and bilateral kyphoplasty in patients with osteoporotic vertebral compression fracture.^[21] The meta-analysis included nine studies, six randomized controlled trials and three retrospective comparative studies, on the use of unipedicular balloon kyphoplasty in the treatment of 870 patients with OVCFs. The patients were followed for periods ranging from 2 weeks to 42.2 months with a mean age of 68.85 years. After unilateral balloon kyphoplasty, the mean postoperative visual analog score (VAS) ranged from 1.74 to 4.77, mean postoperative kyphotic angle ranged from 5.9° to 11.22°, and complications involving cement leaks ranged from 6.8 to 21.9% or adjacent level fractures was from 0 to 5.6%). Unilateral kyphoplasty had significantly shorter operative time, and less bone-cement volume; however, the postoperative VAS, Oswestry Disability Index (ODI), vertebral height restoration rate, and cement leakage and adjacent vertebral fracture rate were similar to bilateral kyphoplasty. The sample size (six RCTs and three retrospective comparative studies) limited the level of evidence for the analysis. Heterogeneity was also detected among the studies once they were pooled. In addition, incomplete data recording was discovered once clinical outcomes were extracted.

In a Bayesian network meta-analysis, Zhao (2017) examined the efficacy and safety of vertebroplasty, kyphoplasty, and conservative treatment (CT) for the treatment of osteoporotic vertebral compression fractures.^[22] Sixteen RCTs were identified (total N=2046 participants; vertebroplasty, 816; kyphoplasty, 478; CT = 752). Eleven of the RCTs compared vertebroplasty with CT; two RCTs compared kyphoplasty with CT; and three RCTs compared kyphoplasty with vertebroplasty. Each trial assessed at least one of the following: VAS, the RMDQ, the European Quality of Life-5 Dimensions (EQ-5D), and the observance of any new fractures. Network meta-analysis demonstrated that kyphoplasty was superior to conservative therapy as assessed by VAS (mean difference [MD], 0.94; 95% CI, -0.40 to 2.39), EQ-5D (MD -0.10; 95% CI, -0.17 to -0.01), and RMDQ (MD=5.72; 95% CI, 1.05 to 10.60). Insufficient data were present to complete pairwise comparison of kyphoplasty with CT for some metrics. No significant differences were found between vertebroplasty and kyphoplasty for pain relief, daily function, and quality of life. Kyphoplasty was associated with the lowest risk of new fractures, while vertebroplasty was the most effective treatment for pain relief. This review was limited by significant heterogeneity across measured outcomes and length of follow-up in studies; the presence of performing and reporting bias in studies was also a concern.

Huang (2014) published a systematic review with meta-analysis evaluating whether unilateral or bilateral KP was superior for effectiveness and safety.^[23] No large studies were found that directly compared these two approaches. Five studies with 253 patients met inclusion criteria. No clinically significant difference between unilateral and bilateral KP was found for improvement in pain and function, kyphosis angle reduction, and anterior vertebral height restoration. The rate of adverse events was also similar between the two approaches. However, the quality of the evidence was graded as very low and the authors recommended high-quality RCTs be conducted to resolve this question.

RANDOMIZED CONTROLLED TRIALS

In 2009, Wardlaw reported on the findings of the FREE trial, a nonblinded industry-sponsored multisite RCT in which 300 adult participants with 1 to 3 painful osteoporotic vertebral compression fractures (VCF) of less than 3 months in duration were assigned to undergo kyphoplasty or conservative care.^[24] Twenty-four-month results of this study were reported by Boonen in 2011 and by Van Meirhaeghe in 2013.^[25, 26] Scores for the primary outcome, 1-

month change in 36-Item Short-Form Health Survey (SF-36) Physical Component Summary (PCS) score, were significantly higher for those in the kyphoplasty group. The difference between the 2 groups was 5.2 points (95% confidence interval, 2.9 to 7.4; $p < 0.001$). Kyphoplasty was associated with greater improvements in SF-36 PCS scores at 6-month follow-up (3.39 points), but not at 12 or 24 months. Greater improvement in back pain was observed over 24 months for kyphoplasty (-1.49 points) and remained statistically significant at 24 months. Participants in the kyphoplasty group also reported greater improvements in quality of life and Roland-Morris Disability Questionnaire (RMDQ) score at short-term follow-up. At 12 months, fewer kyphoplasty patients (26.4% vs 42.1%) had received physical therapy or walking aids, back braces, wheelchairs, miscellaneous aids, or other therapy. Fewer kyphoplasty patients used opioid medications through 6 months (29.8% vs 42.9%) and fewer pain medications through 12 months (51.7% vs. 68.3%). Other differences between the groups were no longer apparent at 12 months; possibly due to natural healing of fractures.

In 2011, Berenson published results from an international randomized multicenter clinical trial.^[27] They enrolled 134 patients with cancer who were at least 21 years of age. Participants had at least one and not more than 3 painful vertebral compression fractures (VCF). (These appear to be due to osteoporosis, rather than from a metastatic lesion.) The primary outcome was change in functional status from baseline at 1 month as measured by the Roland Morris Disability Questionnaire (RMDQ). Treatment allocation was not blinded, and the primary outcome at 1 month was analyzed using all participants with data both at baseline and at 1 month. Crossover to the balloon kyphoplasty arm was allowed after 1 month. The authors report baseline scores in the kyphoplasty and nonsurgical groups of 17.6 and 18.2, respectively and 9.10 and 18.0 at 1-month follow-up. P-value for the between group difference in scores $p = 0.0001$. However, conclusions based upon this trial are limited due to lack of blinding and short-term (one month) follow-up.

In 2011, Edidin reported mortality risk in Medicare patients who had VCFs and had been treated with vertebroplasty, kyphoplasty, or nonoperatively.^[28] This study was industry-funded. Using the U.S. Medicare dataset, they identified 858,978 patients who had VCFs between 2005 and 2008. The data set included 119,253 kyphoplasty patients and 63,693 vertebroplasty patients. Survival was calculated from the index diagnosis date until death or the end of follow-up (up to 4 years). Cox regression was used to evaluate the joint effect of multiple covariates, which included sex, age, race/ethnicity, patient health status, type of diagnosed fracture, site of service, physician specialty, socioeconomic status, year of diagnosis, and census region. After adjusting for covariates, patients in the operated cohort (vertebroplasty or kyphoplasty) were found to have a higher adjusted survival rate (60.8%) than patients in the nonoperated cohort (50.0%) and were 37% less likely to die. The adjusted survival rates for vertebroplasty or kyphoplasty were 57.3% and 62.8%, respectively, a 23% lower relative risk for kyphoplasty. As noted by the authors, a causal relationship cannot be determined from this study.

SECTION SUMMARY

Two moderate-sized unblinded RCTs compared kyphoplasty to conservative care and found short-term benefits in pain and other outcomes. Other RCTs, as summarized in a meta-analysis, reported that outcomes for kyphoplasty and vertebroplasty are similar. Two randomized trials that compared mechanical vertebral augmentation (Kiva) to kyphoplasty reported similar outcomes for the two procedures. A major limitation of the RCTs is the lack of a comparator sham procedure. Due to the possible placebo effect observed in the recent trials of vertebroplasty, the validity of results from non-sham-controlled trials are questionable. There

are no RCTs of kyphoplasty for vertebral body metastasis. However, as with vertebroplasty, the use of kyphoplasty has become increasingly widespread as a treatment of refractory vertebral fracture.

MECHANICAL VERTEBRAL AUGMENTATION WITH KIVA® OR SPINEJACK

SYSTEMATIC REVIEWS

Mattie (2021) conducted a systematic review and meta-analysis of seven RCTs (N=476) that compared the magnitude and duration of pain relief with vertebral augmentation (ie, balloon kyphoplasty or percutaneous vertebroplasty), with or without additional therapy, to any other intervention or placebo/sham for the treatment of cancer-related vertebral compression fractures.^[29] In five of the seven studies, vertebral augmentation alone comprised one group; comparative treatments included nonsurgical management, Kiva implantation, and combinations of percutaneous vertebroplasty and radiofrequency therapy, chemotherapy, intrasomatic steroid injection, or 125I seeds. Results revealed an overall positive and statistically significant effect of vertebral augmentation for the management of cancer-related vertebral compression fractures. This effect was particularly pronounced when comparing vertebral augmentation to nonsurgical management, radiofrequency ablation, or chemotherapy alone. The authors noted that there was much heterogeneity among the included studies regarding the treatment methods in the control groups and one study allowed patients to crossover to the intervention group, potentially leading to biased results.

RANDOMIZED CONTROLLED TRIALS

Noriega (2019) reported the pivotal multicenter non-inferiority trial of the SpineJack vertebral augmentation system. Patients (n=152) with osteoporotic vertebral compression fractures (OVCF) less than 3 months old were randomized to treatment with SpineJack or balloon kyphoplasty. The primary outcome was a composite measure that included improvement in VAS for pain of greater than 20 mm, maintenance or improvement in ODI, and lack of adverse events. Vertebral height was prespecified to be included if the primary outcome was achieved. Non-inferiority was achieved with 89.8% of SpineJack patients achieving the composite of clinical success compared to 87.3% for balloon kyphoplasty. When including the restoration of vertebral body height, the SpineJack procedure was found to be superior to balloon kyphoplasty at 6 months (88.1% vs. 60.9%) and at 12 months (79.7% vs. 59.3%). There was also a reduction in adjacent vertebral fractures with the mechanical augmentation system (12.9% vs. 27.3%). Interpretation of this study is limited by the lack of sham control group.

Tutton (2015) published results from an RCT that where vertebral augmentation with the Kiva® VCF System® was compared with balloon kyphoplasty in the KAST trial, a 2015 pivotal non-inferiority RCT.^[30] This industry-sponsored multicenter open-label trial was conducted in 300 patients with one or two osteoporotic vertebral compression fractures. The Kiva group included 153 patients and the KP group included 147 patients. Included were patients with VAS for back pain of at least 70 mm out of 100 after two to six weeks of conservative care or a VAS of at least 50 mm after six weeks of conservative care, and an ODI of at least 30%. The primary endpoint at 12 months was a composite of a reduction in fracture pain by at least 15 mm on VAS, maintenance or improvement in function on ODI, and absence of device-related serious adverse events. The primary endpoint was met for 94.5% of patients treated with Kiva® and 97.6% of patients treated with kyphoplasty (Bayesian posterior probability of 99.92% for non-inferiority, using as-treated analysis). In the 285 treated patients, Kiva® resulted in a mean improvement of 70.8 points in VAS, compared with a 71.8 point improvement for kyphoplasty.

There was a 38.1 point improvement in ODI for the Kiva® group, compared to a 42.2 point improvement for the kyphoplasty group. There were no device-related serious adverse events. The total volume of cement was 50% less with Kiva® and there was lower cement leakage compared with kyphoplasty (16.9% vs 25.8%, respectively).

In 2013, Korovessis reported a randomized trial comparing mechanical vertebral augmentation with the Kiva device versus balloon kyphoplasty in 180 patients with osteoporotic vertebral body fractures.^[31] Mean follow-up was 14 months (range 13-15 months). The groups showed similar improvements in VAS for back pain, SF-36, and ODI. For example, there was a greater than 5.5 point improvement in VAS in 54% of patients in the Kiva group and 43% of patients in the balloon kyphoplasty group. Radiological measures of vertebral height were similar in the two groups. Kiva reduced the Gardner kyphotic angle, while residual kyphosis of more than five degrees was more frequently observed in the balloon kyphoplasty group. Patients and outcome assessors were reported to be unaware of the group assignment, although it is not clear if the Kiva device was apparent in the radiographs. Cement leakage into the canal occurred in two patients treated with balloon kyphoplasty, necessitating decompression, compared with none following the Kiva procedure.

SECTION SUMMARY

The Kiva procedure appears to be at least equivalent to KP in pain reduction and improved functional outcomes with a lower rate of cement leakage. Mechanical vertebral augmentation with SpineJack was found to be non-inferior to balloon kyphoplasty for success on a composite outcome measure and superior to balloon kyphoplasty when vertebral height restoration was included in the composite.

RADIOFREQUENCY-ASSISTED MECHANICAL VERTEBRAL AUGMENTATION

Current published evidence is limited to a few very small, short-term feasibility studies. These preliminary studies do not permit conclusions due to methodological limitations, including but not limited to the lack of randomized comparison to alternative treatments, small study populations, and the lack of long-term follow-up. One meta-analysis by Feng (2017) compared radiofrequency kyphoplasty with balloon kyphoplasty, however the conclusions are limited by the same methodological parameters outlined above.^[32]

VERTEBRAL BODY STENTING

SYSTEMATIC REVIEW

In 2015, Martin-Lopez published results from a systematic review that examined the effectiveness and safety of stentoplasty in patients with osteoporotic vertebral body fractures.^[33] Five studies were included in the review, two clinical trials and three observational studies. The authors found there was no difference between the two procedures in terms of reduction of kyphosis, time of exposure to radiation or postoperative loss of cement. Although stentoplasty in comparison to vertebroplasty showed an improvement of restoration of vertebral height ($P=0.042$), kyphosis correction and volume of bone cement, no differences were found between two procedures in terms of loss of vertebral body volume. Based on observational studies, stentoplasty improved vertebral height, pain and functional disability at six and 12 months follow-up, and corrected the angle vertebral fractures in patients with

osteoporotic vertebral body. The authors concluded that there was no advantage of stentoplasty over balloon kyphoplasty.

SACROPLASTY AND COCCYGEOPLASTY

Sacroplasty is an evolving technique with numerous methods (short axis, long axis, balloon-assisted short axis, and iliosacral screws). No randomized trials of sacroplasty have been reported. The evidence on sacroplasty is limited to several small case reports or series.^[34-53] These initial pilot studies reported rapid pain relief with few complications. Due to the small size of the evidence base, harms associated with sacroplasty have not been adequately studied. There are complications of cement leakage with sacroplasty that are not observed with vertebroplasty. Leakage of PMMA into the presacral space, spinal canal, sacral foramen, or sacroiliac joint may result in pelvic injection of PMMA, sacral nerve root or sacral spinal canal compromise, or sacroiliac joint dysfunction.^[54] Coccygeoplasty has been reported, but no adequate clinical trial data has been published.

ADVERSE EVENTS

The most commonly reported adverse events for vertebroplasty or balloon kyphoplasty are new compression fractures and cement leakage. Cement leakage remains a concern, though it has been shown to be reduced in kyphoplasty relative to vertebroplasty. Most incidents of cement leakage were reported to be asymptomatic. In addition, there are case reports of cardiac perforation, cardiac tamponade, and embolism of cement into pulmonary vessels.

Zhang (2017) published a meta-analysis of comparative studies to evaluate the incidence of new vertebral fractures between vertebral augmentation, such as vertebroplasty and kyphoplasty, and no operation.^[55] Twelve comparative studies were identified with a total of 1,328 patients (768 underwent operation and 560 did not undergo operation). There were no significant differences between groups for new nor existing vertebral fractures. In addition, there were no significant differences in bone mineral density (lumbar and femoral neck regions).

Zhan (2016) evaluated risk factors for cement leakage after vertebroplasty or kyphoplasty. Zhan conducted a systematic review with meta-analysis that included 32 studies with 2,872 patients.^[56] Cement leakage incidence were 54.7% for vertebroplasty and 18.4% for kyphoplasty. The authors found that the significant risk factors were intravertebral cleft (OR=1.40; 95% CI, 1.09-1.78), cortical disruption (OR=5.56; 95% CI, 1.84-16.81), cement viscosity (OR=3.32; 95% CI, 1.36-8.07) and injected cement volume (WMD=0.59; 95% CI, 0.02-1.17). Other factors were not associated with significant risk including age, sex, and fracture type. The authors also concluded that more randomized controlled trials are needed to validate these findings.

Xiao (2015) published results from a systematic review and meta-analysis that compared complications between percutaneous vertebroplasty (PVP) and balloon kyphoplasty (BK) for the treatment of osteoporotic vertebral compression fractures.^[57] Nineteen studies encompassing 1,787 patients (887 PVP, and 900 BKP) were included in the analysis. The authors reported that the two procedures suffer from equal risk of subsequent spinal fractures; however, PVP had a higher cement leakage rate when compare to BK.

Zhang conducted a systematic review with meta-analysis of four RCTs^[5, 16, 58, 59] (N=454) published through April 2013.^[60] The aim was to analyze the causal relationship between

vertebroplasty and new-onset vertebral fractures in osteoporotic vertebral fracture patients. Authors noted several limitations to the meta-analysis. The authors reported that the RCTs did not support a conclusion that VP significantly increased the occurrence of new postoperative or adjacent vertebral fractures. Due to inclusion of only four RCTs, some with modest sample sizes, could allow over- or under-estimation of results. In addition, there was heterogeneity between RCTs in fracture age, intervention, duration of follow-up, and study design. Attrition bias was also of concern due to moderately high drop-out rates.

Yi (2014) evaluated the occurrence of new vertebral compression fractures after treatment with cement augmenting procedures (vertebroplasty or kyphoplasty) versus conservative treatment in an RCT with 290 patients (363 affected vertebrae).^[17] Patients treated conservatively (offered pain medication, bed rest, a body brace, and physiotherapy) had a mean length of stay of 13.7 days. Return to usual activity occurred at one week for 87.6% of operatively-treated patients and at two months for 59.2% of conservatively-treated patients. At a mean follow-up of 49.4 months (range, 36-80), 10.7% of patients had experienced 42 new symptomatic vertebral compression fractures. There was no significant difference in the incidence of new vertebral fractures between the operative (18 total, 9 adjacent and 9 nonadjacent) and conservative (24 total, 5 adjacent, 16 nonadjacent, and 3 same level) groups, but the mean time to a new fracture was significantly shorter in the operative compared to nonoperative group (9.7 vs 22.4 months).

In 2014 a study was published using the SWISSspine registry (SSR) which included 375 single-level osteoporotic vertebral fracture patients followed for a mean follow-up of 3.6 months following KP.^[61] Post-KP adjacent segment fractures were found in 37 (10%) patients, occurring on average at 2.8 months postoperatively. Significant risk factors included preoperative segmental kyphosis >30 degrees ($p=0.026$), rheumatoid arthritis ($p=0.038$), and cardiovascular disease ($p=0.047$). Patients with postoperative adjacent segment fractures had significantly higher back pain at final follow-up.

In a retrospective analysis of 171 post-KP patients at a mean follow-up of 41 months, Civelek also found higher preoperative kyphotic angle to be a risk factor significantly associated with adjacent level fractures.^[62] Female sex was also found to be a significant risk factor. The severity of osteoporosis was not a determining factor.

PRACTICE GUIDELINE SUMMARY

AMERICAN ACADEMY OF ORTHOPAEDIC SURGEONS

In 2010, the American Academy of Orthopaedic Surgeons (AAOS) Board of Directors approved a clinical practice guideline on the treatment of osteoporotic spinal compression fractures.^[63] The Board approved a strong recommendation against the use of vertebroplasty for patients who “present with an osteoporotic spinal compression fracture on imaging with correlating clinical signs and symptoms and who are neurologically ‘intact’.” This recommendation was based on five RCTs^[5, 6, 64-66], two of which were graded Level I (defined as reliable), and three of which were graded Level II (defined as moderately reliable). In coming out with a strong recommendation, the committee expressed their confidence that future evidence is unlikely to overturn the existing evidence. The Board also downgraded the recommendation supporting the use of kyphoplasty from “moderate” to “limited” based upon low quality and inconclusive evidence comparing this procedure with conservative care and vertebroplasty, respectively.

AMERICAN SOCIETY OF ANESTHESIOLOGISTS AND THE AMERICAN SOCIETY OF REGIONAL ANESTHESIA AND PAIN MEDICINE

Practice guidelines from the American Society of Anesthesiologist (ASA) and the American Society of Regional Anesthesia and Pain Medicine (ASRA) support the use of “minimally invasive spinal procedures” (including vertebroplasty and vertebral augmentation), stating: “Consultants, ASA members, and ASRA members strongly agree that minimally invasive spinal procedures should be performed for pain related to vertebral compression fractures.”^[67] The practice guidelines go on to make the specific recommendation in favor of these procedures in “treatment of pain related to vertebral compression fractures” despite a review of the literature which found that available randomized sham-controlled trials had either not found differences associated with treatment groups, or that differences were inconsistent across available studies.

AMERICAN COLLEGE OF RADIOLOGY, AMERICAN SOCIETY OF NEURORADIOLOGY, AMERICAN SOCIETY OF SPINE RADIOLOGY, THE SOCIETY OF INTERVENTIONAL RADIOLOGY, AND THE SOCIETY OF NEUROINTERVENTIONAL SURGERY

The American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American Society of Spine Radiology (ASSR), the Society of Interventional Radiology (SIR), and the Society of NeuroInterventional Surgery (SNIS) along with the American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CAN), and the Canadian Interventional Radiology Association (CIRA) published a joint position statement on percutaneous vertebral augmentation in 2014^[68], which was revised in 2017^[69]. This document states that percutaneous vertebral augmentation using vertebroplasty or kyphoplasty and performed in a manner in accordance with public standards is a safe, efficacious, and durable procedure in appropriate patients with symptomatic osteoporotic and neoplastic fractures. The document also states that these procedures are offered only when nonoperative medical therapy has not provided adequate pain relief or pain is significantly altering the patient’s quality of life.

An updated 2012 joint practice guideline addresses the performance of vertebral augmentation in general and refers to all available percutaneous techniques used to achieve internal vertebral body stabilization, including vertebroplasty, balloon kyphoplasty, radiofrequency ablation and coblation, mechanical void creation, and injection of bone graft material or bone substitutes.^[69] The ACR, ASN, ASSR, SIR, and SNIS consider vertebral augmentation to be an established and safe procedure, and provide guidelines for patient selection, qualifications and responsibilities of personnel, specifications of the procedure, equipment quality control, and quality improvement and documentation.

SOCIETY OF NEUROINTERVENTIONAL SURGERY

In 2014 the Society of NeuroInterventional Surgery (SNIS) published a report that included a systematic review with meta-analysis and the following recommendations:^[70]

1. Kyphoplasty in selected patients is superior to conservative medical therapy in reducing back pain, disability and improving Karnofsky performance status and quality of life for patients with cancer
2. and disabling back pain from a vertebral fracture (AHA Class IIA, Level of Evidence B).

3. Vertebroplasty and kyphoplasty are reasonable therapeutic options in selected patients with cancer and severe back pain from a vertebral fracture that is refractory to conservative medical therapy (AHA Class IIA, Level of Evidence B).
4. Vertebroplasty and kyphoplasty are reasonable therapeutic options in selected patients with severe back pain from an osteoporotic vertebral fracture that is refractory to conservative medical therapy (AHA Class IIA, Level of Evidence B).

SOCIETY OF INTERVENTIONAL RADIOLOGY

In a 2014 quality improvement guideline on percutaneous vertebroplasty from SIR, vertebral augmentation was recommended for compression fractures refractory to medical therapy. Failure of medical therapy includes the following situations:^[71]

1. Patients who are rendered nonambulatory because of pain from a weakened or fractured vertebral body, pain persisting at a level that prevents ambulation despite 24 hours of analgesic therapy;
2. Patients with sufficient pain from a weakened or fractured vertebral body that physical therapy is intolerable, pain persisting at that level despite 24 hours of analgesic therapy; or
3. Patient with a weakened or fractured vertebral body, and unacceptable side effects such as excessive sedation, confusion, or constipation because of the analgesic therapy necessary to reduce pain to a tolerable level.

AMERICAN COLLEGE OF RADIOLOGY

The American College of Radiology (ACR) published updated appropriateness criteria on the management of vertebral compression fractures in 2014.^[72] While generally supportive of vertebroplasty and kyphoplasty in specified conditions, the guidelines state the following:

- Conservative management is the traditional first-line management for osteoporotic compression fractures.
- Controversy exists over the use of vertebral augmentation due to two previous independent level 1 trials that demonstrated no clinical validity for VP over the sham control groups. Conclusions from these studies have divided the medical community with respect to the efficacy of vertebral augmentation.
- Despite this controversy, increased use of vertebral augmentation for managing painful osteoporotic and malignant vertebral fractures has been the trend, with the literature favoring patient outcomes over conservative medical management up to 1 year.
- If VP is recommended for osteoporosis or malignant fractures, it should be used for patients who have failed or cannot tolerate conservative or traditional management.
- Kyphoplasty data are less extensive but have shown similar results to VP for uncomplicated vertebral compression fractures.
- Kyphoplasty may have an advantage over traditional VP in complex cases (e.g., burst fractures with neurological compromise) or fractures in which height restoration or deformity correction may be beneficial.
- This slight mechanical advantage over VP may also affect long-term outcomes.
- More level one studies are needed to determine the medical and societal cost of the palliative effect on pain related morbidity associated with osteoporotic vertebral compression fractures. Smaller sample studies and use trends indicate vertebral augmentation has benefits over conservative medical management for the first year.

NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE

The National Institute for Health and Care Excellence (NICE) issued a 2013 technology appraisal guidance TA279, which stated that percutaneous vertebroplasty and percutaneous balloon kyphoplasty are recommended as treatment options for treating osteoporotic vertebral compression fractures in persons having severe, ongoing pain after a recent unhealed vertebral fracture, despite optimal pain management, and whose pain has been confirmed through physical exam and imaging to be at the level of the fracture. This appraisal does not address balloon kyphoplasty with stenting, because the manufacturer of the stenting system (Synthes) states there is limited evidence for vertebral body stenting as it has only recently become available.^[73]

In 2008, NICE issued CG75 on the diagnosis and management of adults with metastatic spinal cord compression. This guidance was last reviewed in 2014, and placed on the static list (no major ongoing studies identified and the next review will occur in 5 years). The guideline states that vertebroplasty or kyphoplasty should be considered for patients who have vertebral metastases, and no evidence of spinal cord compression or spinal instability, if they have mechanical pain resistant to conventional pain management and vertebral body collapse. Surgery should only be performed when all appropriate specialists, including the oncologist, interventional radiologist, and spinal surgeon agree. At present, there are relatively few patients in England receiving surgery; however, there is evidence to suggest that in a selected subset of patients, early surgery may be more effective at maintaining mobility than radiotherapy.^[74]

AMERICAN SOCIETY OF PAIN AND NEUROSCIENCE

In 2021, the American Society of Pain and Neuroscience (ASPN) published practice guidelines for the interventional management of cancer-associated pain.^[75] The guideline included a best practice statement that stated "vertebral augmentation should be strongly considered for patients with symptomatic vertebral compression fractures from spinal metastases (evidence level 1-A)." However, ASPN noted that there is little data to suggest the superiority of either vertebroplasty or kyphoplasty when treating malignant vertebral compression fractures.

SUMMARY

There is enough research to show that percutaneous vertebroplasty, kyphoplasty, SpineJack, or Kiva for the treatment of vertebral fractures improves health outcomes including but not limited to pain and/or function in select patients. In addition, practice guidelines recommend these procedures for vertebral fractures. Therefore, vertebroplasty, kyphoplasty, SpineJack, or Kiva may be considered medically necessary in select patients with vertebral fractures when policy Criteria are met.

There is not enough research to show that percutaneous vertebroplasty, kyphoplasty, SpineJack, or Kiva for all other indications improves health outcomes including mechanical vertebral augmentation using techniques other than balloon kyphoplasty. In addition, there are no practice guidelines that recommend these techniques. Therefore, percutaneous vertebroplasty, kyphoplasty, SpineJack, or Kiva for all other indications is considered investigational including but not limited to percutaneous mechanical vertebral augmentation

techniques using devices other than balloon devices (e.g., radiofrequency-assisted vertebral augmentation or vertebral body stents).

There is not enough research to show that sacroplasty or coccygeoplasty improves health outcomes. In addition, no practice guidelines recommend these procedures for any indication. Therefore, sacroplasty and coccygeoplasty are considered investigational.

REFERENCES

1. Zhu HT, Ding DG, Wang S, et al. Comparison Between Percutaneous Kyphoplasty and Percutaneous Vertebroplasty in Terms of Efficacy in Osteoporotic Vertebral Compression Fractures: A Meta-analysis. *Altern Ther Health Med*. 2022;28(5):49-53. PMID: 35648693
2. Halvachizadeh S, Stalder AL, Bellut D, et al. Systematic Review and Meta-Analysis of 3 Treatment Arms for Vertebral Compression Fractures: A Comparison of Improvement in Pain, Adjacent-Level Fractures, and Quality of Life Between Vertebroplasty, Kyphoplasty, and Nonoperative Management. *JBJs Rev*. 2021;9(10). PMID: 34695056
3. Xie L, Zhao ZG, Zhang SJ, et al. Percutaneous vertebroplasty versus conservative treatment for osteoporotic vertebral compression fractures: An updated meta-analysis of prospective randomized controlled trials. *Int J Surg*. 2017;47:25-32. PMID: 28939236
4. Buchbinder R, Golmohammadi K, Johnston RV, et al. Percutaneous vertebroplasty for osteoporotic vertebral compression fracture. *The Cochrane database of systematic reviews*. 2015;4:CD006349. PMID: 25923524
5. Buchbinder R, Osborne RH, Ebeling PR, et al. A randomized trial of vertebroplasty for painful osteoporotic vertebral fractures. *N Engl J Med*. 2009;361(6):557-68. PMID: 19657121
6. Kallmes DF, Comstock BA, Heagerty PJ, et al. A randomized trial of vertebroplasty for osteoporotic spinal fractures. *N Engl J Med*. 2009;361(6):569-79. PMID: 19657122
7. Staples MP, Kallmes DF, Comstock BA, et al. Effectiveness of vertebroplasty using individual patient data from two randomised placebo controlled trials: meta-analysis. *BMJ*. 2011;343:d3952. PMID: 21750078
8. Chew C, Craig L, Edwards R, et al. Safety and efficacy of percutaneous vertebroplasty in malignancy: a systematic review. *Clin Radiol*. 2011;66(1):63-72. PMID: 21147301
9. Firanescu CE, de Vries J, Lodder P, et al. Vertebroplasty versus sham procedure for painful acute osteoporotic vertebral compression fractures (VERTOS IV): randomised sham controlled clinical trial. *BMJ*. 2018;361:k1551. PMID: 29743284
10. Clark W, Bird P, Gonski P, et al. Safety and efficacy of vertebroplasty for acute painful osteoporotic fractures (VAPOUR): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet*. 2016;388(10052):1408-16. PMID: 27544377
11. Kroon F, Staples M, Ebeling PR, et al. Two-year results of a randomized placebo-controlled trial of vertebroplasty for acute osteoporotic vertebral fractures. *J Bone Miner Res*. 2014;29(6):1346-55. PMID: 24967454
12. Leali PT, Solla F, Maestretti G, et al. Safety and efficacy of vertebroplasty in the treatment of osteoporotic vertebral compression fractures: a prospective multicenter international randomized controlled study. *Clin Cases Miner Bone Metab*. 2016;13(3):234-36. PMID: 28228788
13. Yang EZ, Xu JG, Huang GZ, et al. Percutaneous Vertebroplasty Versus Conservative Treatment in Aged Patients With Acute Osteoporotic Vertebral Compression Fractures:

- A Prospective Randomized Controlled Clinical Study. *Spine (Phila Pa 1976)*. 2016;41(8):653-60. PMID: 26630417
14. Chen D, An ZQ, Song S, et al. Percutaneous vertebroplasty compared with conservative treatment in patients with chronic painful osteoporotic spinal fractures. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2014;21(3):473-7. PMID: 24315046
 15. Farrokhi MR, Alibai E, Maghami Z. Randomized controlled trial of percutaneous vertebroplasty versus optimal medical management for the relief of pain and disability in acute osteoporotic vertebral compression fractures. *J Neurosurg Spine*. 2011;14(5):561-9. PMID: 21375382
 16. Klazen CA, Lohle PN, de Vries J, et al. Vertebroplasty versus conservative treatment in acute osteoporotic vertebral compression fractures (Vertos II): an open-label randomised trial. *Lancet*. 2010;376(9746):1085-92. PMID: 20701962
 17. Yi X, Lu H, Tian F, et al. Recompression in new levels after percutaneous vertebroplasty and kyphoplasty compared with conservative treatment. *Archives of orthopaedic and trauma surgery*. 2014;134(1):21-30. PMID: 24287674
 18. Sun HB, Jing XS, Tang H, et al. Clinical and radiological subsequent fractures after vertebral augmentation for treating osteoporotic vertebral compression fractures: a meta-analysis. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2020;29(10):2576-90. PMID: 32776263
 19. Hinde K, Maingard J, Hirsch JA, et al. Mortality Outcomes of Vertebral Augmentation (Vertebroplasty and/or Balloon Kyphoplasty) for Osteoporotic Vertebral Compression Fractures: A Systematic Review and Meta-Analysis. *Radiology*. 2020;295(1):96-103. PMID: 32068503
 20. Wang B, Zhao CP, Song LX, et al. Balloon kyphoplasty versus percutaneous vertebroplasty for osteoporotic vertebral compression fracture: a meta-analysis and systematic review. *J Orthop Surg Res*. 2018;13(1):264. PMID: 30348192
 21. Xiang GH, Tong MJ, Lou C, et al. The Role of Unilateral Balloon Kyphoplasty for the Treatment of Patients with OVCFS: A Systematic Review and Meta-Analysis. *Pain physician*. 2018;21(3):209-18. PMID: 29871365
 22. Frey ME, Warner C, Thomas SM, et al. Sacroplasty: A Ten-Year Analysis of Prospective Patients Treated with Percutaneous Sacroplasty: Literature Review and Technical Considerations. *Pain physician*. 2017;20(7):E1063-E72. PMID: 29149151
 23. Huang Z, Wan S, Ning L, et al. Is unilateral kyphoplasty as effective and safe as bilateral kyphoplasties for osteoporotic vertebral compression fractures? A meta-analysis. *Clinical orthopaedics and related research*. 2014;472(9):2833-42. PMID: 24964889
 24. Wardlaw D, Cummings SR, Van Meirhaeghe J, et al. Efficacy and safety of balloon kyphoplasty compared with non-surgical care for vertebral compression fracture (FREE): a randomised controlled trial. *Lancet*. 2009;373(9668):1016-24. PMID: 19246088
 25. Boonen S, Van Meirhaeghe J, Bastian L, et al. Balloon kyphoplasty for the treatment of acute vertebral compression fractures: 2-year results from a randomized trial. *J Bone Miner Res*. 2011;26(7):1627-37. PMID: 21337428
 26. Van Meirhaeghe J, Bastian L, Boonen S, et al. A Randomized Trial of Balloon Kyphoplasty and Non-Surgical Management for Treating Acute Vertebral Compression Fractures: Vertebral Body Kyphosis Correction and Surgical Parameters. *Spine (Phila Pa 1976)*. 2013. PMID: 23446769

27. Berenson J, Pflugmacher R, Jarzem P, et al. Balloon kyphoplasty versus non-surgical fracture management for treatment of painful vertebral body compression fractures in patients with cancer: a multicentre, randomised controlled trial. *Lancet Oncol.* 2011;12(3):225-35. PMID: 21333599
28. Edidin AA, Ong KL, Lau E, et al. Mortality risk for operated and nonoperated vertebral fracture patients in the medicare population. *J Bone Miner Res.* 2011;26(7):1617-26. PMID: 21308780
29. Mattie R, Brar N, Tram JT, et al. Vertebral Augmentation of Cancer-Related Spinal Compression Fractures: A Systematic Review and Meta-Analysis. *Spine (Phila Pa 1976).* 2021;46(24):1729-37. PMID: 33958537
30. Tutton SM, Pflugmacher R, Davidian M, et al. KAST Study: The Kiva(R) System as a Vertebral Augmentation Treatment - A Safety and Effectiveness Trial: A Randomized, Non-inferiority Trial Comparing the Kiva(R) System to Balloon Kyphoplasty in Treatment of Osteoporotic Vertebral Compression Fractures. *Spine (Phila Pa 1976).* 2015. PMID: 25822543
31. Korovessis P, Vardakastanis K, Repantis T, et al. Balloon kyphoplasty versus KIVA vertebral augmentation--comparison of 2 techniques for osteoporotic vertebral body fractures: a prospective randomized study. *Spine (Phila Pa 1976).* 2013;38(4):292-9. PMID: 23407406
32. McConnell CT, Jr., Wippold FJ, 2nd, Ray CE, Jr., et al. ACR appropriateness criteria management of vertebral compression fractures. *Journal of the American College of Radiology : JACR.* 2014;11(8):757-63. PMID: 24935074
33. Martin-Lopez JE, Pavon-Gomez MJ, Romero-Tabares A, et al. Stentoplasty effectiveness and safety for the treatment of osteoporotic vertebral fractures: a systematic review. *Orthopaedics & traumatology, surgery & research : OTSR.* 2015;101(5):627-32. PMID: 26194207
34. Frey ME, DePalma MJ, Cifu DX, et al. Efficacy and safety of percutaneous sacroplasty for painful osteoporotic sacral insufficiency fractures: a prospective, multicenter trial. *Spine (Phila Pa 1976).* 2007;32(15):1635-40. PMID: 17621211
35. Frey ME, Depalma MJ, Cifu DX, et al. Percutaneous sacroplasty for osteoporotic sacral insufficiency fractures: a prospective, multicenter, observational pilot study. *Spine J.* 2008;8(2):367-73. PMID: 17981097
36. Garant M. Sacroplasty: a new treatment for sacral insufficiency fracture. *J Vasc Interv Radiol.* 2002;13(12):1265-7. PMID: 12471192
37. Layton KF, Thielen KR, Wald JT. Percutaneous sacroplasty using CT fluoroscopy. *AJNR Am J Neuroradiol.* 2006;27(2):356-8. PMID: 16484410
38. Sciubba DM, Wolinsky JP, Than KD, et al. CT fluoroscopically guided percutaneous placement of transiliosacral rod for sacral insufficiency fracture: case report and technique. *AJNR Am J Neuroradiol.* 2007;28(8):1451-4. PMID: 17846189
39. Smith DK, Dix JE. Percutaneous sacroplasty: long-axis injection technique. *AJR Am J Roentgenol.* 2006;186(5):1252-5. PMID: 16632714
40. Tjardes T, Paffrath T, Baethis H, et al. Computer assisted percutaneous placement of augmented iliosacral screws: a reasonable alternative to sacroplasty. *Spine (Phila Pa 1976).* 2008;33(13):1497-500. PMID: 18520946
41. Binaghi S, Guntern D, Schnyder P, et al. A new, easy, fast, and safe method for CT-guided sacroplasty. *Eur Radiol.* 2006;16(12):2875-8. PMID: 17058081
42. Brook AL, Mirsky DM, Bello JA. Computerized tomography guided sacroplasty: a practical treatment for sacral insufficiency fracture: case report. *Spine (Phila Pa 1976).* 2005;30(15):E450-4. PMID: 16094265

43. Butler CL, Given CA, 2nd, Michel SJ, et al. Percutaneous sacroplasty for the treatment of sacral insufficiency fractures. *AJR Am J Roentgenol*. 2005;184(6):1956-9. PMID: 15908561
44. Deen HG, Nottmeier EW. Balloon kyphoplasty for treatment of sacral insufficiency fractures. Report of three cases. *Neurosurg Focus*. 2005;18(3):e7. PMID: 15771397
45. Gjertsen O, Schellhorn T, Nakstad PH. Fluoroscopy-guided sacroplasty: special focus on preoperative planning from three-dimensional computed tomography. *Acta Radiol*. 2008;49(9):1042-8. PMID: 18759150
46. Heron J, Connell DA, James SL. CT-guided sacroplasty for the treatment of sacral insufficiency fractures. *Clin Radiol*. 2007;62(11):1094-100; discussion 101-3. PMID: 17920869
47. Pommersheim W, Huang-Hellinger F, Baker M, et al. Sacroplasty: a treatment for sacral insufficiency fractures. *AJNR Am J Neuroradiol*. 2003;24(5):1003-7. PMID: 12748113
48. Strub WM, Hoffmann M, Ernst RJ, et al. Sacroplasty by CT and fluoroscopic guidance: is the procedure right for your patient? *AJNR Am J Neuroradiol*. 2007;28(1):38-41. PMID: 17213421
49. Whitlow CT, Mussat-Whitlow BJ, Mattern CW, et al. Sacroplasty versus vertebroplasty: comparable clinical outcomes for the treatment of fracture-related pain. *AJNR Am J Neuroradiol*. 2007;28(7):1266-70. PMID: 17698526
50. Kamel EM, Binaghi S, Guntern D, et al. Outcome of long-axis percutaneous sacroplasty for the treatment of sacral insufficiency fractures. *Eur Radiol*. 2009;19(12):3002-7. PMID: 19533145
51. Talmadge J, Smith K, Dykes T, et al. Clinical impact of sacroplasty on patient mobility. *J Vasc Interv Radiol*. 2014;25(6):911-5. PMID: 24713417
52. Andresen R, Radmer S, Ludtke CW, et al. Balloon sacroplasty as a palliative pain treatment in patients with metastasis-induced bone destruction and pathological fractures. *RoFo : Fortschritte auf dem Gebiete der Rontgenstrahlen und der Nuklearmedizin*. 2014;186(9):881-6. PMID: 24557599
53. Zhao S, Xu CY, Zhu AR, et al. Comparison of the efficacy and safety of 3 treatments for patients with osteoporotic vertebral compression fractures: A network meta-analysis. *Medicine*. 2017;96(26):e7328. PMID: 28658144
54. Zaman FM, Frey M, Slipman CW. Sacral stress fractures. *Curr Sports Med Rep*. 2006;5(1):37-43. PMID: 16483515
55. Zhang H, Xu C, Zhang T, et al. Does Percutaneous Vertebroplasty or Balloon Kyphoplasty for Osteoporotic Vertebral Compression Fractures Increase the Incidence of New Vertebral Fractures? A Meta-Analysis. *Pain physician*. 2017;20(1):E13-E28. PMID: 28072794
56. Zhan Y, Jiang J, Liao H, et al. Risk factors for cement leakage after vertebroplasty or kyphoplasty: a meta-analysis of published evidence. *World neurosurgery*. 2017. PMID: 28192270
57. Xiao H, Yang J, Feng X, et al. Comparing complications of vertebroplasty and kyphoplasty for treating osteoporotic vertebral compression fractures: a meta-analysis of the randomized and non-randomized controlled studies. *European journal of orthopaedic surgery & traumatology : orthopedie traumatologie*. 2015;25 Suppl 1:S77-85. PMID: 24989933
58. Blasco Andaluz J, Martinez-Ferrer A, Macho Fernandez J, et al. Effect of vertebroplasty on pain relief, quality of life and the incidence of new vertebral fractures. A 12-month randomised follow-up, controlled trial. *J Bone Miner Res*. 2012. PMID: 22307990

59. Rousing R, Hansen KL, Andersen MO, et al. Twelve-months follow-up in forty-nine patients with acute/semiacute osteoporotic vertebral fractures treated conservatively or with percutaneous vertebroplasty: a clinical randomized study. *Spine (Phila Pa 1976)*. 2010;35:478-82. PMID: 20190623
60. Zhang YZ, Kong LD, Cao JM, et al. Incidence of subsequent vertebral body fractures after vertebroplasty. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia*. 2014;21(8):1292-7. PMID: 24973334
61. Spross C, Aghayev E, Kocher R, et al. Incidence and risk factors for early adjacent vertebral fractures after balloon kyphoplasty for osteoporotic fractures: analysis of the SWISSspine registry. *European spine journal : official publication of the European Spine Society, the European Spinal Deformity Society, and the European Section of the Cervical Spine Research Society*. 2014;23(6):1332-8. PMID: 24197481
62. Civelek E, Cansever T, Yilmaz C, et al. The retrospective analysis of the effect of balloon kyphoplasty to the adjacent-segment fracture in 171 patients. *Journal of spinal disorders & techniques*. 2014;27(2):98-104. PMID: 24795949
63. American Academy of Orthopaedic Surgeons Guideline and Evidence Report. 2010. Treatment of Symptomatic Osteoporotic Spinal Compression Fractures. [cited 6/10/2024]. 'Available from:'
64. Rousing R, Andersen MO, Jespersen SM, et al. Percutaneous vertebroplasty compared to conservative treatment in patients with painful acute or subacute osteoporotic vertebral fractures: three-months follow-up in a clinical randomized study. *Spine (Phila Pa 1976)*. 2009;34(13):1349-54. PMID: 19478654
65. Voormolen MH, Mali WP, Lohle PN, et al. Percutaneous vertebroplasty compared with optimal pain medication treatment: short-term clinical outcome of patients with subacute or chronic painful osteoporotic vertebral compression fractures. The VERTOS study. *AJNR Am J Neuroradiol*. 2007;28(3):555-60. PMID: 17353335
66. Diamond TH, Bryant C, Browne L, et al. Clinical outcomes after acute osteoporotic vertebral fractures: a 2-year non-randomised trial comparing percutaneous vertebroplasty with conservative therapy. *Med J Aust*. 2006;184(3):113-7. PMID: 16460295
67. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. 2010;112(4):810-33. PMID: 20124882
68. Barr JD, Jensen ME, Hirsch JA, et al. Position statement on percutaneous vertebral augmentation: a consensus statement developed by the Society of Interventional Radiology (SIR), American Association of Neurological Surgeons (AANS) and the Congress of Neurological Surgeons (CNS), American College of Radiology (ACR), American Society of Neuroradiology (ASNR), American Society of Spine Radiology (ASSR), Canadian Interventional Radiology Association (CIRA), and the Society of NeuroInterventional Surgery (SNIS). *J Vasc Interv Radiol*. 2014;25(2):171-81. PMID: 24325929
69. ACR–ASNR–ASSR–SIR–SNIS Practice guideline for the performance of vertebral augmentation. Revised 2017. [cited 6/10/2024]. 'Available from:' <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/vertebraaug.pdf>.
70. Chandra RV, Meyers PM, Hirsch JA, et al. Vertebral augmentation: report of the Standards and Guidelines Committee of the Society of NeuroInterventional Surgery. *J Neurointerv Surg*. 2014;6:7-15. PMID: 24198272

71. Baerlocher MO, Saad WE, Dariushnia S, et al. Quality improvement guidelines for percutaneous vertebroplasty. *J Vasc Interv Radiol.* 2014;25(2):165-70. PMID: 24238815
72. Feng L, Shen JM, Feng C, et al. Comparison of radiofrequency kyphoplasty (RFK) and balloon kyphoplasty (BKP) in the treatment of vertebral compression fractures: A meta-analysis. *Medicine.* 2017;96(25):e7150. PMID: 28640091
73. National Institute for Health and Clinical Excellence (NICE). Percutaneous vertebroplasty and percutaneous balloon kyphoplasty for treating osteoporotic vertebral compression fractures [TA279]. [cited. 'Available from:'] <https://www.nice.org.uk/guidance/ta279>.
74. National Institute for Health and Clinical Excellence (NICE). Metastatic spinal cord compression in adults: risk assessment, diagnosis and management [CG75]. [cited. 'Available from:'] <https://www.nice.org.uk/guidance/cg75/chapter/1-Guidance>.
75. Aman MM, Mahmoud A, Deer T, et al. The American Society of Pain and Neuroscience (ASPN) Best Practices and Guidelines for the Interventional Management of Cancer-Associated Pain. *J Pain Res.* 2021;14:2139-64. PMID: 34295184

CODES

Codes	Number	Description	
CPT	22510	Percutaneous vertebroplasty (bone biopsy included when performed), 1 vertebral body, unilateral or bilateral injection, inclusive of all imaging guidance; cervicothoracic	
	22511	;lumbosacral	
	22512	;each additional cervicothoracic or lumbosacral vertebral body (List separately in addition to code for primary procedure)	
	22513	Percutaneous vertebral augmentation, including cavity creation (fracture reduction and bone biopsy included when performed) using mechanical device (eg, kyphoplasty), 1 vertebral body, unilateral or bilateral cannulation, inclusive of all imaging guidance; thoracic	
	22514	;lumbar	
	22515	;each additional thoracic or lumbar vertebral body (List separately in addition to code for primary procedure)	
	0200T	Percutaneous sacral augmentation (sacroplasty), unilateral injection(s), including the use of a balloon or mechanical device, when used, 1 or more needles, includes imaging guidance and bone biopsy, when performed	
	0201T	Percutaneous sacral augmentation (sacroplasty), bilateral injection(s), including the use of a balloon or mechanical device, when used, 2 or more needles, includes imaging guidance and bone biopsy, when performed	
	HCPCS	C1062	Intravertebral body fracture augmentation with implant (e.g., metal, polymer)
		C7504	Percutaneous vertebroplasties (bone biopsies included when performed), first cervicothoracic and any additional cervicothoracic or lumbosacral vertebral bodies, unilateral or bilateral injection, inclusive of all imaging guidance
C7505		Percutaneous vertebroplasties (bone biopsies included when performed), first lumbosacral and any additional cervicothoracic or lumbosacral vertebral bodies, unilateral or bilateral injection, inclusive of all imaging guidance	
C7507		Percutaneous vertebral augmentations, first thoracic and any additional thoracic or lumbar vertebral bodies, including cavity creations (fracture reductions and bone biopsies included when performed) using mechanical device (eg, kyphoplasty), unilateral or bilateral cannulations, inclusive of all imaging guidance	

Codes	Number	Description
	C7508	Percutaneous vertebral augmentations, first lumbar and any additional thoracic or lumbar vertebral bodies, including cavity creations (fracture reductions and bone biopsies included when performed) using mechanical device (eg, kyphoplasty), unilateral or bilateral cannulations, inclusive of all imaging guidance

Date of Origin: July 2000